

CATEGORY $\mathcal{O}$ AND $\mathfrak{sl}_k$ LINK INVARIANTS

JOSHUA SUSSAN

CONTENTS

1. Introduction	1
2. Representation Theory of $\mathfrak{sl}_k$	3
3. Categorification of $\mathfrak{sl}_k$ Modules	7
4. Diagram Relations	14
5. Graded Category $\mathcal{O}$	39
6. Crossings	61
References	74

1. INTRODUCTION

The program of categorification via category $\mathcal{O}$ was introduced by J. Bernstein, I. Frenkel, and M. Khovanov in [BFK] where tensor powers of the standard two dimensional representation of $\mathfrak{sl}_2$ were recognized as Grothendieck groups of certain subcategories of $\mathcal{O}$ for various $\mathfrak{gl}_n$. They had two different constructions. One was based on studying certain blocks with singular generalized central characters. The other was based on examining the trivial regular block but by considering various parabolic subcategories. In the first case the action of $\mathfrak{sl}_2$ was categorified by projective functors acting on these singular categories. The intertwiners were categorified by derived Zuckerman functors acting on the derived category $\mathcal{O}$. In the latter case, the action of the Lie algebra was lifted to Zuckerman functors and the intertwiners became projective functors. These two constructions are related by Koszul duality where the projective functors get exchanged for Zuckerman functors and visa-versa [Rh], [MOS].

Several conjectures posed in [BFK] were solved in [Str2]. The most important result of that work was that the Jones polynomial was lifted to a functorial invariant of tangles. Natural transformations between these functors became invariants of 2-tangles acting as cobordisms between two different tangles [Str4]. One of the principle goals of categorification is to obtain invariants for $n + 1$ dimensional topological objects by homological realizations of classical n dimensional invariants.

There is a parallel approach to categorification of link invariants due to Khovanov. In [Khov], a bi-graded homology theory was constructed whose graded Euler characteristic is the Jones polynomial. M. Khovanov and L. Rozansky extended this work to a homology theory categorifying the HOMFLYPT polynomial [KR]. The theory of matrix factorization is their primary tool. H. Murakami, T. Ohtsuki, and S. Yamada have an interpretation of the HOMFLYPT polynomial through a polynomial assigned to certain planar graphs [MOY]. Khovanov and Rozansky categorified these polynomials which is a crucial step in the categorification of the HOMFLYPT polynomial.

The goal of this paper is to categorify the $\mathfrak{sl}_k$ tangle invariant via category $\mathcal{O}$. It is important to understand all of the planar graphs representation theoretically. The planar graphs give a graphical interpretation of intertwiners between various tensor products $\Lambda^{i_1} V_{k-1} \otimes \cdots \otimes \Lambda^{i_r} V_{k-1}$ of fundamental $\mathfrak{sl}_k$ representations. The first step is to categorify tensor powers of the standard k - dimensional representation of $\mathfrak{sl}_k$. In order to accomplish this, we look at more general singular blocks of category $\oplus_{\mathbf{d}} \mathcal{O}_{\mathbf{d}}(\mathfrak{gl}_n)$. Here, $\mathbf{d}$ denotes a k - tuple

$(d_{k-1}, \dots, d_0)$. Then $\mathcal{O}_{\mathbf{d}}(\mathfrak{gl}_n)$ is a block corresponding to an integral dominant weight whose stabilizer under the dot action of the Weyl group is $S_{d_{k-1}} \times \dots \times S_{d_0}$. Parabolic subcategories of these blocks, $\mathcal{O}_{\mathbf{d}}^{\mathbf{p}}(\mathfrak{gl}_n)$ will provide a categorification of tensor products of fundamental representations. The action of the Lie algebra $\mathfrak{sl}_k$ will be lifted to projective functors. These are functors given by tensoring with a finite dimensional representation and then projecting onto a certain block. Morphisms between various $\Lambda^{i_1} V_{k-1} \otimes \dots \otimes \Lambda^{i_r} V_{k-1}$ will become inclusion and derived Zuckerman functors between various parabolic subcategories of these blocks. Relations between the intertwiners become functorial isomorphisms between various compositions of inclusion and derived Zuckerman functors. It should be possible to do this construction in the Koszul dual situation as well. In this case projective functors categorify the intertwiners and derived Zuckerman functors categorify the action of the Lie algebra.

This construction only gives rise to a categorification of $\mathfrak{sl}_k$ -modules. In order to get a categorification of the quantum group, we consider the above categories to be a category of graded modules. Then in the Grothendieck group the shift functor will descend to the action of multiplication by the quantum parameter q . The theory of graded category $\mathcal{O}$ is due to W. Soergel and has been studied extensively by C. Stroppel [Str1]. Projective functors have graded lifts. This allows us to give a categorification of the quantum Serre relations. Zuckerman functors and inclusion functors also have graded lifts. Graded functors may now be assigned to all flat tangles. In order to extend this construction to tangle with crossings, we consider adjunction morphisms between shifted compositions of inclusion and Zuckerman functors for a minimal parabolic and the identity functor. Cones of these morphisms are assigned to the crossings. In this graded setup it is easy to see that these cones satisfy the $\mathfrak{sl}_k$ skein relation. A cobordism between two tangles should give rise to a natural transformation between the functors assigned to the tangles. This natural transformation should become an invariant of 2-tangles as in [Str4], [Khov3], [Jac].

In section 2 we review the necessary finite dimensional representation theory of $\mathfrak{sl}_k$. A categorification of the fundamental representations will be given in section 3. An equivalence of categories between certain parabolic subcategories which generalizes corollary 5 in [BFK] is also given in this section. This equivalence is used in the definition of the functors assigned to the flat tangles. In section 4 we provide functorial analogues of the graphic relations presented in section 2. Each of the graphical relations will give rise to a functorial isomorphism. Natural transformations are constructed and shown that when restricted to generalized Verma modules, they are isomorphisms. Most of the section is devoted to calculations of the relevant functors on generalized Verma modules. In section 5 graded category $\mathcal{O}$ will be discussed and all of the results in the previous sections will be lifted to the graded case. We define in section 6 our functor valued tangle invariant $\mathbb{F}$ and prove that it satisfies the Reidemeister relations.

Suppose that tangles T and T' are morphisms from r points to r' points labeled from the set $\{1, k-1\}$. Denote by n and n' the sum of the labels for the r and r' points respectively. These compositions of n and n' naturally give rise to parabolic subalgebras $\mathfrak{p}$ and $\mathfrak{p}'$ of $\mathfrak{gl}_n$ and $\mathfrak{gl}_{n'}$. Theorem 7 is the main result of this work.

Theorem. *If tangles T and T' are ambient isotopic tangles, then $\mathbb{F}(T)$ and $\mathbb{F}(T')$ are isomorphic as derived functors from $D^b(\oplus_{\mathbf{d}} \mathcal{O}_{\mathbf{d}}^{\mathbf{p}}(\mathfrak{gl}_n))$ to $D^b(\oplus_{\mathbf{d}'} \mathcal{O}_{\mathbf{d}'}^{\mathbf{p}'}(\mathfrak{gl}_{n'}))$.*

The functorial isomorphisms of section 4 play a crucial role in proving this theorem. When restricted to $(0,0)$ -tangles, the invariant is a complex of graded vector spaces and thus homology groups may be assigned to links.

It follows immediately from the definition of $\mathbb{F}$ that on the Grothendieck group, the following equality holds:

Theorem. *Let Ω_i and Π_i be the functors assigned to the crossings. Then $q^k[\Omega_i] - q^{-k}[\Pi_i] = (-1)^{k-1}(q^1 - q^{-1})[\text{Id}]$*

An explicit relationship between the functorial tangle invariant introduced by Khovanov in [Khov5] and the category $\mathcal{O}$ invariant of [Str2] was given in [Str5]. Stroppel considered the subcategory of the trivial block of $\mathcal{O}(\mathfrak{sl}_{2n})$ of modules locally finite with respect to a parabolic subalgebra whose reductive part is $\mathfrak{gl}_n \oplus \mathfrak{gl}_n$ and whose projective presentation only consists of projective-injective modules. This subcategory categorifies the space of invariants in tensor products of the standard two dimensional of $\mathfrak{sl}_2$. This subcategory is equivalent to a category of modules over an associative algebra which is isomorphic to the algebra constructed in

[Khov5]. A natural question is to find a connection between the functorial $\mathfrak{sl}_k$ invariant in this work and the invariant constructed via matrix factorization in [KR] or Soergel bimodules in [Khov4]. One should probably look for a subcategory of $\mathcal{O}$ that categorifies the space of invariants in tensor products of the standard k -dimensional representation of $\mathfrak{sl}_k$ and then study the associative algebra governing that subcategory.

Acknowledgements: The author is grateful to his advisor, Igor Frenkel for his encouragement and support throughout the development of this project. In addition, the author is very thankful to Mikhail Khovanov, Raphael Rouquier, Catharina Stroppel, and Gregg Zuckerman for helpful conversations and comments on preliminary versions of this work.

2. REPRESENTATION THEORY OF $\mathfrak{sl}_k$

2.1. Basic Definitions. We begin by reviewing the finite dimensional representation theory of $\mathfrak{sl}_k$. Recall that $\mathfrak{sl}_k$ is the Lie algebra of $k \times k$ matrices with entries in $\mathbb{C}$. Denote the matrix with only 1 in the (i, j) entry by $e_{i,j}$. There is the triangular decomposition $\mathfrak{sl}_k = \mathfrak{n}^- + \mathfrak{h} + \mathfrak{n}^+$ where $\mathfrak{n}^-$ is the subalgebra of lower triangular matrices, $\mathfrak{n}^+$ is the subalgebra of upper triangular matrices and $\mathfrak{h}$ is the Cartan subalgebra of diagonal matrices.

The dual $\mathfrak{h}^*$ of the Cartan subalgebra has a basis $\alpha_1 = e_{1,1}^* - e_{2,2}^*, \dots, \alpha_i = e_{i,i}^* - e_{i+1,i+1}^*, \dots, \alpha_{k-1} = e_{k-1,k-1}^* - e_{k,k}^*$ where $e_{i,j}^*(e_{r,s}) = \delta_{i,r}\delta_{j,s}$. These α_i are called the positive simple roots for $\mathfrak{sl}_k$.

A linear map $\lambda: \mathfrak{h} \rightarrow \mathbb{C}$ may be written in coordinates $\lambda = \lambda_1 e_{1,1}^* + \dots + \lambda_k e_{k,k}^*$.

The finite dimensional irreducible representations of $\mathfrak{sl}_k$ are indexed by these weights λ . There exists a unique irreducible representation of highest weight $\lambda = \lambda_1 e_{1,1}^* + \dots + \lambda_k e_{k,k}^*$ when $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$. For our purposes, one of the most important representations is the standard k -dimensional vector space $\mathbb{C}^k$ where $\mathfrak{sl}_k$ acts on it naturally as matrices. Call this space V_{k-1} . The basis for this vector space is given by $e_1, \dots, e_k$ and $e_{ij}(e_m) = \delta_{j,m}e_i$. Notice that the highest weight of this representation is $e_{1,1}^*$. We denote the i th exterior power of a module V by $\Lambda^i V$.

Proposition 1. *Let V_{k-1} be the module defined above. Then $\Lambda^i V_{k-1}$ is an irreducible module of highest weight $e_{1,1}^* + \dots + e_{i,i}^*$.*

Proof. See [FH] page 221. □

Now we define the algebra $\mathcal{U}_q(\mathfrak{sl}_k)$ and its fundamental representations.

Definition 1. The quantum group $\mathcal{U}_q(\mathfrak{sl}_k)$ is the associative algebra over $\mathbb{C}(q)$ with generators E_i, F_i, K_i, K_i^{-1} for $i = 1, \dots, k-1$ satisfying the following conditions:

- (1) $K_i K_i^{-1} = K_i^{-1} K_i = 1$
- (2) $K_i K_j = K_j K_i$
- (3) $K_i E_j = q^{c_{i,j}} E_j K_i$
- (4) $K_i F_j = q^{-c_{i,j}} F_j K_i$
- (5) $E_i F_j - F_j E_i = \delta_{i,j} \frac{K_i - K_i^{-1}}{q - q^{-1}}$
- (6) $E_i E_j = E_j E_i$ if $|i - j| > 1$
- (7) $F_i F_j = F_j F_i$ if $|i - j| > 1$
- (8) $E_i^2 E_{i\pm 1} - (q + q^{-1}) E_i E_{i\pm 1} E_i + E_{i\pm 1} E_i^2 = 0$
- (9) $F_i^2 F_{i\pm 1} - (q + q^{-1}) F_i F_{i\pm 1} F_i + F_{i\pm 1} F_i^2 = 0$

where

$$c_{i,j} = \begin{cases} 2 & \text{if } j = i \\ -1 & \text{if } j = i \pm 1 \\ 0 & \text{if } |i - j| > 1. \end{cases}$$

The most basic representation of $\mathcal{U}(\mathfrak{sl}_k)$ is V_{k-1} . It is the k -dimensional vector space with basis $v_0, \dots, v_{k-1}$. The algebra acts on this space as follows:

$$\begin{aligned} E_i v_j &= 0 & \text{if } j \neq i-1 \\ E_i v_j &= v_i & \text{if } j = i-1 \\ F_i v_j &= 0 & \text{if } j \neq i \\ F_i v_j &= v_{i-1} & \text{if } j = i \\ K_i^{\pm 1} v_j &= q^{\pm 1} v_i & \text{if } j = i \\ K_i^{\pm 1} v_j &= q^{\mp 1} v_{i-1} & \text{if } j = i-1 \\ K_i^{\pm 1} v_j &= v_j & \text{if } j \neq i-1, i. \end{aligned}$$

There are several intertwiners between various representations that will be important for later. There is the map $\Lambda^k V_{k-1} \rightarrow V_{k-1} \otimes \Lambda^{k-1} V_{k-1}$ given by

$$v_{k-1} \wedge \dots \wedge v_0 \rightarrow \sum_{j=0}^{k-1} (-1)^j q^{k-j-1} v_j \otimes (v_{k-1} \wedge \dots \wedge \hat{v}_j \wedge \dots \wedge v_0),$$

where $\hat{v}_j$ means that the term is omitted from the expression.

There is the map $\Lambda^k V_{k-1} \rightarrow \Lambda^{k-1} V_{k-1} \otimes V_{k-1}$ given by

$$v_{k-1} \wedge \dots \wedge v_0 \rightarrow \sum_{j=0}^{k-1} (-1)^{k-1-j} q^j (v_{k-1} \wedge \dots \wedge \hat{v}_j \wedge \dots \wedge v_0 \otimes v_j).$$

There is a map in the other direction: $V_{k-1} \otimes \Lambda^{k-1} V_{k-1} \rightarrow \Lambda^k V_{k-1}$ given by

$$v_j \otimes (v_{k-1} \wedge \dots \wedge \hat{v}_j \wedge \dots \wedge v_0) \rightarrow (-1)^{k-j-1} q^{-j} v_{k-1} \wedge \dots \wedge v_0.$$

There is also the map $\Lambda^{k-1} V_{k-1} \otimes V_{k-1} \rightarrow \Lambda^k V_{k-1}$ given by

$$(v_{k-1} \wedge \dots \wedge \hat{v}_j \wedge \dots \wedge v_0) \otimes v_j \rightarrow (-1)^j q^{j-k+1} v_{k-1} \wedge \dots \wedge v_0.$$

The inclusion map $\Lambda^2 V_{k-1} \rightarrow V_{k-1} \otimes V_{k-1}$ is determined by

$$v_i \wedge v_j \rightarrow v_i \otimes v_j - q^1 v_j \otimes v_i$$

where $i > j$.

The projection map $V_{k-1} \otimes V_{k-1} \rightarrow \Lambda^2 V_{k-1}$ is given by

$$\begin{aligned} v_i \otimes v_j &\rightarrow q^{-1} v_i \wedge v_j & \text{if } i > j \\ &\rightarrow -v_i \wedge v_j & \text{if } i < j. \end{aligned}$$

More generally, let

$$\pi_{r_1} \otimes \dots \otimes \pi_{r_t} : V_{k-1}^{\otimes(r_1+\dots+r_t)} \rightarrow \Lambda^{r_1} V_{k-1} \otimes \dots \otimes \Lambda^{r_t} V_{k-1}$$

be the canonical projection map and

$$i_{r_1} \otimes \dots \otimes i_{r_t} : \Lambda^{r_1} V_{k-1} \otimes \dots \otimes \Lambda^{r_t} V_{k-1} \rightarrow V_{k-1}^{\otimes(r_1+\dots+r_t)}$$

be the canonical inclusion map. Since we will not need these more general intertwiners, we will omit the precise formulas.

2.2. Graphical Calculus. There is a graphical description of the intertwiners between tensor products of fundamental representations via colored trivalent graphs. A line segment labeled by an integer i where $1 \leq i \leq k$, will depict the representation $\Lambda^i V_{k-1}$. Note that an edge labeled by k depicts the trivial representation and thus such an edge may be added or removed at will. The figure below depicts the canonical projection map $\Lambda^i V_{k-1} \otimes \Lambda^j V_{k-1} \rightarrow \Lambda^{i+j} V_{k-1}$.

The inclusion $\Lambda^{i+j} V_{k-1} \rightarrow \Lambda^i V_{k-1} \otimes \Lambda^j V_{k-1}$ is given by the diagram below.

There are relations between these maps. Graphically, the relations are depicted by the following five diagrams.




FIGURE 1.




FIGURE 2.




FIGURE 3. Diagram Relation 1




FIGURE 4. Diagram Relation 2

If the trivalent graph has external edges labeled only by k , then it represents a composition of intertwiners from the the trivial representation to itself so a Laurent polynomial may be assigned to it. We will say that such a graph is closed.

Theorem 1. *There is a Laurent polynomial $\langle D \rangle_k$ in $\mathbb{Z}[q, q^{-1}]$ which may be assigned to closed, colored trivalent graphs D which satisfy the five relations above. It is invariant under ambient isotopy of $\mathbb{R}^2$.*

Proof. See sections 1 and 2 of [MOY]. □

The Laurent polynomial in the theorem above coincides with the representation-theoretic polynomial.

A version of the HOMFLYPT for links may then be defined. Each crossing may be resolved in the following two ways:

Let D_+, D_-, D_0 be identical link diagrams except near a crossing as given by the figure below.

Definition 2. Let D be a planar projection of a link. Let

- (1) $\langle D_+ \rangle_k = q^1 \langle D_0 \rangle_k - \langle D_* \rangle_k$,
- (2) $\langle D_- \rangle_k = q^{-1} \langle D_0 \rangle_k - \langle D_* \rangle_k$,




FIGURE 5. Diagram Relation 3




FIGURE 6. Diagram Relation 4




FIGURE 7. Diagram Relation 5




FIGURE 8.

(3) $P_k(D) = q^{(k)(-w(D))} \langle D \rangle_k$ where $w(D)$ is the difference between the number of positive and negative crossings.

Theorem 2. (1) $P_k(D)$ satisfies the Reidemeister moves and thus is an invariant of a link L which has planar projection D .

(2) The invariant above satisfies the skein relation for the one variable $\mathfrak{sl}_k$ specialization of the HOM-FLYPT polynomial:

$$q^k P_k(D_+) - q^{-k} P_k(D_-) = (q^1 - q^{-1}) P_k(D_0).$$




FIGURE 9.

Proof. This is theorem 3.2 of [MOY]. □

3. CATEGORIFICATION OF $\mathfrak{sl}_k$ MODULES

The most important representations to categorify are the tensor powers $V_{k-1}^{\otimes n}$. When $k = 2$, the categorifications were constructed in [BFK]. The authors worked on the most singular blocks of category $\mathcal{O}$ which correspond to maximal subgroups of the symmetric group. They also suggested what to do for the $\mathfrak{sl}_k$ case: consider other blocks of category $\mathcal{O}$.

3.1. Categorification of $\Lambda^{i_1} V_{k-1} \otimes \cdots \otimes \Lambda^{i_r} V_{k-1}$.

Definition 3. (1) Denote the Grothendieck group of an abelian or triangulated category $\mathcal{C}$ by $[\mathcal{C}]$. It is the free abelian group generated by the symbols $[M]$ where M is an object of $\mathcal{C}$. The only relations in this group are of the form $[N] = [M] + [P]$ when there is a short exact sequence or distinguished triangle of the form

$$0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0.$$

- (2) The image of an object M or an exact functor F in the Grothendieck group will be indicated by $[M]$ and $[F]$ respectively.
- (3) Given an abelian category $\mathcal{C}$, let $D^b(\mathcal{C})$ denote the corresponding bounded derived category.
- (4) Denote by $H^j: D^b(\mathcal{C}) \rightarrow \mathcal{C}$, the j th cohomology functor. If F is a left exact functor, the j th derived functor $R^j F$ is defined to be $H^j \circ RF$. If F is right exact, the j th derived functor $L_j F$ is defined to be $H^{-j} \circ LF$.

The truncation functors $\tau^{\leq n}$ and $\tau^{\geq n}$ are well defined on the derived category.

Proposition 2. *Let X be an object in $D^b(\mathcal{C})$. The following triangles exist in the derived category:*

$$\begin{aligned} \tau^{\leq n} X &\rightarrow X \rightarrow \tau^{\geq n+1} X \\ \tau^{\leq n-1} X &\rightarrow \tau^{\leq n} X \rightarrow H^n(X)[-n] \end{aligned}$$

Proof. See proposition 4.1.8 of [Sch]. □

Definition 4. Let $\mathcal{O}(\mathfrak{gl}_n)$ be the category of $\mathfrak{gl}_n$ modules which satisfy the following properties:

- (1) Finitely generated as $\mathcal{U}(\mathfrak{gl}_n)$ -modules.
- (2) Diagonalizable under the action of the Cartan subalgebra $\mathfrak{h}$.
- (3) Locally finite under the action of the Borel subalgebra $\mathfrak{b} = \mathfrak{h} + \mathfrak{n}^+$.

This category decomposes into a direct sum of subcategories corresponding to the generalized central characters.

Definition 5. (1) Let $\mathcal{O}_{(d_{k-1}, d_{k-2}, \dots, d_0)} = \mathcal{O}_{\mathbf{d}}$ be the block of $\mathcal{O}(\mathfrak{gl}_n)$ for the central character corresponding to the weight $\sum_{i=0}^{k-1} \sum_{j=1}^{d_i} i e_{d_0 + \dots + d_{i-1} + j} - \rho$, where

$$\rho = \frac{n-1}{2} e_1 + \frac{n-3}{2} e_2 + \cdots + \frac{1-n}{2} e_n$$

is half the sum of the positive roots.

- (2) Let $M(a_1, \dots, a_n)$ be the Verma module with highest weight $a_1 e_1 + \cdots + a_n e_n - \rho$.
- (3) Let $L(a_1, \dots, a_n)$ be the Verma module with highest weight $a_1 e_1 + \cdots + a_n e_n - \rho$.

There are d_i terms in the weight from the definition above with coefficient i . Note that $\sum_{j=0}^{k-1} d_j = n$.

Proposition 3. *Assume that the following direct sum is over all $\mathbf{d}$ such that the entries are non-negative integers and the sum of the entries is n . Then $\mathbb{C} \otimes_{\mathbb{Z}} [\oplus \mathcal{O}_{\mathbf{d}}] \cong V_{k-1}^{\otimes n}$.*

Proof. The image of the Verma module $[M(a, \dots, a_n)]$ gets mapped to $v_{a_1} \otimes \dots \otimes v_{a_n}$. $\square$

This proposition is the first step towards categorification of $\mathfrak{sl}_k$ -modules. Next we would like to categorify the action of the Lie algebra. The desired functors come directly from [BFK]. It is essentially the projective functor of tensoring with the n -dimensional representation V_{n-1} . One only has to be careful about projecting onto the various blocks. This is done next.

Definition 6. (1) Let $\mathcal{E}_i: \mathcal{O}_{(d_{k-1}, d_{k-2}, \dots, d_0)} \rightarrow \mathcal{O}_{(d_{k-1}, \dots, d_i+1, d_{i-1}-1, \dots, d_0)}$ be the functor defined by

$$\mathcal{E}_i M = \text{proj}_{(d_{k-1}, \dots, d_i+1, d_{i-1}-1, \dots, d_0)}(V_{n-1} \otimes M).$$

(2) Let $\mathcal{F}_i: \mathcal{O}_{(d_{k-1}, d_{k-2}, \dots, d_0)} \rightarrow \mathcal{O}_{(d_{k-1}, \dots, d_i-1, d_{i-1}+1, \dots, d_0)}$ by

$$\mathcal{F}_i M = \text{proj}_{(d_{k-1}, \dots, d_i-1, d_{i-1}+1, \dots, d_0)}(V_{n-1}^* \otimes M).$$

(3) Let $\mathcal{H}_i: \mathcal{O}_{(d_{k-1}, d_{k-2}, \dots, d_0)} \rightarrow \mathcal{O}_{(d_{k-1}, d_{k-2}, \dots, d_0)}$ be $\text{Id}^{\oplus(d_i-d_{i-1})}$.

Theorem 3. *Functorial Serre relations are satisfied on $\oplus_{\mathbf{d}} \mathcal{O}(\mathfrak{gl}_n)$.*

- (1) $\mathcal{H}_i \mathcal{H}_j \cong \mathcal{H}_j \mathcal{H}_i$.
- (2) If $d_i > d_{i-1}$ then $\mathcal{E}_i \mathcal{F}_i \cong \mathcal{F}_i \mathcal{E}_i \oplus \mathcal{H}_i$.
- (3) If $d_i = d_{i-1}$ then $\mathcal{E}_i \mathcal{F}_i \cong \mathcal{F}_i \mathcal{E}_i$.
- (4) If $d_i < d_{i-1}$ then $\mathcal{F}_i \mathcal{E}_i \cong \mathcal{E}_i \mathcal{F}_i \oplus \mathcal{H}_i$.
- (5) If $i \neq j$, then $\mathcal{E}_i \mathcal{F}_j \cong \mathcal{F}_j \mathcal{E}_i$.
- (6) If $d_i - d_{i-1} \geq 0$, then $\mathcal{H}_i \mathcal{E}_i \cong \mathcal{E}_i \mathcal{H}_i \oplus \mathcal{E}_i^{\oplus 2}$.
- (7) If $d_i - d_{i-1} \leq -2$, then $\mathcal{E}_i \mathcal{H}_i \cong \mathcal{H}_i \mathcal{E}_i \oplus \mathcal{E}_i^{\oplus 2}$.
- (8) If $d_i - d_{i-1} = -1$, then $\mathcal{H}_i \mathcal{E}_i \cong \mathcal{E}_i \mathcal{H}_i$.
- (9) If $d_i - d_{i-1} \geq 2$, then $\mathcal{F}_i \mathcal{H}_i \cong \mathcal{H}_i \mathcal{F}_i \oplus \mathcal{F}_i^{\oplus 2}$.
- (10) If $d_i - d_{i-1} \leq 0$, then $\mathcal{H}_i \mathcal{F}_i \cong \mathcal{F}_i \mathcal{H}_i \oplus \mathcal{F}_i^{\oplus 2}$.
- (11) If $d_i - d_{i-1} = 1$, then $\mathcal{H}_i \mathcal{F}_i \cong \mathcal{F}_i \mathcal{H}_i$.
- (12) If $|i - j| > 1$, then $\mathcal{H}_i \mathcal{E}_j \cong \mathcal{E}_j \mathcal{H}_i$.
- (13) If $|i - j| > 1$, then $\mathcal{H}_i \mathcal{F}_j \cong \mathcal{F}_j \mathcal{H}_i$.
- (14) If $|i - j| > 1$, then $\mathcal{E}_i \mathcal{E}_j \cong \mathcal{E}_j \mathcal{E}_i$.
- (15) If $|i - j| > 1$, then $\mathcal{F}_i \mathcal{F}_j \cong \mathcal{F}_j \mathcal{F}_i$.
- (16) If $j = i + 1$, then $\mathcal{E}_i \mathcal{E}_i \mathcal{E}_j \oplus \mathcal{E}_j \mathcal{E}_i \mathcal{E}_i \cong \mathcal{E}_i \mathcal{E}_j \mathcal{E}_i \oplus \mathcal{E}_i \mathcal{E}_j \mathcal{E}_i$.
- (17) If $j = i + 1$, then $\mathcal{F}_i \mathcal{F}_i \mathcal{F}_j \oplus \mathcal{F}_j \mathcal{F}_i \mathcal{F}_i \cong \mathcal{F}_i \mathcal{F}_j \mathcal{F}_i \oplus \mathcal{F}_i \mathcal{F}_j \mathcal{F}_i$.
- (18) If $d_i - d_{i-1} - 1 \geq 0$, then $\mathcal{H}_i \mathcal{E}_{i+1} \oplus \mathcal{E}_{i+1} \cong \mathcal{E}_{i+1} \mathcal{H}_i$.
- (19) If $d_i - d_{i-1} \leq 0$, then $\mathcal{E}_{i+1} \mathcal{H}_i \oplus \mathcal{E}_{i+1} \cong \mathcal{H}_i \mathcal{E}_{i+1}$.
- (20) If $d_i - d_{i-1} - 1 \geq 0$, then $\mathcal{H}_i \mathcal{E}_{i-1} \oplus \mathcal{E}_{i-1} \cong \mathcal{E}_{i-1} \mathcal{H}_i$.
- (21) If $d_i - d_{i-1} \leq 0$, then $\mathcal{E}_{i-1} \mathcal{H}_i \oplus \mathcal{E}_{i-1} \cong \mathcal{H}_i \mathcal{E}_{i-1}$.
- (22) If $d_i - d_{i-1} \geq 0$, then $\mathcal{F}_{i+1} \mathcal{H}_i \oplus \mathcal{F}_{i+1} \cong \mathcal{H}_i \mathcal{F}_{i+1}$.
- (23) If $d_i - d_{i-1} + 1 \geq 0$, then $\mathcal{H}_i \mathcal{F}_{i+1} \oplus \mathcal{F}_{i+1} \cong \mathcal{F}_{i+1} \mathcal{H}_i$.
- (24) If $d_i - d_{i-1} \geq 0$, then $\mathcal{F}_{i-1} \mathcal{H}_i \oplus \mathcal{F}_{i-1} \cong \mathcal{H}_i \mathcal{F}_{i-1}$.
- (25) If $d_i - d_{i-1} + 1 \geq 0$, then $\mathcal{H}_i \mathcal{F}_{i-1} \oplus \mathcal{F}_{i-1} \cong \mathcal{F}_{i-1} \mathcal{H}_i$.

One only has to compute in the Grothendieck group to check these relations [BG]. Details will be provided later when a functorial version of the quantum Serre relations is proved.

Our next goal is to categorify tensor products of exterior powers of V_{k-1} . The case $k = 2$ was addressed in [BFK]. When $k = 2$, $\Lambda^2 V_{k-1}$ is just a one dimensional vector space. The main ingredient in the categorification of $V_1^{\otimes n} \rightarrow V_1^{\otimes(n-2)}$ is the Zuckerman functor and parabolic subcategories. Thus for general k , we expect a locally finite subcategory to categorify a tensor product of fundamental representations. We now give a definition of various parabolic subalgebras of $\mathfrak{sl}_k$ and the corresponding locally finite subcategories first introduced by A. Rocha-Caridi in [Ro].

Definition 7. (1) The subalgebra $\mathfrak{p}_{(r_1, \dots, r_t)}$ is the parabolic subalgebra whose reductive subalgebra is $\mathfrak{gl}_{r_1} \oplus \dots \oplus \mathfrak{gl}_{r_t}$, where $r_1 + \dots + r_t = n$.

(2) Denote by $\mathcal{O}_{\mathbf{d}}^{\mathfrak{p}}$ the full subcategory of $\mathcal{O}_{\mathbf{d}}$ of modules locally finite with respect to the subalgebra $\mathfrak{p}$. By abuse of notation, let $\mathcal{O}_{\mathbf{d}}^{(r_1, \dots, r_t)}$ be the category $\mathcal{O}_{\mathbf{d}}^{\mathfrak{p}_{(r_1, \dots, r_t)}}$.

It is important to know some objects which are in these locally finite categories.

Definition 8. (1) Let S denote the subset of simple roots defining the parabolic subalgebra $\mathfrak{p}$.
(2) Let $P_{\mathfrak{p}}^+ = \{\lambda \in \mathfrak{h}^* | \langle \lambda, \alpha \rangle \in \mathbb{N}, \forall \alpha \in S\}$.

Given such a $\lambda \in P_{\mathfrak{p}}^+$, we may define the generalized Verma module $M^{\mathfrak{p}}(\lambda) = \mathcal{U}(\mathfrak{g}) \otimes_{\mathcal{U}(\mathfrak{p})} E(\lambda)$, where $E(\lambda)$ is the simple $\mathfrak{p}$ -module with highest weight λ .

Now it is easy to give conditions for which modules in these singular blocks are locally finite. Let $\lambda = a_1 e_1^* + \dots + a_n e_n^*$. We need to give a condition for λ to be in this set. Since $\langle \rho, h_i \rangle = 1$, the condition that $\langle \lambda - \rho, h_i \rangle \geq 0$ simply becomes $a_i > a_{i+1}$. We now could state which simple modules and generalized Verma modules are in the locally finite subcategories.

Lemma 1. (1) $L(a_1, \dots, a_n)$ is a simple module in $\mathcal{O}_{\mathfrak{d}}^{\mathfrak{p}}$ if $a_i > a_{i+1}$ whenever α_i is a simple root of $\mathfrak{p}$.
(2) $M^{\mathfrak{p}}(a_1, \dots, a_n)$ is a generalized Verma module in $\mathcal{O}_{\mathfrak{d}}^{\mathfrak{p}}$ if $a_i > a_{i+1}$ whenever α_i is a simple root of $\mathfrak{p}$.

Proof. This follows directly from above by evaluating at all simple roots α_i which define the parabolic subalgebra $\mathfrak{p}$. $\square$

We will often group coefficients in the highest weight of a generalized module such as

$$M^{\mathfrak{p}}(a_1, \dots, a_{i-1}, \underbrace{a_i, \dots, a_r}_{\text{grouped}}, \dots, a_n)$$

to stress that $a_i > \dots > a_r$ so that it is locally finite with respect to a certain subalgebra $\mathfrak{p}$.

Proposition 4. $\mathbb{C} \otimes_{\mathbb{Z}} [\oplus_{\mathfrak{d}} \mathcal{O}_{\mathfrak{d}}^{(r_1, \dots, r_t)}] \cong \Lambda^{r_1} V_{k-1} \otimes \dots \otimes \Lambda^{r_t} V_{k-1}$.

Proof. It suffices to show that this is an isomorphism on the basis of generalized Verma modules. Let $\mathfrak{p}$ be the subalgebra given above. The isomorphism sends $[M^{\mathfrak{p}}(a_1, \dots, a_n)]$ to

$$(v_{a_1} \wedge \dots \wedge v_{r_1}) \otimes \dots \otimes (v_{a_{r_1+\dots+r_{t-1}+1}} \wedge \dots \wedge v_{a_{r_1+\dots+r_t}}).$$

This is clearly a bijection. $\square$

Remark 1. If any of the r_i above is larger than k , then the category contains no non-trivial objects.

The Zuckerman functor plays an obvious role in this setup. It categorifies projection maps of these modules.

Definition 9. (1) Let the Zuckerman functor $\Gamma^{(r_1, \dots, r_t)}: \mathcal{O}_{\mathfrak{d}} \rightarrow \mathcal{O}_{\mathfrak{d}}^{(r_1, \dots, r_t)}$ be the functor of taking the maximal locally $\mathfrak{p}_{(r_1, \dots, r_t)}$ finite submodule.
(2) Let the dual Zuckerman $Z^{(r_1, \dots, r_t)}: \mathcal{O}_{\mathfrak{d}} \rightarrow \mathcal{O}_{\mathfrak{d}}^{(r_1, \dots, r_t)}$ be the functor of taking the maximal locally $\mathfrak{p}_{(r_1, \dots, r_t)}$ finite quotient.
(3) Let $\epsilon_{(r_1, \dots, r_t)}: \mathcal{O}_{\mathfrak{d}}^{(r_1, \dots, r_t)} \rightarrow \mathcal{O}_{\mathfrak{d}}$ be the natural inclusion functor.

Remark 2. The dual Zuckerman functor is also known as the Bernstein functor. See [KV].

If there is an inclusion of parabolic subalgebras $\mathfrak{q} \subset \mathfrak{p}$, there is an obvious generalization of these definitions for categories locally finite with respect to these algebras. For example, one may take a module locally finite with respect to $\mathfrak{p}$ and apply the Zuckerman functor $\Gamma_{\mathfrak{q}}^{\mathfrak{p}}$.

The Zuckerman functor is left exact. One usually studies its right derived functor and its cohomology functors. Taking the derived functor is important in categorification so that it becomes exact as a functor on the derived category. We denote its right derived functor shifted by j by $R\Gamma[j]$. The dual Zuckerman functor is right exact and similarly, one should consider its left derived functor. On the Grothendieck group, LZ and $R\Gamma$ descend to the canonical map from $V_{k-1}^{\otimes n}$ to $\Lambda^{r_1} V_{k-1} \otimes \dots \otimes \Lambda^{r_t} V_{k-1}$.

Proposition 5. (1) $[LZ^{(r_1, \dots, r_t)}] = \pi_{r_1} \otimes \dots \otimes \pi_{r_t}$.
(2) $[\epsilon_{(r_1, \dots, r_t)}[-\sum_{r=1}^t \frac{i_r(i_r-1)}{2}]] = i_{r_1} \otimes \dots \otimes i_{r_t}$.

Proof. By proposition 5.5 of [ES], one may easily prove the first part by computing on the basis of Verma modules. The second part follows from the generalized BGG resolution. $\square$

Let d be the codimension of $\mathfrak{q}$ in $\mathfrak{p}$.

Lemma 2. *The derived Zuckerman functor $R\Gamma_{\mathfrak{q}}^{\mathfrak{p}}$ and the inclusion functor $\epsilon_{\mathfrak{q}}^{\mathfrak{p}}$ satisfy the following adjointness properties in the derived category:*

- (1) $\text{Hom}(\epsilon_{\mathfrak{p}}^{\mathfrak{q}}X, Y) \cong \text{Hom}(X, R\Gamma_{\mathfrak{q}}^{\mathfrak{p}}Y)$
- (2) $\text{Hom}(X, \epsilon_{\mathfrak{p}}^{\mathfrak{q}}[2d]Y) \cong \text{Hom}(R\Gamma_{\mathfrak{q}}^{\mathfrak{p}}X, Y).$

Proof. See the proof of theorem 5 in [BFK]. □

Lemma 3. *The derived dual Zuckerman functor $LZ_{\mathfrak{q}}^{\mathfrak{p}}$ and the inclusion functor $\epsilon_{\mathfrak{q}}^{\mathfrak{p}}$ satisfy the following adjointness properties in the derived category:*

- (1) $\text{Hom}(\epsilon_{\mathfrak{p}}^{\mathfrak{q}}[-2d]X, Y) \cong \text{Hom}(X, LZ_{\mathfrak{q}}^{\mathfrak{p}}Y)$
- (2) $\text{Hom}(X, \epsilon_{\mathfrak{p}}^{\mathfrak{q}}Y) \cong \text{Hom}(LZ_{\mathfrak{q}}^{\mathfrak{p}}X, Y).$

Corollary 1. *The derived Zuckerman and dual Zuckerman functors are related by $R\Gamma_{\mathfrak{q}}^{\mathfrak{p}}[2d] \cong LZ_{\mathfrak{q}}^{\mathfrak{p}}$.*

Proof. By the previous two lemmas these functors are adjoints of the same functor so they are the same up to isomorphism. □

3.2. Equivalences of Categories. We look to generalize section 3.2.2 of [BFK] to the subcategories with generalized central character considered here. In order to make the notation more compact, we will have the following notation for several important subalgebras.

Definition 10. Let

- (1) $\mathfrak{p}_i$ to be the parabolic subalgebra corresponding to $(\underbrace{1, \dots, 1}_{i-1}, k, 1, \dots, 1).$
- (2) $\mathfrak{q}_i$ to be the parabolic subalgebra corresponding to $(\underbrace{1, \dots, 1}_{i-1}, k-1, 1, \dots, 1).$
- (3) $\mathfrak{r}_i$ be the parabolic subalgebra corresponding to $(\underbrace{1, \dots, 1}_i, k-2, 1, \dots, 1).$
- (4) $\mathfrak{s}_i$ be the parabolic subalgebra corresponding to $(\underbrace{1, \dots, 1}_{i-1}, 2, 1, \dots, 1).$
- (5) $\mathfrak{t}_i$ be the parabolic subalgebra corresponding to $(\underbrace{1, \dots, 1}_{i-1}, 3, 1, \dots, 1).$

Clearly $\mathfrak{p}_j \supset \mathfrak{q}_j$ and $\mathfrak{p}_j \supset \mathfrak{q}_{j-1}$. Recall from earlier the definitions of the functors $\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j}$, $\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_{j-1}}$, $LZ_{\mathfrak{q}_j}^{\mathfrak{p}_j}$, and $LZ_{\mathfrak{q}_{j-1}}^{\mathfrak{p}_j}$. Note that the derived Zuckerman functors in this setup could have non-zero cohomology functors only in degrees 0 through $2(k-1)$. As in [BFK], the middle cohomology functor plays a significant role. Given a generalized Verma module $M^{\mathfrak{p}_j}(\alpha)$, we would like to compute its image under the functor $L_{(k-1)}Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} \circ \epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j}$.

The main result of this subsection is there are equivalences of categories:

$$\mathcal{O}_{(d_{k-1}, \dots, d_0)}^{\mathfrak{p}_j} \cong \mathcal{O}_{(d_{k-1}, \dots, d_0)}^{\mathfrak{p}_{j+1}}.$$

The equivalences are compositions of inclusions into larger categories and the middle cohomology of the derived Zuckerman functor. The plan of the proof is exactly that from [BFK]. First the action of these functors on generalized Verma modules is computed. Then after some exactness and adjointness statements are proved, the equivalence will follow from lemma 2 of [BFK].

First we recall the generalized BGG resolution. Let S be a subset of simple roots defining a parabolic subalgebra $\mathfrak{p}$ and $W_{\mathfrak{p}}$ the corresponding Weyl group. Denote by $W^{\mathfrak{p}}$ the set of shortest coset representatives in $W/W_{\mathfrak{p}}$ and ρ_S half the sum of the positive roots of the subalgebra $\mathfrak{p}$.

Theorem 4. *Let μ be dominant integral. For all $j = 0, \dots, \dim u$, define*

$$C_j^S = \oplus_{w \in W^{\mathfrak{p}}, l(w)=j} M^{\mathfrak{p}}(w(\mu + \rho) - \rho_S).$$

Then there is an exact sequence

$$0 \rightarrow C_{\dim u}^S \rightarrow \dots \rightarrow C_0^S \rightarrow L(\mu + \rho) \rightarrow 0$$

with all maps nontrivial.

Let $\alpha = (a_1, \dots, a_n)$. In order for the generalized Verma module $M^{\mathfrak{p}_j}(\alpha)$ to be locally $\mathcal{U}(\mathfrak{p}_j)$ -finite,

$$a_j = k - 1, a_{j+1} = k - 2, \dots, a_{j+k-1} = 0.$$

From the previous theorem, it follows that there is a resolution of this generalized Verma module in terms of generalized Verma modules in larger parabolic categories.

Corollary 2. *There is an exact sequence in $\mathcal{O}_{\mathfrak{d}}^{q_j}$ of the form:*

$$0 \rightarrow M^{q_j}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha) \rightarrow M^{q_j}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow \cdots \rightarrow M^{q_j}(\alpha) \rightarrow M^{\mathfrak{p}_j}(\alpha) \rightarrow 0,$$

where σ_i permutes the elements a_{j+i-1} and a_{j+i} of $(a_j, \dots, a_{j+k-1})$.

Proof. This follows from the theorem and parabolic induction. $\square$

The cohomology functors on the modules $M^{q_j}(\sigma_1 \cdots \sigma_l \cdot \alpha)$ play a critical role.

Lemma 4. *Suppose $a_{k+j} = k - l - 1$. Then, $L_i Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{q_j}(\sigma_1 \cdots \sigma_l \cdot \alpha) \cong$*

$$\begin{aligned} M^{\mathfrak{p}_{j+1}}(a_1, \dots, a_{j-1}, k - l - 1, k - 1, \dots, 0, a_{k+j-1}, \dots, a_n) & \quad \text{if } i = k - l - 1 \\ 0 & \quad \text{if } i \neq k - l - 1. \end{aligned}$$

Proof. It is easily seen that

$$M^{q_j}(\sigma_1 \cdots \sigma_l \cdot \alpha) = M^{q_j}(a_1, \dots, a_{j-1}, k - l - 1, k - 1, k - 2, \dots, k - l, k - l - 2, \dots, 0, a_{k+j}, \dots, a_n).$$

Thus $L_i Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{q_j}(\sigma_1 \cdots \sigma_l \cdot \alpha) = 0$ if $a_{k+j} \neq k - l - 1$.

Now suppose S_k permutes the elements $(k - 1, k - 2, \dots, k - l, k - l - 2, \dots, 0, a_{k+j})$, by

$$\begin{aligned} \sigma_{l+1} \cdots \sigma_{k-1} \cdot (k - 1, k - 2, \dots, k - l, k - l - 2, \dots, 0, a_{k+j}) = \\ (k - 1, k - 2, \dots, k - l, a_{k+j}, k - l - 2, \dots, 0). \end{aligned}$$

The length of this element in the symmetric group is $k - l - 1$. Thus from [ES] proposition 5.5,

$$L_i Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{q_j}(\sigma_1 \cdots \sigma_l \cdot \alpha) \cong M^{\mathfrak{p}_{j+1}}(a_1, \dots, a_{j-1}, k - l - 1, k - 1, \dots, 0, a_{k+j-1}, \dots, a_n),$$

if $i = (k - l - 1)$. $\square$

Now we are in position to compute $L_i Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} \circ \epsilon_{\mathfrak{p}_j}^{q_j} M^{\mathfrak{p}_j}(\alpha)$. Let

$$\beta = (a_1, \dots, a_{j-1}, a_{j+k}, a_j, \dots, a_{j+k-1}, a_{j+k+1}, \dots, a_n).$$

Lemma 5. *There are isomorphisms: $L_{(k-1)} Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} \circ \epsilon_{\mathfrak{p}_j}^{q_j} M^{\mathfrak{p}_j}(\alpha) \cong$*

$$\begin{aligned} M^{\mathfrak{p}_{j+1}}(\beta) & \quad \text{if } i = k - 1 \\ 0 & \quad \text{if } i \neq k - 1 \end{aligned}$$

Proof. Consider the generalized BGG resolution. This gives rise to short exact sequences:

$$0 \rightarrow K_0 \rightarrow M^{q_j}(\alpha) \rightarrow M^{\mathfrak{p}_j}(\alpha) \rightarrow 0$$

$$0 \rightarrow K_1 \rightarrow M^{q_j}(\sigma_1 \cdot \alpha) \rightarrow K_0 \rightarrow 0$$

$$0 \rightarrow K_2 \rightarrow M^{q_j}(\sigma_1 \sigma_2 \cdot \alpha) \rightarrow K_1 \rightarrow 0$$

...

$$0 \rightarrow K_{k-2} \rightarrow M^{q_j}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow K_{k-3} \rightarrow 0$$

$$0 \rightarrow K_{k-1} \rightarrow M^{q_j}(\sigma_1 \cdots \sigma_{k-2} \sigma_{k-1} \cdot \alpha) \rightarrow K_{k-2} \rightarrow 0.$$

Assume $a_{k+j} = k - l - 1$. By the previous lemma, for all s

$$L_s Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{q_j}(e) \cong L_s Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{q_j}(\sigma_1 \cdot \alpha) \cong \cdots \cong L_s Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{q_j}(\sigma_1 \cdots \sigma_{l-1} \cdot \alpha) = 0.$$

Therefore

$$L_s Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} M^{\mathfrak{p}_j}(\alpha) \cong L_{s-1} Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} K_0 \cong \cdots \cong L_{s-l} Z_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} K_{l-1}.$$

Also by the previous lemma, for all s ,

$$L_s Z_{q_j}^{p_{j+1}} M^{q_j}(\sigma_{l+1} \cdot \alpha) \cong \dots \cong L_s Z_{q_j}^{p_{j+1}} M^{q_j}(\sigma_1 \dots \sigma_{k-1} \cdot \alpha) \cong 0.$$

Therefore for all s ,

$$L_s Z_{q_j}^{p_{j+1}} K_{k-2} \cong \dots \cong L_{s+l} Z_{q_j}^{p_{j+1}} K_l \cong 0.$$

Now consider the short exact sequence

$$0 \rightarrow K_l \rightarrow M^{q_j}(\sigma_1 \dots \sigma_l \cdot \alpha) \rightarrow K_{l-1} \rightarrow 0.$$

Due to the above

$$L_s Z_{q_j}^{p_{j+1}} M^{q_j}(\sigma_1 \dots \sigma_l \cdot \alpha) \cong L_s Z_{q_j}^{p_{j+1}} K_{l-1} \cong L_{s+l} Z_{q_j}^{p_{j+1}} \epsilon_{p_j}^{q_j} M^{p_j}(\alpha).$$

By the previous lemma

$$L_s Z_{q_j}^{p_{j+1}} M^{q_j}(\sigma_1 \dots \sigma_l \cdot \alpha) \cong M^{p_{j+1}}(\beta)$$

for $s = k - l - 1$, and it is zero otherwise.

Thus $L_s Z_{q_j}^{p_{j+1}} M^{p_j}(\alpha) \cong M^{p_{j+1}}(\beta)$ if $s = k - 1$ and is zero otherwise. $\square$

Lemma 6. Suppose $i \neq k - 1$. Then for any module M in $\mathcal{O}_{(d_{k-1}, d_{k-2}, \dots, d_0)}^{p_j}$, there is an isomorphism

$$L_i Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} M = 0.$$

Proof. We know this is true by the previous lemma for generalized Verma modules. It is then true for projective modules by induction on the length of their Verma flag. Now consider a simple module S . It has a projective resolution which gives rise to several short exact sequences:

$$0 \rightarrow K_1 \rightarrow P_1 \rightarrow S \rightarrow 0$$

$$0 \rightarrow K_2 \rightarrow P_2 \rightarrow K_1 \rightarrow 0$$

...

$$0 \rightarrow K_r \rightarrow P_r \rightarrow K_{r-1} \rightarrow 0.$$

The first short exact sequence implies $L_i Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} S \cong L_{(i-1)} Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} K_1$ for $i < (k - 1)$. Continuing in this way we get $L_i Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} S \cong L_{i-r} Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} P_r$ for $i < (k - 1)$. Since $K_r \cong 0$, $L_i Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} S \cong 0$, for $i < (k - 1)$. By the duality theorem for Zuckerman functors, this isomorphism is true for all $i \neq (k - 1)$. Now the lemma follows for any M by induction on the length of its composition series. $\square$

Proposition 6. There is an equivalence of categories

$$L_{k-1} Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} : \mathcal{O}_{\mathbf{d}}^{p_j} \xrightarrow{\cong} \mathcal{O}_{\mathbf{d}}^{p_{j+1}}.$$

The inverse equivalence is given by $L_{k-1} Z_{q_j}^{p_j} \circ \epsilon_{p_{j+1}}^{q_j}$.

Proof. These functors send generalized Verma modules to generalized Verma modules. By examining the long exact sequence for the functor and using that the cohomology functors vanish in all but one degree, it is clear that this functor is exact. It suffices by lemma 2 of [BFK] to show that they are adjoint. In the derived category $LZ_{q_j}^{p_{j+1}}$ is left adjoint to $\epsilon_{p_{j+1}}^{q_j}$ by lemma 3. Furthermore, $LZ_{q_j}^{p_{j+1}}[-2(k-1)]$ is right adjoint to $\epsilon_{p_{j+1}}^{q_j}$. Thus we have

$$\begin{aligned} \text{Hom}_{\mathcal{O}}(L_{k-1} Z_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j} M, N) &\cong \text{Hom}_{D^b(\mathcal{O})}(LZ_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j}[-(k-1)]M, N) \\ &\cong \text{Hom}_{D^b(\mathcal{O})}(\epsilon_{p_j}^{q_j} M, \epsilon_{p_{j+1}}^{q_j}[k-1]N) \\ &\cong \text{Hom}_{D^b(\mathcal{O})}(M, LZ_{q_j}^{p_j} \epsilon_{p_{j+1}}^{q_j}[-(k-1)]N) \\ &\cong \text{Hom}_{\mathcal{O}}(M, L_{k-1} Z_{q_j}^{p_j} \epsilon_{p_{j+1}}^{q_j} N). \end{aligned}$$

$\square$

Remark 3. The functor $LZ_{q_j}^{p_{j+1}} \circ \epsilon_{p_j}^{q_j}$ gives an equivalence of the corresponding derived categories.

Next assume that all of the a_i are 1 or $k-1$.

Proposition 7. *There is an equivalence of categories*

$$\mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_i, k, a_{i+1}, \dots, a_n)}(\mathfrak{gl}_n) \cong \mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_{i+1}, k, a_{i+2}, \dots, a_n)}(\mathfrak{gl}_n)$$

given by the functor

$$L_{(k-1)} Z_{(a_1, \dots, a_i, a_{i+1}, k-a_{i+1}, a_{i+1}, \dots, a_n)}^{(a_1, \dots, a_{i+1}, k, \dots, a_n)} \epsilon_{(a_1, \dots, a_i, k, a_{i+1}, \dots, a_n)}^{(a_1, \dots, a_i, a_{i+1}, k-a_{i+1}, a_{i+1}, \dots, a_n)}.$$

Proof. The proof is exactly the same as the previous proposition. $\square$

Composing the equivalences given by the proposition above, we get the following equivalence.

$$\Delta_{(a_1, \dots, a_i, k, \dots, a_r)}^{(a_1, \dots, a_j, k, \dots, a_r)} : \mathcal{O}_{\mathbf{d}}^{(a_1, a_i, k, \dots, a_r)} \rightarrow \mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_j, k, \dots, a_r)}.$$

Next we define functors between parabolic categories for Lie algebras of different ranks.

Definition 11. (1) Let Y be the one dimensional $\mathfrak{gl}_k \oplus \mathfrak{gl}_n$ - module with weight

$$-\frac{n}{2}(e_1 + \dots + e_k) + \frac{k}{2}(e_{k+1} + \dots + e_{k+n}).$$

(2) Let Y' be the one dimensional $\mathfrak{gl}_n$ - module with weight $-\frac{k}{2}(e_1 + \dots + e_n)$.

(3) Let W be the finite dimensional irreducible $\mathfrak{gl}_k$ - module with highest weight $(k-1)e_1 + \dots + (0)e_k - \rho_k$.

(4) The functor $\zeta : \mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_r)}(\mathfrak{gl}_n) \rightarrow \mathcal{O}_{\mathbf{d}'}^{(k, a_1, \dots, a_r)}(\mathfrak{gl}_{k+n})$ is given by

$$\zeta(M) = \text{Ind}_{\mathcal{U}(\mathfrak{gl}_k + \mathfrak{gl}_n)}^{\mathcal{U}(\mathfrak{gl}_{k+n})}(Y \otimes W \otimes M)$$

where $\mathbf{d}' = (d_{k-1} + 1, \dots, d_0 + 1)$.

(5) The functor $\nu : \mathcal{O}_{\mathbf{d}'}^{(k, a_1, \dots, a_r)}(\mathfrak{gl}_{k+n}) \rightarrow \mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_r)}(\mathfrak{gl}_n)$ is given by taking the sum of the weight spaces of a module M for the weights of the form

$$(k-1)e_1 + (k-2)e_2 + \dots + (0)e_k + x_{k+1}e_{k+1} + \dots + x_{k+n}e_{k+n} - \rho,$$

tensor with Y' , where the $x_i \in \mathbb{Z}$.

The next lemma is a direct generalization of proposition 17 of [BFK].

Lemma 7. *The functors ζ and ν are inverse equivalences of categories.*

Next we assign functors to various oriented tangles. If an edge is labeled by $k-1$, then we give it a down arrow. If an edge is labeled by 1, then it is given an up arrow.




FIGURE 10.

To the tangle in Figure 10 we assign the functor $\cup_{i,+r} : D^b(\mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_r)}) \rightarrow D^b(\mathcal{O}_{\mathbf{d}'}^{(a_1, \dots, a_{i-1}, 1, k-1, a_i, \dots, a_r)})$. It is given by

$$\cup_{i,+r}(M) = \epsilon_{(a_1, \dots, a_{i-1}, k, \dots, a_r)}^{(a_1, \dots, a_{i-1}, 1, k-1, \dots, a_r)} [-(k-1)] \Delta_{(k, a_1, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, a_r)} \zeta(M).$$




FIGURE 11.

To the tangle in Figure 11 we assign the functor $\cup_{i,-,r}: D^b(\mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_r)}) \rightarrow D^b(\mathcal{O}_{\mathbf{d}'}^{(a_1, \dots, a_{i-1}, k-1, 1, a_i, \dots, a_r)})$. It is given by

$$\cup_{i,-,r}(M) = \epsilon_{(a_1, \dots, a_{i-1}, k, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k-1, 1, \dots, a_r)} [-(k-1)] \Delta_{(k, a_1, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, a_r)} \zeta(M).$$




FIGURE 12.

To the tangle in Figure 12 we assign the functor $\cap_{i,+,r}: D^b(\mathcal{O}_{\mathbf{d}'}^{(a_1, \dots, a_r)}) \rightarrow D^b(\mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_{i-1}, a_{i+2}, \dots, a_r)})$. It is given by

$$\cap_{i,+,r}(M) = \nu \Delta_{(a_1, \dots, a_{i-1}, k, a_{i+2}, \dots, a_r)}^{(k, a_1, \dots, a_{i-1}, a_{i+2}, \dots, a_r)} LZ_{(a_1, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, a_{i+2}, \dots, a_r)}.$$




FIGURE 13.

To the tangle in Figure 13 we assign the functor $\cap_{i,-,r}: D^b(\mathcal{O}_{\mathbf{d}'}^{(a_1, \dots, a_r)}) \rightarrow D^b(\mathcal{O}_{\mathbf{d}}^{(a_1, \dots, a_{i-1}, a_{i+2}, \dots, a_r)})$. It is given by

$$\cap_{i,-,r}(M) = \nu \Delta_{(a_1, \dots, a_{i-1}, k, a_{i+2}, \dots, a_r)}^{(k, a_1, \dots, a_{i-1}, a_{i+2}, \dots, a_r)} LZ_{(a_1, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, a_{i+2}, \dots, a_r)}.$$

4. DIAGRAM RELATIONS

In this section we verify that the relations between various intertwiners which have a graphical interpretation given in figures 3, 4, 5, 6, and 7 gives rise to relations between various inclusion and Zuckerman functors.

4.1. Diagram 1. Our goal is to prove that $LZ_{\mathbf{q}_j}^{\mathbf{p}_j} \epsilon_{\mathbf{p}_j}^{\mathbf{q}_j}$ is a direct sum of shifted identity functors. First we compute the cohomology functors on the generalized Verma module $M^{\mathbf{p}_j}(\alpha)$ where $\alpha = (a_1, \dots, a_{j-1}, k-1, \dots, 0, a_{j+k}, \dots, a_n)$.

Lemma 8. $L_i Z_{\mathbf{q}_j}^{\mathbf{p}_j} \epsilon_{\mathbf{p}_j}^{\mathbf{q}_j} M^{\mathbf{p}_j}(\alpha) \cong M^{\mathbf{p}_j}(\alpha)$ if i is even and $0 \leq i \leq 2(k-1)$. Otherwise it is 0.

Proof. First we note that $L_s Z M^{\mathbf{q}_j}(\sigma_1 \dots \sigma_t, \alpha) \cong$

$$\begin{aligned} M^{\mathbf{p}_j}(\alpha) & \quad \text{if} \quad s = t \\ 0 & \quad \text{if} \quad s \neq t. \end{aligned}$$

Suppose $i = 2r$. Then the above fact and the short exact sequences from the previous section imply

$$\begin{aligned} L_i Z_{\mathbf{q}_j}^{\mathbf{p}_j} \epsilon_{\mathbf{p}_j}^{\mathbf{q}_j} M^{\mathbf{p}_j}(\alpha) & \cong L_{i-1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_0 \\ L_{i-1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_0 & \cong L_{i-2} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_1 \\ & \dots \\ L_{i-r+1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_{r-2} & \cong L_{i-r} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_{r-1}. \end{aligned}$$

Now consider $0 \rightarrow K_r \rightarrow M^{\mathbf{q}_j}(\sigma_1 \dots \sigma_r, \alpha) \rightarrow K_{r-1} \rightarrow 0$. This gives a long exact sequence

$$\begin{aligned} 0 \rightarrow L_{i-r+1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_{r-1} \rightarrow L_{i-r} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_r \rightarrow L_{i-r} Z_{\mathbf{q}_j}^{\mathbf{p}_j} M^{\mathbf{q}_j}(\sigma_1 \dots \sigma_r, \alpha) \cong M^{\mathbf{p}_j}(\alpha) \rightarrow \\ L_{i-r-1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_{r-1} \rightarrow L_{i-r-1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} K_r \rightarrow L_{i-r-1} Z_{\mathbf{q}_j}^{\mathbf{p}_j} M^{\mathbf{q}_j}(\sigma_1 \dots \sigma_r, \alpha) \cong 0. \end{aligned}$$

Thus it suffices to show $L_{i-r}Z_{q_j}^{\mathfrak{p}_j}K_r \cong L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r \cong 0$. Using the first fact of the proof we easily see

$$\begin{aligned} L_{i-r}Z_{q_j}^{\mathfrak{p}_j}K_r &\cong L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_{r+1} \\ L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_{r+1} &\cong L_{i-r-2}Z_{q_j}^{\mathfrak{p}_j}K_{r+2} \\ &\dots \\ L_{i-k+2}Z_{q_j}^{\mathfrak{p}_j}K_{k-2} &\cong L_{i-k+1}Z_{q_j}^{\mathfrak{p}_j}K_{k-1} \cong 0. \end{aligned}$$

Thus $L_{i-r}Z_{q_j}^{\mathfrak{p}_j}K_r \cong 0$. Again using the first fact we easily see that

$$\begin{aligned} L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r &\cong L_{i-r-2}Z_{q_j}^{\mathfrak{p}_j}K_{r+1} \\ L_{i-r-2}Z_{q_j}^{\mathfrak{p}_j}K_{r+1} &\cong L_{i-r-3}Z_{q_j}^{\mathfrak{p}_j}K_{r+2} \\ &\dots \\ L_{i-k+1}Z_{q_j}^{\mathfrak{p}_j}K_{k-2} &\cong L_{i-k}Z_{q_j}^{\mathfrak{p}_j}K_{k-1} \cong 0. \end{aligned}$$

Thus $L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r \cong 0$.

Now we proceed for odd i . Let $i = 2r + 1$. Then we get

$$\begin{aligned} L_iZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j}M^{\mathfrak{p}_j}(\alpha) &\cong L_{i-1}Z_{q_j}^{\mathfrak{p}_j}K_0 \\ L_{i-1}Z_{q_j}^{\mathfrak{p}_j}K_0 &\cong L_{i-2}Z_{q_j}^{\mathfrak{p}_j}K_1 \\ &\dots \\ L_{i-r+1}Z_{q_j}^{\mathfrak{p}_j}K_{r-2} &\cong L_{i-r}Z_{q_j}^{\mathfrak{p}_j}K_{r-1}. \end{aligned}$$

Now consider $0 \rightarrow K_r \rightarrow M^{\mathfrak{q}_j}(\sigma_1 \cdots \sigma_r.\alpha) \rightarrow K_{r-1} \rightarrow 0$. This gives a long exact sequence

$$\rightarrow L_{i-r}Z_{q_j}^{\mathfrak{p}_j}M^{\mathfrak{q}_j}(\sigma_1 \cdots \sigma_r.\alpha) \cong 0 \rightarrow L_{i-r}Z_{q_j}^{\mathfrak{p}_j}K_{r-1} \rightarrow L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r \rightarrow L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}M^{\mathfrak{q}_j}(\sigma_1 \cdots \sigma_r.\alpha) \rightarrow .$$

Thus it suffices to prove $L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r \cong 0$. Again using the first fact, it is easy to see that

$$\begin{aligned} L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r &\cong L_{i-r-2}Z_{q_j}^{\mathfrak{p}_j}K_{r+1} \\ L_{i-r-2}Z_{q_j}^{\mathfrak{p}_j}K_{r+1} &\cong L_{i-r-3}Z_{q_j}^{\mathfrak{p}_j}K_{r+2} \\ &\dots \\ L_{i-k+1}Z_{q_j}^{\mathfrak{p}_j}K_{k-2} &\cong L_{i-k}Z_{q_j}^{\mathfrak{p}_j}K_{k-1} \cong 0. \end{aligned}$$

Thus $L_{i-r-1}Z_{q_j}^{\mathfrak{p}_j}K_r \cong 0$. □

Lemma 9. Let $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} T(X)$ be a distinguished triangle in a triangulated category. If $h = 0$, then this triangle is isomorphic to $X \rightarrow X \oplus Z \rightarrow Z \rightarrow T(X)$.

Proof. Since h is zero, using the axioms of a triangulated category one could construct a morphism from the triangle $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} T(X)$ to the triangle $X \rightarrow X \oplus Z \rightarrow Z \rightarrow T(X)$. The map from Y to $X \oplus Z$ is an isomorphism because the other two maps in the morphism of triangles are isomorphisms. □

Proposition 8. $LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} \cong \oplus_{r=0}^{k-1} \text{Id}[2r]$.

Proof. We claim that for $0 \leq n \leq k-1$,

$$\tau^{\leq -2n} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \cong \oplus_{r=0}^{(k-1)-n} M^{\mathfrak{p}_j}(\alpha)[2(k-1) - 2r].$$

We proceed by induction. The base case is $n = k-1$. Assume by induction

$$\tau^{\leq -2n} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \cong \oplus_{r=0}^{(k-1)-n} M^{\mathfrak{p}_j}(\alpha)[2(k-1) - 2r].$$

Consider the distinguished triangle

$$\tau^{\leq (-2n+1)} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \rightarrow \tau^{\leq (-2n+2)} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \rightarrow H^{-2n+2} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha)[2n-2].$$

This triangle is isomorphic to

$$\tau^{\leq (-2n)} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \rightarrow \tau^{\leq (-2n+2)} LZ_{q_j}^{\mathfrak{p}_j}\epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \rightarrow M^{\mathfrak{p}_j}(\alpha)[2n-2].$$

We would like to compute

$$\mathrm{Hom}(M^{\mathfrak{p}_j}(\alpha)[2n-2], \tau^{\leq -2n} LZ_{\mathfrak{q}_j}^{\mathfrak{p}_j} \epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}[1]).$$

By the induction hypothesis this is equal to

$$\mathrm{Hom}(M^{\mathfrak{p}_j}(\alpha)[2n-2], \oplus_{r=0}^{2(k-1)-n} M^{\mathfrak{p}_j}(\alpha)[2(k-1)-2r+1]).$$

Since there are no non-trivial morphism between a generalized Verma module and itself in different degrees, the above must be zero. Thus by the previous lemma

$$\tau^{\leq -2n+2} LZ_{\mathfrak{q}_j}^{\mathfrak{p}_j} \epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \cong \tau^{\leq -2n} LZ_{\mathfrak{q}_j}^{\mathfrak{p}_j} \epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} M^{\mathfrak{p}_j}(\alpha) \oplus M^{\mathfrak{p}_j}(\alpha)[2n-2] \cong \oplus_{r=0}^{2(k-1)-(n-1)} M^{\mathfrak{p}_j}(\alpha)[2(k-1)-2r]$$

which completes the induction.

In the Koszul dual situation, the inclusion functor and derived Zuckerman functor become graded translation off an intersection of walls and graded translation back respectively. For this fact see [MOS] or [Rh]. Then by theorem 4.12 of [J], the composition of these functors is a direct sum of identity functors with various shifts in the grading. By Koszul duality these become identity functors with shifts in the grading and homological shifts. The computation above for the generalized module indicate exactly what the homological shifts are. $\square$

Corollary 3. $LZ_{\mathfrak{q}_j}^{\mathfrak{p}_j} \epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} [-(k-1)] \cong \oplus_{r=0}^{k-1} \mathrm{Id}[2r - (k-1)].$

Remark 4. Lemma 3.5.4 of [BGS] is a geometric version of this result.

4.2. Diagram 2. This subsection is a restatement of proposition 16 of [BFK]. It could also be proven using the techniques for diagram 1, but here we just refer to [BFK]. The following is a functorial version of the second relation in the graphical calculus of [MOY].

Proposition 9. *There is an isomorphism of functors $LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_i} \cong \mathrm{Id} \oplus \mathrm{Id}[2]$.*

Proof. See [BFK] proposition 16. $\square$

Corollary 4. $\epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i} \cong \epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i}[1] \oplus \epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i}[-1].$

Proof. This is a direct consequence of the previous proposition. $\square$

4.3. Diagram 3. We would like to have a functorial isomorphism corresponding to the third graphical relation.

Recall the definitions of the algebras, $\mathfrak{p}_i$, $\mathfrak{q}_i$, and $\mathfrak{s}_i$. There are obvious inclusions of subalgebras: $\mathfrak{q}_i \subset \mathfrak{p}_i$, $\mathfrak{q}_{i+1} \subset \mathfrak{s}_i + \mathfrak{q}_{i+1}$, $\mathfrak{q}_{i-1} \subset \mathfrak{p}_i$.

We begin by studying the generalized Verma module with highest weight α , $M^{\mathfrak{p}_{i+1}}(\alpha) =$

$$M^{\mathfrak{p}_{i+1}}(a_1 \dots a_i, k-1 \dots, 0, a_{i+k+1} \dots, a_n).$$

For notational purposes, we will omit the components $a_1, \dots, a_{i-1}, a_{i+k+1}, \dots, a_n$ from the weights in the generalized Verma modules in this subsection. Denote by $M^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_t \alpha)$ the module

$$M^{\mathfrak{q}_{i+1}}(a_i, k-1-t, k-1, \dots, \widehat{k-1-t}, \dots, 0).$$

(The term $k-1-t$ is omitted.)

The generalized BGG resolution gives rise to the following long exact sequence

$$0 \rightarrow M^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-1} \alpha) \rightarrow M^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-2} \alpha) \rightarrow \dots \rightarrow M^{\mathfrak{q}_{i+1}}(e \alpha) \rightarrow M^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \rightarrow 0.$$

Then the above sequence gives rise to the following set of short exact sequences:

$$\begin{aligned} 0 &\rightarrow K_0 \rightarrow M^{\mathfrak{q}_{i+1}}(e \alpha) \rightarrow M^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \rightarrow 0 \\ 0 &\rightarrow K_1 \rightarrow M^{\mathfrak{q}_{i+1}}(\sigma_1 \alpha) \rightarrow K_0 \rightarrow 0 \\ &\dots \\ 0 &\rightarrow K_{k-2} \rightarrow M^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-2} \alpha) \rightarrow K_{k-3} \rightarrow 0 \\ 0 &\rightarrow M^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-1} \alpha) \rightarrow K_{k-2} \rightarrow 0. \end{aligned}$$

Our first goal is to compute $LZ_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0)$.

Lemma 10. *Let $a_i = l$.*

(1) *Suppose $s = 0$ or $s = 2$. Then*

$$L_s Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0) = 0$$

(2) *Otherwise we have isomorphisms:*

$$\begin{aligned} L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0) &\cong L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(e.\alpha)/L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 \\ L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 &\cong L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1.\alpha)/L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 \\ &\dots \\ L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-3} &\cong L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l-2}.\alpha)/L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-2} \\ L_s Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-2} &\cong L_{s-1} Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-1} \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-1} &\cong L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l}.\alpha)/L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l} \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l} &\cong L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l+1}.\alpha)/L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l+1} \\ &\dots \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-4} &\cong L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-3}.\alpha)/L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-3} \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-3} &\cong L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-2}.\alpha)/L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-1}.\alpha). \end{aligned}$$

Proof. By proposition 5.5 of [ES], $L_s Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_t.\alpha) \cong$

$$\begin{aligned} M^{\mathbf{s}_i+\mathbf{q}_{i+1}}(a_i, k-1-t, k-1, \dots, k-\hat{1}-t, \dots, 0) &\quad \text{if } s=0, l > k-1-t \\ M^{\mathbf{s}_i+\mathbf{q}_{i+1}}(k-1-t, a_i, k-1, \dots, k-\hat{1}-t, \dots, 0) &\quad \text{if } s=1, l < k-1-t \\ 0 &\quad \text{if otherwise} \end{aligned}$$

Now the functor $LZ_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}$ induces long exact sequences for all the short exact sequences stated before the lemma.

$$\begin{aligned} 0 \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 &\rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(e.\alpha) \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}M^{\mathbf{p}_{i+1}}(\alpha) \rightarrow \\ L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 &\rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(e) \rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}M^{\mathbf{p}_{i+1}}(\alpha) \rightarrow \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 &\rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(e) \rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}M^{\mathbf{p}_{i+1}}(\alpha) \rightarrow 0 \end{aligned}$$

$$\begin{aligned} 0 \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 &\rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1.\alpha) \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 \rightarrow \\ L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 &\rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1.\alpha) \rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 \rightarrow \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 &\rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1.\alpha) \rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_0 \rightarrow 0 \end{aligned}$$

$$\begin{aligned} 0 \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_2 &\rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1\sigma_2.\alpha) \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 \rightarrow \\ L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_2 &\rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1\sigma_2.\alpha) \rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 \rightarrow \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_2 &\rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1\sigma_2.\alpha) \rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_1 \rightarrow 0 \end{aligned}$$

...

$$\begin{aligned} 0 \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-2} &\rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l-2}.\alpha) \rightarrow L_2 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-3} \rightarrow \\ L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-2} &\rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l-2}.\alpha) \rightarrow L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-3} \rightarrow \\ L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-2} &\rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}M^{\mathbf{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l-2}.\alpha) \rightarrow L_0 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i+\mathbf{q}_{i+1}}K_{k-l-3} \rightarrow 0. \end{aligned}$$

This concludes the first set of exact sequences. Then we have the following isomorphisms.

$$\begin{aligned} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-1} \cdot \alpha) &\cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-1} \cdot \alpha) \\ &\cong L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-1} \cdot \alpha) \\ &\cong 0. \end{aligned}$$

Now we have this second set of exact sequences:

$$\begin{aligned} 0 \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l} &\rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l} \cdot \alpha) \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-1} \rightarrow \\ L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l} &\rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l} \cdot \alpha) \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-1} \rightarrow \\ L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l} &\rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l} \cdot \alpha) \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-1} \rightarrow 0 \\ \\ 0 \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+1} &\rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l+1} \cdot \alpha) \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l} \rightarrow \\ L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+1} &\rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l+1} \cdot \alpha) \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l} \rightarrow \\ L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+1} &\rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l+1} \cdot \alpha) \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l} \rightarrow 0 \\ \\ 0 \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+2} &\rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l+2} \cdot \alpha) \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+1} \rightarrow \\ L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+2} &\rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l+2} \cdot \alpha) \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+1} \rightarrow \\ L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+2} &\rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l+2} \cdot \alpha) \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l+1} \rightarrow 0 \\ &\dots \end{aligned}$$

$$\begin{aligned} 0 \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-2} &\rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3} \rightarrow \\ L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-2} &\rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3} \rightarrow \\ L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-2} &\rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3} \rightarrow 0. \end{aligned}$$

Now we know

$$L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-2} \cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-2} \cong 0,$$

and that

$$L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-2} \cong M^{s_i+q_{i+1}}(a_i, 0, k-1, \dots, 1).$$

In the first set of sequences, $L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_t \cdot \alpha) = 0$. Thus $L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{t-1} \cong 0$ and $L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{p_{i+1}}(\alpha) \cong 0$. Also $L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_t \cdot \alpha) \cong 0$ so $L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_t \cong 0$. Therefore each sequence in the first set produces a short exact sequence

$$0 \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_t \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_t \cdot \alpha) \rightarrow L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{t-1} \rightarrow 0.$$

Thus we have part of the lemma:

$$\begin{aligned} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{p_{i+1}}(\alpha) &\cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(e \cdot \alpha) / L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_0 \\ L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_0 &\cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdot \alpha) / L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_1 \\ &\dots \\ L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-3} &\cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-2} \cdot \alpha) / L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-2}. \end{aligned}$$

In the second set of exact sequences

$$\begin{aligned} L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_t \cdot \alpha) &\cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_t \cdot \alpha) \\ &\cong 0. \end{aligned}$$

Thus

$$\begin{aligned} L_2 Z_{q_{i+1}}^{s_i+q_{i+1}} K_t &\cong L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{t+1} \\ &\cong 0. \end{aligned}$$

Therefore each sequence in the second set produces a short exact sequence

$$0 \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_t \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_t \cdot \alpha) \rightarrow L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{t-1} \rightarrow 0.$$

These short exact sequences give the rest of the lemma. $\square$

Next we would like to include this object back into $\mathcal{O}^{q_{i+1}}$ and then apply $LZ_{q_{i+1}}^{p_{i+1}}$. That is, we would like to compute

$$LZ_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} LZ_{q_{i+1}}^{s_i+q_{i+1}} M^{p_{i+1}}(a_i, k-1, \dots, 0).$$

Therefore we want to compute

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{p_{i+1}}(a_i, k-1, \dots, 0).$$

We begin with something simpler.

Lemma 11. *There is an isomorphism:*

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) / L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha)) \cong \\ M^{p_{i+1}}(a_i, k-1, \dots, 0) \quad \text{if } s = k-2, k \\ 0 \quad \text{if } s \neq k-2, k. \end{aligned}$$

Proof. Consider the short exact sequence

$$0 \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha) \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3} \rightarrow 0.$$

Once we understand the first two terms, we will be able to understand the functor applied to K_{k-3} .

So first we compute $L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha)$. This is isomorphic to

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} M^{s_i+q_{i+1}}(a_i, 0, k-1, \dots, 1).$$

For this, consider the short exact sequence

$$0 \rightarrow M^{q_{i+1}}(0, a_i, k-1, \dots, 1) \rightarrow M^{q_{i+1}}(a_i, 0, k-1, \dots, 1) \rightarrow M^{s_i+q_{i+1}}(a_i, 0, k-1, \dots, 1) \rightarrow 0.$$

Now $L_s Z_{q_{i+1}}^{p_{i+1}} M^{q_{i+1}}(0, a_i, k-1, \dots, 1) \cong 0$ for all s . Also,

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} M^{q_{i+1}}(a_i, 0, k-1, \dots, 1) &\cong \\ M^{p_{i+1}}(a_i, k-1, \dots, 0) &\quad \text{if } s = k-1 \\ 0 &\quad \text{if } s \neq k-1. \end{aligned}$$

Thus $L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha) \cong$

$$\begin{aligned} M^{p_{i+1}}(a_i, k-1, \dots, 0) &\quad \text{if } s = k-1 \\ 0 &\quad \text{if } s \neq k-1. \end{aligned}$$

Next we compute $L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha)$. This is isomorphic to

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} M^{s_i+q_{i+1}}(a_i, 1, k-1, \dots, 2, 0).$$

For this, consider the short exact sequence

$$0 \rightarrow M^{q_{i+1}}(1, a_i, k-1, \dots, 2, 0) \rightarrow M^{q_{i+1}}(a_i, 1, k-1, \dots, 2, 0) \rightarrow M^{s_i+q_{i+1}}(a_i, 1, k-1, \dots, 2, 0) \rightarrow 0.$$

Now $L_s Z_{q_{i+1}}^{p_{i+1}} M^{q_{i+1}}(1, a_i, k-1, \dots, 2, 0) \cong 0$ for all s . Also,

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} M^{q_{i+1}}(a_i, 1, k-1, \dots, 2, 0) &\cong \\ M^{p_{i+1}}(a_i, k-1, \dots, 0) &\quad \text{if } s = k-2 \\ 0 &\quad \text{if } s \neq k-2. \end{aligned}$$

Thus

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) &\cong \\ M^{p_{i+1}}(a_i, k-1, \dots, 0) &\quad \text{if } s = k-2 \\ 0 &\quad \text{if } s \neq k-2. \end{aligned}$$

Finally, we consider the long exact sequence for the functor $LZ_{q_{i+1}}^{p_{i+1}}$ for the short exact sequence considered in the beginning of the proof. It is

$$\begin{aligned} \cdots \rightarrow & L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha) \rightarrow L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow \\ & L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3} \rightarrow L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-1} \cdot \alpha) \rightarrow \\ & L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-2} \cdot \alpha) \rightarrow \cdots \end{aligned}$$

If $s = k - 2$ a portion of this long exact sequence reduces to

$$0 \rightarrow M^{p_{i+1}}(a_i, k-1, \dots, 0) \rightarrow L_{k-2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3}) \rightarrow 0.$$

Thus

$$L_{k-2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} K_{k-3} \cong M^{p_{i+1}}(a_i, k-1, \dots, 0).$$

If $s - 1 = k - 1$, then $s = k$ and the sequence reduces to

$$0 \rightarrow L_k Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-3}) \rightarrow M^{p_{i+1}}(a_i, k-1, \dots, 0) \rightarrow 0.$$

Thus

$$L_k Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} K_{k-3} \cong M^{p_{i+1}}(a_i, k-1, \dots, 0).$$

For all other values it is 0. $\square$

Lemma 12. Suppose $3 \leq t \leq l+1$ and $a_i = l$. If $s = k - t + 1, k - t + 3, \dots, k + t - 3$, then

$$\begin{aligned} & L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) / L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1}) \\ & \cong M^{p_{i+1}}(a_i, k-1, \dots, 0). \end{aligned}$$

Otherwise it is zero.

Proof. We proceed by induction on t . The base case is the previous lemma.

As in the proof of the previous lemma, we find that

$$\begin{aligned} & L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \cong \\ & M^{p_{i+1}}(a_i, k-1, \dots, 0) \quad \text{if } s = k - t + 1 \\ & 0 \quad \text{if } s \neq k - t + 1. \end{aligned}$$

We also have analogously a long exact sequence

$$\begin{aligned} \cdots \rightarrow & L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \rightarrow L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \rightarrow \\ & L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \rightarrow L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \rightarrow \\ & L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \rightarrow \cdots \end{aligned}$$

Now note that by the induction hypothesis $L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \cong 0$ unless $s = k - (t-1) + 1, \dots, k + (t-1) - 3$.

Suppose $s \leq k - t$. Then

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \cong L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \cong 0.$$

Now just apply the induction hypothesis.

Suppose $s \geq k - t + 3$. Then

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \cong L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \cong 0.$$

Now just apply the induction hypothesis.

Suppose $s = k - t + 1$. Then the long exact sequence and the induction hypothesis gives

$$0 \rightarrow M^{p_{i+1}}(a_i, k-1, \dots, 0) \rightarrow L_{k-t+1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \rightarrow L_{k-t} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \cong 0.$$

Suppose $s = k - t + 2$. Then the long exact sequence and the induction hypothesis gives

$$0 \rightarrow L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \rightarrow L_{k-t+1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \cong 0.$$

□

Corollary 5. If $s = k - l, k - l + 2, \dots, k + l - 2$,

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-1} \cong M^{p_{i+1}}(a_i, k - 1, \dots, 0)$$

Otherwise it is zero.

Lemma 13. If $s = k - l - 1, k - l + 1, \dots, k + l - 1$ then

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-2} \cdot \alpha) / L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-2}) \\ \cong M^{p_{i+1}}(a_i, k - 1, \dots, 0). \end{aligned}$$

Otherwise it is zero.

Proof. From the previous corollary we know that

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-1} &\cong \\ M^{p_{i+1}}(a_i, k - 1, \dots, 0) &\quad \text{if } s = k - l, k - l + 2, \dots, k + l - 2 \\ 0 &\quad \text{if } s \neq k - l, k - l + 2, \dots, k + l - 2. \end{aligned}$$

By lemma 10, $L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-2} \cong L_0 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-1}$. Thus

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-2} &\cong \\ M^{p_{i+1}}(a_i, k - 1, \dots, 0) &\quad \text{if } s = k - l, k - l + 2, \dots, k + l - 2 \\ 0 &\quad \text{if } s \neq k - l, k - l + 2, \dots, k + l - 2. \end{aligned}$$

Next we compute $L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-2} \cdot \alpha)$. It is isomorphic to

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} M^{s_i+q_{i+1}}(l + 1, a_i, k - 1, \dots, l + 2, l, \dots, 0).$$

In order to handle this, consider the short exact sequence

$$M^{q_{i+1}}(l + 1, k - 1, \dots, l + 2, l, \dots, 0) \hookrightarrow M^{q_{i+1}}(l + 1, a_i, k - 1, \dots, l + 2, l, \dots, 0) \twoheadrightarrow M^{s_i+q_{i+1}}(l + 1, a_i, k - 1, \dots, l + 2, l, \dots, 0).$$

The long exact sequence for the functor $LZ_{q_{i+1}}^{p_{i+1}}$ gives

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-2} \cdot \alpha) &\cong \\ M^{p_{i+1}}(a_i, k - 1, \dots, 0) &\quad \text{if } s = k - l - 1 \\ 0 &\quad \text{if } s \neq k - l - 1. \end{aligned}$$

Now consider the short exact sequence

$$0 \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-2} \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}}(\sigma_1 \cdots \sigma_{k-l-2} \cdot \alpha) \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-3} \rightarrow 0.$$

This gives rise to a long exact sequence for $LZ_{q_{i+1}}^{p_{i+1}}$. The long exact sequence together with the first two paragraphs of the proof easily give for $s \geq k - l + 1$

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-3} \cong M^{p_{i+1}}(a_i, k - 1, \dots, 0)$$

if $s = k - l + 1, \dots, k + l - 1$ and zero otherwise.

If $s = k - l$ the long exact sequence gives

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-3} \cong 0.$$

If $s = k - l - 1$, the long exact sequence gives

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-3} \cong M^{p_{i+1}}(a_i, k - 1, \dots, 0).$$

If $s < k - l - 1$, then $L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-l-3} \cong 0$.

□

Lemma 14. Suppose $l + 3 \leq t \leq k + 1$. If $s = k - t + 2, k - t + 4, \dots, k + t - 4$ then

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} (L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) / L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1}) \cong M^{p_{i+1}}(a_i, k-1, \dots, 0).$$

Otherwise it is zero.

Proof. We proceed by induction. The base case is lemma 13.

As in lemma 13, we compute

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) &\cong \\ L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (a_1, \dots, a_i, t-2, k-1, \dots, t-1, t-3, \dots, 0, \dots, a_n) &\cong \\ L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} M^{s_i+q_{i+1}} (t-2, a_i, k-1, \dots, t-1, t-3, \dots, 0). \end{aligned}$$

Now consider the short exact sequence

$$\begin{aligned} 0 \rightarrow M^{q_{i+1}}(a_i, t-2, k-1, \dots, t-1, t-3, \dots, 0) \rightarrow \\ M^{q_{i+1}}(t-2, a_i, k-1, \dots, t-1, t-3, \dots, 0) \rightarrow \\ M^{s_i+q_{i+1}}(t-2, a_i, k-1, \dots, t-1, t-3, \dots, 0) \rightarrow 0. \end{aligned}$$

The long exact sequence for $LZ_{q_{i+1}}^{p_{i+1}}$ gives

$$L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \cong M^{p_{i+1}}(a_1, \dots, a_i, k-1, \dots, 0, \dots, a_n)$$

if $s = k - t + 2$ and zero otherwise.

Now consider the short exact sequence

$$0 \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \rightarrow \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \rightarrow 0.$$

This gives rise to a long exact sequence for $LZ_{q_{i+1}}^{p_{i+1}}$.

Now by the induction hypothesis, the long exact sequence and the first paragraph, the lemma is true for $s \geq k - t + 4$.

Suppose $s = k - t + 3$. Then the long exact sequence becomes

$$0 \rightarrow L_{k-t+3} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \rightarrow L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \rightarrow \cdots$$

By the induction hypothesis

$$L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \cong 0,$$

so

$$L_{k-t+3} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \cong 0.$$

Suppose $s = k - t + 2$. Then the long exact sequence becomes

$$\begin{aligned} 0 \rightarrow L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \rightarrow L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \rightarrow \\ L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} \rightarrow L_{k-t+1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \rightarrow \cdots \end{aligned}$$

By the induction hypothesis

$$L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \cong L_{k-t+1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t+1} \cong 0.$$

Thus

$$\begin{aligned} L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} K_{k-t} &\cong L_{k-t+2} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) \\ &\cong M^{p_{i+1}}(a_i, k-1, \dots, 0). \end{aligned}$$

Thus the claim is verified for $s = k - t + 2$.

Finally suppose $s \leq k - t + 1$. Then

$$\begin{aligned} L_s Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) &\cong \\ L_{s-1} Z_{q_{i+1}}^{p_{i+1}} \epsilon_{s_i+q_{i+1}}^{q_{i+1}} L_1 Z_{q_{i+1}}^{s_i+q_{i+1}} M^{q_{i+1}} (\sigma_1 \cdots \sigma_{k-t+1} \cdot \alpha) &\cong 0. \end{aligned}$$

This case is then true by the induction hypothesis. $\square$

Corollary 6. *If $s = 1, 3, \dots, 2(k-1) - 1$ then*

$$L_s Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}} M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0) \cong M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0)$$

and zero otherwise.

Proof. This is a direct consequence of the previous lemma. $\square$

Corollary 7. *For $s = 0, 2, \dots, 2(k-1) - 2$*

$$L_s Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}} M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0) \cong M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0).$$

It is zero otherwise.

Proof. This follows from the previous corollary and noting that

$$L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}} M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0) \cong L_1 Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}} M^{\mathbf{p}_{i+1}}(a_i, k-1, \dots, 0)[1].$$

$\square$

By adjointness, there is an isomorphism

$$\text{Hom}(L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}, L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}) \cong \text{Hom}(\epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}, \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}).$$

Call the image of the identity morphism under this isomorphism ϕ . Then

$$L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \phi: L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}} \rightarrow L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}.$$

By proposition 8, we know $L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}} \cong \bigoplus_{r=0}^{k-1} \text{Id}[2r]$.

Finally we have an isomorphism of functors lifting the third graphical relation.

Proposition 10. *There is an isomorphism*

$$\tau^{\leq k-2} L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}}[-k] \phi: \bigoplus_{r=1}^{k-1} \text{Id}[2r-k] \rightarrow L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}}[-1] L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}} \epsilon_{\mathbf{p}_{i+1}}^{\mathbf{q}_{i+1}}[-(k-1)].$$

Proof. We need to show that the cohomology functors of $\tau^{\leq k-2} L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \phi$ when applied to the generalized Verma module are isomorphisms. In the previous lemmas we needed to compute

$$L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} M^{\mathbf{s}_i + \mathbf{q}_{i+1}}(\underbrace{a_i, a_{i+1}, a_{i+2}, \dots, a_{i+k}}).$$

The calculation was made by considering the short exact sequence

$$0 \rightarrow M^{\mathbf{q}_{i+1}}(a_{i+1}, a_i, \underbrace{a_{i+2}, \dots, a_{i+k}}) \rightarrow M^{\mathbf{q}_{i+1}}(a_i, a_{i+1}, \underbrace{a_{i+2}, \dots, a_{i+k}}) \rightarrow M^{\mathbf{s}_i + \mathbf{q}_{i+1}}(\underbrace{a_i, a_{i+1}, a_{i+2}, \dots, a_{i+k}}) \rightarrow 0.$$

It is clear that if $\{a_{i+1}, \dots, a_{i+k}\} = \{k-1, \dots, 0\}$, then the isomorphism

$$L Z_{\mathbf{q}_{i+1}}^{\mathbf{p}_{i+1}} \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} M^{\mathbf{s}_i + \mathbf{q}_{i+1}}(\underbrace{a_i, a_{i+1}, a_{i+2}, \dots, a_{i+k}}) \cong M^{\mathbf{p}_{i+1}}(a_i, \underbrace{k-1, \dots, 0})[k-1-a_{i+1}]$$

comes from the natural transformation $\text{Id} \rightarrow \epsilon_{\mathbf{s}_i + \mathbf{q}_{i+1}}^{\mathbf{q}_{i+1}} L Z_{\mathbf{q}_{i+1}}^{\mathbf{s}_i + \mathbf{q}_{i+1}}$. If instead it is the case that $\{a_i, a_{i+2}, \dots, a_{i+k}\} = \{k-1, \dots, 0\}$, we need to demonstrate the isomorphism in a different way so that it is natural.

Consider the short exact sequence in the category $\mathcal{O}(\mathfrak{gl}_{k+1})$

$$0 \rightarrow M^{\mathbf{q}_2}(a_{i+1}, a_i, \underbrace{a_{i+2}, \dots, a_{i+k}}) \rightarrow M^{\mathbf{q}_2}(a_i, a_{i+1}, \underbrace{a_{i+2}, \dots, a_{i+k}}) \rightarrow M^{\mathbf{s}_1 + \mathbf{q}_2}(\underbrace{a_i, a_{i+1}, a_{i+2}, \dots, a_{i+k}}) \rightarrow 0.$$

In this category, the objects at the beginning and end of the sequence are self dual with respect to the duality functor d because they are simple. Thus there is an exact sequence

$$M^{\mathbf{s}_1 + \mathbf{q}_2}(\underbrace{a_i, a_{i+1}, a_{i+2}, \dots, a_{i+k}}) \hookrightarrow d M^{\mathbf{q}_2}(a_i, a_{i+1}, \underbrace{a_{i+2}, \dots, a_{i+k}}) \twoheadrightarrow M^{\mathbf{q}_2}(a_{i+1}, a_i, \underbrace{a_{i+2}, \dots, a_{i+k}}).$$

Now we take an external tensor product on the left with the Verma module $M(a_1, \dots, a_{i-1})$ and on the right with $M(a_{i+k+1}, \dots, a_n)$. This will give an exact sequence in $\mathcal{O}(\mathfrak{gl}_{i-1} \oplus \mathfrak{gl}_{k+1} \oplus \mathfrak{gl}_{n-k-i})$. Then we may induce it to the following exact sequence in $\mathcal{O}(\mathfrak{gl}_n)$:

$$M^{\mathbf{s}_1 + \mathbf{q}_{i+1}}(a_1, \dots, a_{i-1}, \underbrace{a_i, a_{i+1}, a_{i+2}, \dots, a_{i+k}}, \dots, a_n) \hookrightarrow X \twoheadrightarrow M^{\mathbf{q}_{i+1}}(a_1, \dots, a_{i-1}, a_{i+1}, a_i, \underbrace{a_{i+2}, \dots, a_{i+k}}, \dots, a_n).$$

The object X is obtained by taking an external tensor product of Verma modules with a dual generalized Verma module followed by parabolic induction. Since $LZ_{\mathfrak{q}_{i+1}}^{\mathfrak{p}_{i+1}}$ naturally commutes with this composition of functors, it suffice to compute

$$LZ_{\mathfrak{q}_2}^{\mathfrak{p}_1} dM^{\mathfrak{q}_2}(a_i, a_{i+1}, \underbrace{a_{i+2}, \dots, a_{i+k}}).$$

This is easily seen to be zero. Thus the isomorphism

$$LZ_{\mathfrak{q}_{i+1}}^{\mathfrak{p}_{i+1}} \epsilon_{\mathfrak{s}_i + \mathfrak{q}_{i+1}}^{\mathfrak{q}_{i+1}} M^{\mathfrak{s}_i + \mathfrak{q}_{i+1}}(\underbrace{a_i, a_{i+1}}, \underbrace{a_{i+2}, \dots, a_{i+k}}) \cong M^{\mathfrak{p}_{i+1}}(a_i, \underbrace{k-1, \dots, 0}[k-1-a_i]$$

also comes from the natural transformation $\text{Id} \rightarrow \epsilon_{\mathfrak{s}_i + \mathfrak{q}_{i+1}}^{\mathfrak{q}_{i+1}} LZ_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}}$.

Therefore the natural transformation stated in the proposition is an isomorphism for generalized Verma modules. Then it must be true for any projective object by induction on the length of its generalized Verma flag. The theorem is then true for an arbitrary objects by considering a projective resolution of it. $\square$

4.4. Diagram 4. For ease of notation, we abbreviate the algebras defined before for this subsection as follows.

Definition 12. Let

- (1) $\alpha = \mathfrak{p}_i + \mathfrak{q}_{i+k}$
- (2) $\beta = \mathfrak{q}_{i-1} + \mathfrak{q}_{i+k}$
- (3) $\gamma = \mathfrak{q}_{i-1} + \mathfrak{s}_{i+k-1} + \mathfrak{q}_{i+k}$
- (4) $\delta = \mathfrak{q}_{i-1} + \mathfrak{p}_{i+k}$.

We begin with a generalized Verma module

$$M^\alpha(a_1, \dots, a_{i-1}, \underbrace{k-1, \dots, 0}, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}, a_{i+2k}, \dots, a_n).$$

Since the coordinates $a_1, \dots, a_{i-1}, a_{i+2k}, \dots, a_n$ play no role in the computation, we omit them when labeling generalized Verma modules. Suppose $a_{i+k} = l$. Let us denote the module above by $M^\alpha(l)$.

There are short exact sequences arising from the generalized BGG resolution:

$$\begin{aligned} 0 \rightarrow K_0 \rightarrow M^\beta(\underbrace{k-1, \dots, \hat{0}}, 0, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow M^\alpha(l) \rightarrow 0 \\ 0 \rightarrow K_1 \rightarrow M^\beta(\underbrace{k-1, \dots, \hat{1}}, 0, 1, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow K_0 \rightarrow 0 \\ \dots \\ 0 \rightarrow K_{l-1} \rightarrow M^\beta(\underbrace{k-1, \dots, \widehat{l-1}}, 0, l-1, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow K_{l-2} \rightarrow 0 \\ 0 \rightarrow K_l \rightarrow M^\beta(\underbrace{k-1, \dots, \hat{l}}, 0, l, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow K_{l-1} \rightarrow 0 \\ 0 \rightarrow K_{l+1} \rightarrow M^\beta(\underbrace{k-1, \dots, \widehat{l+1}}, 0, l+1, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow K_l \rightarrow 0 \\ \dots \\ 0 \rightarrow K_{k-1} \rightarrow M^\beta(\underbrace{\widehat{k-1}, \dots, 0}, k-1, a_{i+k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow K_{k-2} \rightarrow 0. \end{aligned}$$

Lemma 15. *There are exact sequences:*

$$\begin{aligned} 0 \rightarrow L_1 Z_\beta^\gamma K_0 \rightarrow M^\gamma(\underbrace{k-1, \dots, \hat{0}}, \underbrace{a_{i+k}, 0}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow L_1 Z_\beta^\alpha \epsilon_\alpha^\beta M^\alpha(l) \rightarrow 0 \\ 0 \rightarrow L_1 Z_\beta^\gamma K_1 \rightarrow M^\gamma(\underbrace{k-1, \dots, \hat{1}}, \underbrace{0, 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow L_1 Z_\beta^\gamma K_0 \rightarrow 0 \\ \dots \\ 0 \rightarrow L_1 Z_\beta^\gamma K_{l-1} \rightarrow M^\gamma(\underbrace{k-1, \dots, \widehat{l-1}}, \underbrace{0, l-1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow L_1 Z_\beta^\gamma K_{l-2} \rightarrow 0 \\ L_i Z_\beta^\gamma K_l = L_{i+1} Z_\beta^\gamma K_{l-1}, \forall i \\ 0 \rightarrow L_0 Z_\beta^\gamma K_{l+1} \rightarrow M^\gamma(\underbrace{k-1, \dots, \widehat{l+1}}, \underbrace{0, l+1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}) \rightarrow L_0 Z_\beta^\gamma K_l \rightarrow 0 \\ \dots \end{aligned}$$

$$0 \rightarrow L_0 Z_\beta^\gamma K_{k-1} \rightarrow M^\gamma(\underbrace{k-1, \dots, 0}_{\text{width } k}, \underbrace{k-1, a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}) \rightarrow L_0 Z_\beta^\gamma K_{k-2} \rightarrow 0$$

Proof. In the first set of sequences, $L_0 Z_\beta^\gamma$ must be zero. Also, $L_1 Z_\beta^\gamma K_l = L_2 Z_\beta^\gamma K_{l-1} = 0$. Then $L_1 Z_\beta^\gamma K_{l+1} = L_2 Z_\beta^\gamma K_l = L_3 Z_\beta^\gamma K_{l-1} = 0$. We also have $L_1 Z_\beta^\gamma K_{l+2} = L_2 Z_\beta^\gamma K_{l+1} = 0$ and so on. This gives the second set of exact sequences. $\square$

Corollary 8. *There are exact sequences:*

$$\begin{aligned} 0 \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_0 &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{k-1, \dots, \hat{0}, a_{i+k}, 0}_{\text{width } k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}) \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\alpha \epsilon_\alpha^\beta M^\alpha(l) \rightarrow 0 \\ 0 \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_1 &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{k-1, \dots, \hat{1}, 0, a_{i+k}, 1}_{\text{width } k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}) \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_0 \rightarrow 0 \\ &\dots \\ 0 \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-1} &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{k-1, \dots, \widehat{l-1}, 0, a_{i+k}, l-1}_{\text{width } k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}) \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-2} \rightarrow 0 \\ &\epsilon_\gamma^\beta L_i Z_\beta^\gamma K_l = \epsilon_\gamma^\beta L_{i+1} Z_\beta^\gamma K_{l-1}, \forall i \\ 0 \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma K_{l+1} &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{k-1, \dots, \widehat{l+1}, 0, l+1, a_{i+k}}_{\text{width } k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}) \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma K_l \rightarrow 0 \\ &\dots \\ 0 \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma K_{k-1} &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{k-1, \dots, \widehat{0}, k-1, a_{i+k}}_{\text{width } k}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}) \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma K_{k-2} \rightarrow 0 \end{aligned}$$

Lemma 16. *Let $\{a_{i+k+1}, \dots, a_{i+2k-1}\} = \{k-1, \dots, \hat{m}, \dots, 0\}$. Let $m \neq l$. If $s = k-1$, then*

$$L_s Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma \epsilon_\alpha^\beta M^\alpha = M^\delta(\underbrace{k-1, \dots, \hat{m}, \dots, 0}_{\text{width } k}, \underbrace{a_{i+k}, k-1, \dots, 0}_{\text{width } k}).$$

Otherwise it is zero.

Proof. This follows easily from the previous corollary. $\square$

Now consider

$$M^\delta = M^\delta(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{a_{i+k-1}, k-1}_{\text{width } 2}, \underbrace{k-1, \dots, 0}_{\text{width } k-1})$$

and let $a_{i+k-1} = l$. There are short exact sequences

$$\begin{aligned} 0 \rightarrow J_0 &\rightarrow M^\beta(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{a_{i+k-1}, k-1}_{\text{width } 2}, \underbrace{k-1, \widehat{k-1}, \dots, 0}_{\text{width } k-1}) \rightarrow M^\delta(l) \rightarrow 0 \\ 0 \rightarrow J_1 &\rightarrow M^\beta(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{a_{i+k-1}, k-2}_{\text{width } 2}, \underbrace{k-1, \widehat{k-2}, \dots, 0}_{\text{width } k-1}) \rightarrow J_0 \rightarrow 0 \\ &\dots \\ 0 \rightarrow J_{k-1} &\rightarrow M^\beta(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{a_{i+k-1}, 0}_{\text{width } 2}, \underbrace{k-1, \dots, \hat{0}}_{\text{width } k-1}) \rightarrow J_{k-2} \rightarrow 0. \end{aligned}$$

Corollary 9. *There are exact sequences:*

$$\begin{aligned} 0 \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_0 &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{k-1, a_{i+k-1}}_{\text{width } 2}, \underbrace{\widehat{k-1}, \dots, 0}_{\text{width } k-1}) \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma M^\delta(l) \rightarrow 0 \\ 0 \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_1 &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{k-2, a_{i+k-1}}_{\text{width } 2}, \underbrace{k-1, \widehat{k-2}, \dots, 0}_{\text{width } k-1}) \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_0 \rightarrow 0 \\ &\dots \\ 0 \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-2} &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{l+1, a_{i+k-1}}_{\text{width } 2}, \underbrace{k-1, \widehat{l+1}, \dots, 0}_{\text{width } k-1}) \rightarrow \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-3} \rightarrow 0 \\ &\epsilon_\gamma^\beta L_i Z_\beta^\gamma J_{k-l-1} \cong \epsilon_\gamma^\beta L_{i+1} Z_\beta^\gamma J_{k-l-2}, \forall i \\ 0 \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-l} &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{a_{i+k-1}, l-1}_{\text{width } 2}, \underbrace{k-1, \widehat{l-1}, \dots, 0}_{\text{width } k-1}) \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-l-1} \rightarrow 0 \\ &\dots \\ 0 \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-1} &\rightarrow \epsilon_\gamma^\beta M^\gamma(\underbrace{a_i, \dots, a_{i+k-2}}_{\text{width } k-1}, \underbrace{a_{i+k-1}, 0}_{\text{width } 2}, \underbrace{k-1, \dots, \hat{0}}_{\text{width } k-1}) \rightarrow \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-2} \rightarrow 0 \end{aligned}$$

Proof. This is similar to corollary 8. $\square$

Lemma 17. Let $\{a_i, \dots, a_{i+k-2}\} = \{k-1, \dots, \hat{m}, \dots, 0\}$. Let $m \neq l$. If $s = k-1$, then

$$L_s Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma \epsilon_\delta^\beta M^\delta(l) = M^\alpha(\underbrace{k-1, \dots, 0}_{a_{i+k-1}}, \underbrace{k-1, \dots, \hat{m}, \dots, 0}_{\dots}).$$

Otherwise it is zero.

Proof. This follows from corollary 9. □

We must now study the case when $l = m$.

Lemma 18. Let $m = l$. Then $L_s Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma \epsilon_\alpha^\beta M^\alpha(l) \cong$

$$\begin{aligned} M^\delta(0)/\dots/M^\delta(k-1) & \quad \text{if } s = k-l \\ L_{k-l+1} Z_\beta^\gamma \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} & \quad \text{if } s = k-1 \end{aligned}$$

and there is an exact sequence

$$0 \rightarrow M^\delta(l)/\dots/M^\delta(k-1) \rightarrow L_{k-l+1} Z_\beta^\gamma \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} \rightarrow M^\delta(l+1)/\dots/M^\delta(k-1) \rightarrow 0.$$

Proof. From corollary 8 we get $L_s Z_\beta^\delta \epsilon_\gamma^\beta L_0 Z_\beta^\gamma K_{k-2} \cong$

$$\begin{aligned} M^\delta(k-1) & \quad \text{if } s = k-l-1 \\ 0 & \quad \text{if } s \neq k-l-1. \end{aligned}$$

Continuing in this way and using the corollary we get $L_s Z_\beta^\delta \epsilon_\gamma^\beta L_0 Z_\beta^\gamma K_l \cong$

$$\begin{aligned} M^\delta(l+1)/\dots/M^\delta(k-1) & \quad \text{if } s = k-l-1 \\ 0 & \quad \text{if } s \neq k-l-1 \end{aligned}$$

and then $L_s Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-1} \cong$

$$\begin{aligned} M^\delta(l+1)/\dots/M^\delta(k-1) & \quad \text{if } s = k-l-1 \\ 0 & \quad \text{if } s \neq k-l-1. \end{aligned}$$

Corollary 8 then gives an exact sequence

$$0 \rightarrow M^\delta(l-1) \rightarrow L_{k-l} Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-2} \rightarrow M^\delta(l+1)/\dots/M^\delta(k-1) \rightarrow 0.$$

Next the corollary gives the following diagram:

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ & & M^\delta(l-1) & & & & \\ & & \downarrow & & & & \\ 0 & \longrightarrow & L_{k-l+1} Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} & \longrightarrow & L_{k-l} Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-2} & \longrightarrow & M^\delta(l-2) \longrightarrow L_{k-l} Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} \longrightarrow 0 \\ & & & & \downarrow & & \\ & & & & M^\delta(l+1)/\dots/M^\delta(k-1) & & \\ & & & & \downarrow & & \\ & & & & 0 & & \end{array}$$

Since that the map $M^\delta(l-1) \rightarrow M^\delta(l-2)$ is standard, the above diagram can be extended to the following commutative diagram.

$$\begin{array}{ccccccc}
& 0 & & 0 & & 0 & \\
& \downarrow & & \downarrow & & \downarrow & \\
& M^\delta(l)/\dots/M^\delta(k-1) & \longrightarrow & M^\delta(l-1) & \longrightarrow & M^\delta(l-2) & \\
& \downarrow & & \downarrow & & \downarrow & \\
0 \longrightarrow & L_{k-l+1}Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} & \longrightarrow & L_{k-l}Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-2} & \longrightarrow & M^\delta(l-2) & \longrightarrow L_{k-l}Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} \longrightarrow 0 \\
& \downarrow & & \downarrow & & \downarrow & \\
0 \longrightarrow & M^\delta(l+1)/\dots/M^\delta(k-1) & \longrightarrow & M^\delta(l+1)/\dots/M^\delta(k-1) & \longrightarrow & 0 & \\
& \downarrow & & \downarrow & & \downarrow & \\
& 0 & & 0 & & 0 &
\end{array}$$

The surjection in the first column arises from the snake lemma and formulas for dimensions of spaces of homomorphisms between generalized Verma modules given in [Sh]. It also implies that $L_{k-l}Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} \cong M^\delta(l-2)/\dots/M^\delta(k-1)$ and there is an exact sequence

$$0 \rightarrow M^\delta(l)/\dots/M^\delta(k-1) \rightarrow L_{k-l+1}Z_\beta^\delta \epsilon_\gamma^\beta L_1 Z_\beta^\gamma K_{l-3} \rightarrow M^\delta(l+1)/\dots/M^\delta(k-1) \rightarrow 0.$$

Continuing in this manner and using corollary 8 gives us the lemma. $\square$

Lemma 19. *Let $m = l$. Then $L_s Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma \epsilon_\delta^\beta M^\delta(l) \cong$*

$$\begin{aligned}
& M^\alpha(k-1)/\dots/M^\alpha(0) & \text{if } s = l+1 \\
& L_{l+2}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-4} & \text{if } s = k-1
\end{aligned}$$

and there is an exact sequence

$$0 \rightarrow M^\alpha(l)/\dots/M^\alpha(0) \rightarrow L_{l+2}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-4} \rightarrow M^\alpha(l-1)/\dots/M^\alpha(0) \rightarrow 0.$$

Proof. We use the exact sequences of corollary 9.

If $s = l$, $L_s Z_\beta^\alpha \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-2} \cong M^\alpha(0)$ and is zero otherwise.

If $s = l$, $L_s Z_\beta^\alpha \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-3} \cong M^\alpha(1)/M^\alpha(0)$ and is zero otherwise.

Continuing in this way,

If $s = l$, $L_s Z_\beta^\alpha \epsilon_\gamma^\beta L_0 Z_\beta^\gamma J_{k-l-1} \cong M^\alpha(l-1)/\dots/M^\alpha(0)$ and is zero otherwise.

Next we get an exact sequence

$$0 \rightarrow M^\alpha(l+1) \rightarrow L_{l+1}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-3} \rightarrow M^\alpha(l-1)/\dots/M^\alpha(0) \rightarrow 0.$$

Corollary 9 then produces the following diagram:

$$\begin{array}{ccccccc}
& 0 & & & & & \\
& \downarrow & & & & & \\
& M^\alpha(l+1) & & & & & \\
& \downarrow & & & & & \\
0 \longrightarrow & L_{l+2}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-4} & \longrightarrow & L_{l+1}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-3} & \longrightarrow & M^\alpha(l+2) & \longrightarrow L_{l+1}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-4} \longrightarrow 0 \\
& \downarrow & & & & & \\
& M^\alpha(l-1)/\dots/M^\alpha(0) & & & & & \\
& \downarrow & & & & & \\
& 0 & & & & &
\end{array}$$

Thus $L_{l+1}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-4} \cong M^\alpha(l+2)/\dots/M^\alpha(0)$ and there is an exact sequence

$$0 \rightarrow M^\alpha(l)/\dots/M^\alpha(0) \rightarrow L_{l+2}Z_\beta^\alpha \epsilon_\gamma^\beta L_1 Z_\beta^\gamma J_{k-l-4} \rightarrow M^\alpha(l-1)/\dots/M^\alpha(0) \rightarrow 0.$$

Using the rest of the exact sequences of corollary 9, we easily obtain this lemma. (The details are the same as lemma 18.) $\square$

For the rest of this subsection, let $F = LZ_{\beta}^{\alpha}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}\epsilon_{\delta}^{\beta}$. Let $G = LZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}$.

Lemma 20. *Let $l < k - 1$.*

- (1) $H^{-s}FM^{\delta}(l)/\dots/M^{\delta}(k-1) \cong$

$$\begin{array}{ll} M^{\alpha}(k-1)/\dots/M^{\alpha}(0) & \text{if } s = l+1, l+3, \dots, 2(k-1) - l - 1 \\ M^{\alpha}(l-1)/\dots/M^{\alpha}(0) & \text{if } s = k-1. \end{array}$$
- (2) $H^{-s}GM^{\alpha}(l)/\dots/M^{\alpha}(0) \cong$

$$\begin{array}{ll} M^{\delta}(0)/\dots/M^{\delta}(k-1) & \text{if } s = k-l, k-l+2, \dots, k+l-2 \\ M^{\delta}(l+1)/\dots/M^{\delta}(k-1) & \text{if } s = k-1. \end{array}$$

Proof. (1) Consider the exact sequence

$$0 \rightarrow M^{\delta}(k-1) \rightarrow \dots \rightarrow M^{\delta}(l+1) \rightarrow M^{\delta}(l) \rightarrow M^{\delta}(l)/\dots/M^{\delta}(k-1) \rightarrow 0.$$

By lemma 19, applying F gives the following complex with vertical short exact sequences:

$$\begin{array}{ccccccc} 0 & & 0 & & 0 & & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ M^{\alpha}(k-1)/\dots/M^{\alpha}(0) & & M^{\alpha}(l+1)/\dots/M^{\alpha}(0) & & M^{\alpha}(l)/\dots/M^{\alpha}(0) & & M^{\alpha}(l)/\dots/M^{\alpha}(0) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ L_{k-1}FM^{\delta}(k-1) & \xrightarrow{\quad} & L_{k-1}FM^{\delta}(l+1) & \xrightarrow{\quad} & L_{k-1}FM^{\delta}(l) & \xrightarrow{\quad} & L_{k-1}FM^{\delta}(l)/\dots/M^{\delta}(k-1) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ M^{\alpha}(k-2)/\dots/M^{\alpha}(0) & & M^{\alpha}(l)/\dots/M^{\alpha}(0) & & M^{\alpha}(l-1)/\dots/M^{\alpha}(0) & & M^{\alpha}(l-1)/\dots/M^{\alpha}(0) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & & 0 & & 0 & & 0 \end{array}$$

It suffices to show that the map $L_{k-1}FM^{\delta}(l+1) \rightarrow L_{k-1}FM^{\delta}(l)$ factors through $M^{\alpha}(l)/\dots/M^{\alpha}(0)$. Suppose that the generalized BGG resolutions for $M^{\delta}(l+1)$ and $M^{\delta}(l)$ give rise to short exact sequences involving modules B'_i and B_i respectively. Let $M^{\beta}(r, s) = M^{\beta}(\underbrace{a_i, \dots, a_{i+k-2}}_{}, r, s, \underbrace{k-1, \dots, \hat{s}, \dots, 0}_{})$.

There is a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & B'_{k-l-3} & \longrightarrow & M^{\beta}(l+1, l+2) & \longrightarrow & B'_{k-l-4} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & B_{k-l-3} & \longrightarrow & M^{\beta}(l, l+2) & \longrightarrow & B_{k-l-4} \longrightarrow 0 \end{array}$$

Now apply F to get the following commutative diagram:

$$\begin{array}{ccccc} L_{l+2}FB'_{k-l-4} & \longrightarrow & L_{l+1}FB'_{k-l-3} \cong M^{\alpha}(l)/\dots/M^{\alpha}(0) & \longrightarrow & 0 \\ \downarrow f & & \downarrow i & & \downarrow \\ L_{l+2}FB_{k-l-4} & \xrightarrow{h} & L_{l+1}FB_{k-l-3} & \xrightarrow{j} & M^{\alpha}(l+2) \end{array}$$

Thus $\text{im}(i) \subset \ker(j) = \text{im}(h) = L_{l+2}FB_{k-l-4}$. Thus the map $L_{k-1}FM^{\delta}(l+1) \rightarrow L_{k-1}FM^{\delta}(l)$ factors through $M^{\alpha}(l)/\dots/M^{\alpha}(0)$ as desired.

Now the long exact sequence for F gives $H^{-s}FM^{\delta}(l)/\dots/M^{\delta}(k-1) \cong$

$$\begin{array}{ll} M^{\alpha}(k-1)/\dots/M^{\alpha}(0) & \text{if } s = l+1, l+3, \dots, 2(k-1) - l - 1 \\ M^{\alpha}(l-1)/\dots/M^{\alpha}(0) & \text{if } s = k-1. \end{array}$$

(2) This follows exactly as the first part. □

Lemma 21. $H^{-s}FLZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}M^{\alpha}(l) \cong$

$$\begin{aligned} M^{\alpha}(k-1)/\dots/M^{\delta}(0) & \quad \text{if } s = k-l+1, k-l+3, \dots, k-l+2(k-1)-3 \\ M^{\alpha}(l) & \quad \text{if } s = 2(k-1). \end{aligned}$$

Proof. By lemma 18, there is a distinguished triangle

$$L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1] \rightarrow LZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}M^{\alpha}(l) \rightarrow M^{\delta}(0)/\dots/M^{\delta}(k-1)[k-l].$$

By proposition 10, $H^{-s}FM^{\delta}(0)/\dots/M^{\delta}(k-1)[k-l] \cong M^{\alpha}(k-1)/\dots/M^{\alpha}(0)$ for $s = k-l+1, k-l+3, \dots, k-l+2(k-1)-1$.

From the commutative diagram of lemma 18, we have the short exact sequences:

$$\begin{aligned} 0 \rightarrow L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} & \rightarrow L_{k-l}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-2} \rightarrow M^{\delta}(l-1)/\dots/M^{\delta}(k-1) \rightarrow 0 \\ 0 \rightarrow M^{\delta}(l-1)/\dots/M^{\delta}(k-1) & \rightarrow M^{\delta}(l-2) \rightarrow M^{\delta}(l-2)/\dots/M^{\delta}(k-1) \rightarrow 0. \end{aligned}$$

Therefore we get the following morphisms:

$$M^{\delta}(0)/\dots/M^{\delta}(k-1) \rightarrow M^{\delta}(l-1)/\dots/M^{\delta}(k-1)[l-1] \rightarrow L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[l].$$

This is the morphism $M^{\delta}(0)/\dots/M^{\delta}(k-1)[k-l] \rightarrow L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1]$ in the distinguished triangle. Assume $s \leq k-l+2(k-1)-3$. Suppose

$$H^{-s}F(M^{\delta}(l-1)/\dots/M^{\delta}(k-1)[k-1]) \rightarrow H^{-s+1}F(L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1])$$

is an isomorphism. Then the long exact sequence for the triangle above and F gives a map:

$$H^{-s}FM^{\delta}(0)/\dots/M^{\delta}(k-1)[k-l] \rightarrow H^{-s+1}F(L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1])$$

which is an isomorphism. Then we also have

$$H^{-s}F(L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1]) \cong M^{\alpha}(k-1)/\dots/M^{\alpha}(0).$$

If the map is zero, then

$$H^{-s}F(L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1]) \cong H^{-s+1}F(L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1]) = 0.$$

Therefore in either case,

$$H^{-s}FLZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}M^{\alpha}(l) \cong M^{\alpha}(k-1)/\dots/M^{\delta}(0)$$

for $s = k-l+1, k-l+3, \dots, k-l+2(k-1)-3$.

For $s = 2(k-1)-l-1$, consider again the short exact sequence

$$0 \rightarrow L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \rightarrow L_{k-l}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-2} \rightarrow M^{\delta}(l-1)/\dots/M^{\delta}(k-1) \rightarrow 0.$$

It is easy to calculate

$$H^{-(2(k-1)-l-1)}FL_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \cong M^{\alpha}(k-1)/\dots/M^{\alpha}(0)$$

and

$$H^{-(2(k-1)-l-1)}FL_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-2} \cong H^{-(2(k-1)-l)}FL_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-2} = 0.$$

Thus

$$H^{-(2(k-1)-l)}FM^{\delta}(l-1)/\dots/M^{\delta}(k-1) \rightarrow H^{-(2(k-1)-l-1)}FL_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}$$

is an isomorphism. Therefore

$$H^{-(k-l+2(k-1))}FM^{\delta}(0)/\dots/M^{\delta}(k-1)[k-l] \rightarrow H^{-(k-l+2(k-1)-1)}FL_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1]$$

is an isomorphism so

$$H^{-(k-l+2(k-1)-1)}FLZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}M^{\alpha}(l) = 0.$$

In order to compute $H^{-s}F(L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}[k-1])$ for $s = 2(k-1)$, we use the short exact sequence from lemma 18:

$$0 \rightarrow M^{\delta}(l)/\dots/M^{\delta}(k-1) \rightarrow L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \rightarrow M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow 0.$$

The first goal is to prove that this sequence is non-split. Let $\bar{F} = LZ_{\beta}^{\alpha}\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma}\epsilon_{\delta}^{\beta}[-(k-1)]$. Let the adjoint of $\bar{F}$ be $\bar{G} = LZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}[-(k-1)]$. Then the exact sequence becomes

$$0 \rightarrow M^{\delta}(l)/\dots/M^{\delta}(k-1) \rightarrow H^0\bar{G}M^{\alpha}(l) \rightarrow M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow 0.$$

In order to show that it is non-split, it suffices to show that $\text{Hom}(M^{\delta}(l+1)/\dots/M^{\delta}(k-1), H^0\bar{G}M^{\alpha}(l)) = 0$. Consider the distinguished triangle

$$H^0\bar{G}M^{\alpha}(l) \rightarrow \bar{G}M^{\alpha}(l) \rightarrow M^{\delta}(0)/\dots/M^{\delta}(k-1)[-l+1].$$

It then suffices to prove

$$\text{Hom}(M^{\delta}(l+1)/\dots/M^{\delta}(k-1), \bar{G}M^{\alpha}(l)) \cong \text{Hom}(\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1), M^{\alpha}(l)) = 0.$$

For this consider the distinguished triangle

$$\tau^{\leq 0}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow \bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow \tau^{\geq 1}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1).$$

We will show that this triangle splits. Note that $M^{\alpha}(k-1)/\dots/M^{\alpha}(0) \cong \epsilon_{\alpha+\delta}^{\alpha}M^{\alpha+\delta}(\underbrace{k-1, \dots, 0}_{\text{---}}, \underbrace{k-1, \dots, 0}_{\text{---}})$.

Denote it simply by $M^{\alpha+\delta}$. Assume $l-k$ is even. The other case follows in the same way as below. Clearly,

$$\tau^{\geq 1}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \cong M^{\alpha+\delta}[-k+l+3] \oplus M^{\alpha+\delta}[-k+l+5] \oplus \dots \oplus M^{\alpha+\delta}[-1].$$

Next consider $\tau^{\leq 0}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1)$. There is a distinguished triangle:

$$\begin{aligned} \tau^{\leq -1}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) &\rightarrow \tau^{\leq 0}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow \\ H^0\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) &\cong M^{\alpha}(l)/\dots/M^{\alpha}(0). \end{aligned}$$

By lemma 20,

$$\tau^{\leq -1}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \cong M^{\alpha+\delta}[1] \oplus M^{\alpha+\delta}[3] \oplus \dots \oplus M^{\alpha+\delta}[k-l-3].$$

We easily compute

$$\text{Hom}(M^{\alpha}(l)/\dots/M^{\alpha}(0), \tau^{\leq -1}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1)[1]) = 0.$$

Thus

$$\tau^{\leq 0}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \cong M^{\alpha+\delta}[1] \oplus \dots \oplus M^{\alpha+\delta}[k-l-3] \oplus M^{\alpha}(l)/\dots/M^{\alpha}(0).$$

Next we check that

$$\text{Hom}(\tau^{\geq 1}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1), \tau^{\leq 0}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1)[1]) = 0.$$

This reduces to showing

$$\text{Hom}(M^{\alpha+\delta}[-k+l+3] \oplus M^{\alpha+\delta}[-k+l+5] \oplus \dots \oplus M^{\alpha+\delta}[-1], M^{\alpha}(l)/\dots/M^{\alpha}(0)[1]) = 0.$$

Then the above space of homomorphisms becomes by adjointness:

$$\text{Hom}(M^{\alpha+\delta}[-k+l+3] \oplus \dots \oplus M^{\alpha+\delta}[-1], LZ_{\alpha}^{\alpha+\delta}M^{\alpha}(l)/\dots/M^{\alpha}(0)[-2k+3]).$$

It is easy to calculate

$$LZ_{\alpha}^{\alpha+\delta}M^{\alpha}(l)/\dots/M^{\alpha}(0) \cong M^{\alpha+\delta}[k-l-1] \oplus \dots \oplus M^{\alpha+\delta}[3k-l-3].$$

Therefore the space of homomorphisms is zero so

$$\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \cong M^{\alpha}(l)/\dots/M^{\alpha}(0) \oplus M^{\alpha+\delta}[-k+l+3] \oplus \dots \oplus M^{\alpha+\delta}[k-l-3].$$

It is now easy to see that

$$\text{Hom}(\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1), M^{\alpha}(l)) = 0.$$

Therefore

$$0 \rightarrow M^{\delta}(l)/\dots/M^{\delta}(k-1) \rightarrow L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \rightarrow M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow 0$$

is non-split.

We want to show that when $\bar{F}$ is applied to the above sequence, it remains non-split. Since $\bar{F}$ and $\bar{G}$ are biadjoint, the composition $\bar{G} \rightarrow \bar{G}\bar{F}\bar{G} \rightarrow \bar{G}$ is the identity. This gives that the adjunction morphism induces an isomorphism on cohomology:

$$M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow H^0\bar{G}\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1).$$

Therefore, by the five lemma, the adjunction morphism induces an isomorphism:

$$L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \rightarrow H^0\bar{G}\bar{F}L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3}.$$

This implies that

$$0 \rightarrow H^0\bar{F}M^{\delta}(l)/\dots/M^{\delta}(k-1) \rightarrow H^0\bar{F}L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \rightarrow H^0\bar{F}M^{\delta}(l+1)/\dots/M^{\delta}(k-1) \rightarrow 0$$

is non-split. This is the exact sequence

$$0 \rightarrow M^{\alpha}(l-1)/\dots/M^{\alpha}(0) \rightarrow H^0\bar{F}L_{k-l+1}Z_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}L_1Z_{\beta}^{\gamma}K_{l-3} \rightarrow M^{\alpha}(l)/\dots/M^{\alpha}(0) \rightarrow 0.$$

If the extension is unique up to isomorphism, the middle term is isomorphic to $M^{\alpha}(l)$. This would complete the lemma.

Consider the truncated full subcategory where $M^{\alpha}(l)$ is a projective object. Apply the functor $\text{Hom}(M^{\alpha}(l), \bullet)$ to the short exact sequence

$$0 \rightarrow M^{\alpha}(l-1)/\dots/M^{\alpha}(0) \rightarrow M^{\alpha}(l) \rightarrow M^{\alpha}(l)/\dots/M^{\alpha}(0) \rightarrow 0$$

in the truncated subcategory to get the exact sequence

$$0 \rightarrow \text{Hom}(M^{\alpha}(l), M^{\alpha}(l-1)/\dots/M^{\alpha}(0)) \rightarrow \text{Hom}(M^{\alpha}(l), M^{\alpha}(l)) \rightarrow \text{Hom}(M^{\alpha}(l), M^{\alpha}(l)/\dots/M^{\alpha}(0)) \rightarrow 0.$$

Thus $\text{Hom}(M^{\alpha}(l), M^{\alpha}(l)/\dots/M^{\alpha}(0)) \cong \mathbb{C}$ and $\text{Ext}^1(M^{\alpha}(l), M^{\alpha}(l-1)/\dots/M^{\alpha}(0)) = 0$. If we apply the functor $\text{Hom}(\bullet, M^{\alpha}(l)/\dots/M^{\alpha}(0))$ to the short exact sequence

$$0 \rightarrow M^{\alpha}(l-1)/\dots/M^{\alpha}(0) \rightarrow M^{\alpha}(l) \rightarrow M^{\alpha}(l)/\dots/M^{\alpha}(0) \rightarrow 0,$$

we get the exact sequence

$$0 \rightarrow \text{End}(M^{\alpha}(l)/\dots/M^{\alpha}(0)) \rightarrow \text{Hom}(M^{\alpha}(l), M^{\alpha}(l)/\dots/M^{\alpha}(0)) \rightarrow \dots$$

This implies

$$\text{End}(M^{\alpha}(l)/\dots/M^{\alpha}(0)) \cong \mathbb{C}.$$

Now apply the functor $\text{Hom}(\bullet, M^{\alpha}(l-1)/\dots/M^{\alpha}(0))$ to the short exact sequence

$$0 \rightarrow M^{\alpha}(l-1)/\dots/M^{\alpha}(0) \rightarrow M^{\alpha}(l) \rightarrow M^{\alpha}(l)/\dots/M^{\alpha}(0) \rightarrow 0.$$

This gives a long exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \text{Ext}^1(M^{\alpha}(l)/\dots/M^{\alpha}(0), M^{\alpha}(l-1)/\dots/M^{\alpha}(0)) \rightarrow \text{Ext}^1(M^{\alpha}(l), M^{\alpha}(l-1)/\dots/M^{\alpha}(0)) = 0.$$

Therefore $\text{Ext}^1(M^{\alpha}(l)/\dots/M^{\alpha}(0), M^{\alpha}(l-1)/\dots/M^{\alpha}(0)) \cong \mathbb{C}$ and we have shown that the extension is unique. □

For the rest of this subsection we make the following definition.

Definition 13. Let $S = LZ_{\beta}^{\alpha}\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma}\epsilon_{\delta}^{\beta}[-(k-1)]LZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}[-(k-1)]$.

The identity morphism in $\text{End}(LZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}[-(k-1)])$ gives rise to a morphism $S \rightarrow \text{Id}$. There is an adjunction morphism $\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma} \rightarrow \text{Id}[1]$ that gives rise to a morphism

$$\theta: S \rightarrow LZ_{\beta}^{\alpha}\epsilon_{\delta}^{\beta}[-(k-1)]LZ_{\beta}^{\delta}\epsilon_{\gamma}^{\beta}[-1]LZ_{\beta}^{\gamma}\epsilon_{\alpha}^{\beta}[-(k-1)][1].$$

By proposition 10, this is a map

$$\theta: S \rightarrow \oplus_{r=1}^{k-1}\epsilon_{\alpha+\delta}^{\alpha}[-(k-1)]LZ_{\alpha}^{\alpha+\delta}[1][2r-k].$$

Composing this map with projection gives a map:

$$\mu_2: S \rightarrow \oplus_{r=1}^{k-2}\epsilon_{\alpha+\delta}^{\alpha}[-(k-1)]LZ_{\alpha}^{\alpha+\delta}[2r-(k-1)].$$

Proposition 11. The map $\mu_1 \oplus \mu_2: S \rightarrow \text{Id} \oplus \oplus_{r=1}^{k-2}\epsilon_{\alpha+\delta}^{\alpha+\beta}[-(k-1)]LZ_{\alpha+\beta}^{\alpha+\delta}[2r-(k-1)]$ is an isomorphism.

Proof. This morphism is an isomorphism for generalized Verma modules as in the proof of proposition 10. By considering a generalized Verma flag for a projective object, we see that $\mu_1 \oplus \mu_2$ is an isomorphism for projectives as well. Since any bounded complex is quasi-isomorphic to a complex of projective objects, $\mu_1 \oplus \mu_2$ must be an isomorphism. □

4.5. Diagram 5. Recall the definitions of the subalgebras $\mathfrak{s}_i$ and $\mathfrak{t}_i$. We would like to understand the functor $LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1]$. In [BFK] this functor turned out to be the identity. When working on maximally singular categories, $LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1]$ was an equivalence of categories. On these less singular blocks this will turn out not to be the case. It will be the identity plus some correction term which will depend on the two minimal parabolics involved. The plan is to compute this functor on generalized Verma modules and then to construct some natural transformations which are isomorphisms on the generalized Verma modules. These objects are parametrized by highest weights which satisfy the conditions of lying in the block and locally finite with respect to the appropriate subalgebra. Since in these computations only $\mathfrak{s}_i, \mathfrak{s}_{i+1}$, and $\mathfrak{t}_i$ are involved, for convenience of notation we will omit all the coordinates of the highest weight except the components in positions $i, i+1$, and $i+2$.

Lemma 22. *Let $a_i > a_{i+1} = a_{i+2}$. Then*

$$LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

Proof. There is a short exact sequence

$$0 \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

Thus $L_s Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong 0$ if $s = 0, 2$ and is isomorphic to $M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_i, a_{i+2}})$ if $s = 1$.

Now consider the short exact sequence

$$0 \rightarrow M(a_{i+1}, a_{i+2}, a_i) \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_i, a_{i+2}}) \rightarrow 0.$$

This gives $L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_i, a_{i+2}}) \cong 0$ if $s = 0, 2$ and is isomorphic to $M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$ if $s = 1$.

The lemma now follows. $\square$

Lemma 23. *Let $a_{i+2} = a_i > a_{i+1}$. Then*

$$LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

Proof. Consider the short exact sequence

$$0 \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

Thus $L_s Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong 0$ if $s = 0, 2$ and is isomorphic to $M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}})$ if $s = 1$.

Now consider the short exact sequence

$$0 \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M(a_i, a_{i+2}, a_{i+1}) \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow 0.$$

This gives $L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \cong 0$ if $s = 0, 2$ and is isomorphic to $M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$ if $s = 1$.

The lemma now follows. $\square$

These were the two easiest cases and the computations were made in [BFK]. Now we come to the three new cases.

Lemma 24. *Suppose $a_{i+2} > a_i > a_{i+1}$. Then*

$$LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \oplus \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}).$$

Proof. Consider the short exact sequence

$$0 \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

The long exact sequence for $LZ^{\mathfrak{s}_{i+1}}$ reduces to the short exact sequence

$$0 \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

Thus if $s = 1$

$$L_s Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}) / M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_{i+2}, a_i})$$

and is 0 otherwise.

Now consider

$$0 \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M(a_{i+1}, a_{i+2}, a_i) \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow 0.$$

This gives a long exact sequence for $LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}$ reducing to

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow 0.$$

Thus if $s = 1$,

$$L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \cong M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) / M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$$

and it is zero otherwise.

Now consider

$$0 \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M(a_i, a_{i+2}, a_{i+1}) \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow 0.$$

This gives a long exact sequence for $LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}$ reducing to

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}) \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

Thus the long exact sequence for

$$0 \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0$$

becomes

$$M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) / M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \hookrightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \twoheadrightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

Thus we get the following diagram:

$$\begin{array}{ccccc}
0 & & & & 0 \\
\downarrow & & & & \downarrow h \\
M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) & & & & M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}) \\
\downarrow f & & & & \downarrow i \\
M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) & & & & \\
\downarrow g & & & & \downarrow l \\
M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) / M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) & \xrightarrow{j} & L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(\underbrace{a_i, a_{i+2}, a_{i+1}}) & \xrightarrow{k} & L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \\
& & \downarrow & & \\
& & M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) & & \\
& & \downarrow & & \\
& & 0 & &
\end{array}$$

Assume that $l \circ j \neq 0$. Then the following composition of maps is non-zero:

$$M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) / M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}).$$

A non-zero endomorphism of a generalized Verma module must be an isomorphism so the first map above must be injective as well. This is impossible so $l \circ j = 0$. Thus

$$M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) / M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \hookrightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}).$$

We claim that there is a well defined map from $L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}})$ to $M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}})$. Via k , pull an element back and then apply l . Suppose $k(x) = k(y)$. Then $k(x - y) = 0$ so $x - y = j(z)$ for some $z \in M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}})$. Then $l(x - y) = lj(z)$. This is zero by the observation above. Thus the map is well defined and call it ϕ . It is easy to see that ki is a surjection of $M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})$ onto $\ker \phi$. The kernel of this surjection is obviously $M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}})$. Thus the kernel of ϕ is

$$M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})/(M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}})).$$

Since this module is isomorphic to $\epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})$, there is a short exact sequence

$$0 \rightarrow \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

Finally we must show that this exact sequence splits.

$$\begin{aligned} \text{Hom}(M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}), \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})[1]) &\cong \text{Hom}(LZ_{\mathfrak{s}_i}^{\mathfrak{t}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}), M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})[1]) \\ &\cong \text{Hom}(M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})[2], M^{\mathfrak{t}_i}(\underbrace{a_{i+2}, a_i, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}})[1]). \end{aligned}$$

This last space is zero so the sequence must split. $\square$

Lemma 25. *Suppose $a_i > a_{i+2} > a_{i+1}$. Then*

$$LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \oplus \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}})[-1].$$

Proof. Consider the short exact sequence

$$0 \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

The long exact sequence for $LZ^{\mathfrak{s}_{i+1}}$ reduces to the short exact sequence

$$0 \rightarrow M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

We compute $LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}$ on the first and third terms of this short exact sequence.

Consider

$$0 \rightarrow M(a_{i+1}, a_{i+2}, a_i) \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

The long exact sequence gives

$$L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{a_i, a_{i+1}, a_{i+2}})$$

if $s = 1$ and is zero otherwise.

Next consider

$$0 \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M(a_i, a_{i+2}, a_{i+1}) \rightarrow M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \rightarrow 0.$$

The long exact sequence for $LZ^{\mathfrak{s}_i}$ becomes

$$0 \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \rightarrow L_0 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \rightarrow 0.$$

The middle map is clearly the standard map. Therefore

$$L_0 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \cong \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}})$$

and

$$L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{a_i, a_{i+2}, a_{i+1}}).$$

Now the long exact sequence for the functor $LZ^{\mathfrak{s}_i}$ and

$$0 \rightarrow M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}_{a_i, a_{i+2}, a_{i+1}}) \rightarrow L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow M^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}_{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

is

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i) \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow \\ M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) / M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i) \rightarrow \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{}) \rightarrow L_0 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow 0.$$

Suppose that

$$\text{Hom}(M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) / M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i), \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \neq 0.$$

Then

$$0 \neq \text{Hom}(M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \\ \cong \text{Hom}(L Z_{\mathfrak{s}_i}^{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \\ \cong \text{Hom}(M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})[1], M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \\ = 0.$$

Thus $L_0 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \cong \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})$ and there is the short exact sequence

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i) \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) / M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i) \rightarrow 0.$$

Next we show that

$$\text{Hom}(L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i)) \cong 0$$

so that the sequence does not split. The distinguished triangle

$$H^0 L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow \tau^{\geq 0} L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow \\ \tau^{\geq 1} L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2})$$

is isomorphic to

$$L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}) \rightarrow \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})[-1].$$

Now apply the functor $\text{Hom}(\bullet, M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i))$ to get a long exact sequence

$$\cdots \rightarrow \text{Hom}(L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i)) \rightarrow \\ \text{Hom}(L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i)) \rightarrow 0.$$

We calculate

$$\text{Hom}(L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i)) \cong \\ \text{Hom}(M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), L Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] L Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i)) \cong \\ \text{Hom}(M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}}_{}, a_i) \oplus \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \cong \\ \text{Hom}(M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), \epsilon_{t_i}^{\mathfrak{s}_i} M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \cong \\ \text{Hom}(L Z_{\mathfrak{s}_i}^{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}_{}, a_{i+2}), M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \cong \\ \text{Hom}(M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})[1], M^{t_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{})) \cong 0.$$

The second isomorphism is due to the previous lemma. Therefore the sequence cannot split.

It remains to be shown that the extension is unique up to isomorphism. Consider the full subcategory of objects which have composition factors with weights less than or equal to $(\underbrace{a_i, a_{i+1}, a_{i+2}})$. In this full subcategory the generalized Verma module $M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$ is projective. Thus for any exact sequence

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) \rightarrow X \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) \rightarrow 0$$

there is a nontrivial map $g: M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow X$. Note that all of the objects in the above sequence are in this subcategory. Now we can set up the following diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) & \xrightarrow{j} & M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) & \xrightarrow{f} & M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) \longrightarrow 0 \\ & & \downarrow Id & & \downarrow g & & \downarrow Id \\ 0 & \longrightarrow & M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) & \xrightarrow{i} & X & \xrightarrow{h} & M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}) \longrightarrow 0 \end{array}$$

The square on the right commutes by construction. Take an element in the top left. Applying the map hgj must give zero. Therefore $\text{im}(gj) \subset \ker(h) \cong M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i})$. But then i must be the same as gj up to a scalar since scalar multiplication are the only endomorphisms of $M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i})$. If the scalar is zero then there is a map from $M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})/M^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i})$ to X giving a splitting, contrary to the above where it was shown that the extension must be non-split. Thus the scalar must be non-zero and so the middle map is an isomorphism by the five lemma.

Therefore,

$$L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} L_1 Z^{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

□

Lemma 26. Suppose $a_i > a_{i+1} > a_{i+2}$. Then

$$LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} [-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \oplus \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-2].$$

Proof. Let $X = LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} [-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$.

Consider the short exact sequence

$$0 \rightarrow M(a_{i+1}, a_i, a_{i+2}) \rightarrow M(a_i, a_{i+1}, a_{i+2}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

This induces a long exact sequence for $LZ^{\mathfrak{s}_{i+1}}$

$$L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \hookrightarrow M^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}) \rightarrow M^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+1}, a_{i+2}}) \twoheadrightarrow L_0 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

The middle map is clearly the standard map so

$$L_1 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}})$$

and

$$L_0 Z^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i} M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \cong \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}).$$

Thus

$$\begin{aligned} H^0(X) &\cong M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}}) \\ H^1(X) &\cong \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \\ H^s(X) &\cong 0, \forall s \neq 0, 1. \end{aligned}$$

Therefore we get the distinguished triangle

$$H^0(X)[0] \rightarrow \tau^{\geq 0} X \rightarrow \tau^{\geq 1} X$$

which is

$$(1) \quad M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}}) \rightarrow X \rightarrow \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_{i+1}} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-1].$$

Next we compute $L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}})$. Consider the short exact sequence

$$0 \rightarrow M(a_{i+2}, a_{i+1}, a_i) \rightarrow M(a_{i+2}, a_i, a_{i+1}) \rightarrow M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}}) \rightarrow 0.$$

This gives a long exact sequence for $LZ^{\mathfrak{s}_i}$:

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}}) \rightarrow 0.$$

Thus if $s = 1$

$$L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}}, \dots, a_n) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}) / M^{\mathfrak{s}_i}(\underbrace{a_{i+1}, a_{i+2}})$$

and it is zero otherwise.

Next to compute $L_s Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_{i+1}} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$:

$$\begin{aligned} LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_{i+1}} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) &\cong LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_i} \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \\ &\cong \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \oplus \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[2]. \end{aligned}$$

The last isomorphism follows from a previous section.

Consider (1) again. It gives rise to a distinguished triangle

$$LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}}) \rightarrow LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X \rightarrow LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-1].$$

This gives rise to a long exact sequence

$$\begin{aligned} &\rightarrow H^s(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] M^{\mathfrak{s}_{i+1}}(a_{i+2}, \underbrace{a_i, a_{i+1}})) \rightarrow H^s(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) \rightarrow \\ &H^s(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-1]) \rightarrow \end{aligned}$$

This becomes

$$\begin{aligned} 0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}) / M^{\mathfrak{s}_i}(\underbrace{a_{i+1}, a_{i+2}, a_i}) &\rightarrow H^0(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) \rightarrow \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0 \rightarrow \\ H^1(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) &\rightarrow L_1 Z^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0 \rightarrow \\ H^2(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) &\rightarrow \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0. \end{aligned}$$

Thus

$$H^2(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) \cong \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$$

and

$$H^1 LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X \cong 0.$$

We must prove $H^0(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) \cong M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$. This reduces to showing that

$$0 \rightarrow M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}) / M^{\mathfrak{s}_i}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow H^0(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] X) \rightarrow \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i} M^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0$$

is the unique non-trivial extension up to isomorphism.

The distinguished triangle

$$\begin{aligned} H^0(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) &\rightarrow \tau^{\geq 0}(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) \rightarrow \\ \tau^{\geq 1}(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] M^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) &\rightarrow \end{aligned}$$

is isomorphic to

$$H^0(LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1] M^{s_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) \rightarrow LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1] M^{s_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow \epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-2].$$

Apply the functor $\text{Hom}(\epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), \bullet)$ to get a long exact sequence

$$0 \rightarrow \text{Hom}(\epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), H^0 LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1] M^{s_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) \rightarrow \text{Hom}(\epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1] M^{s_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) \rightarrow \dots$$

We calculate

$$\begin{aligned} & \text{Hom}(\epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1] M^{s_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) \cong \\ & \text{Hom}(M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), LZ_{s_i}^{t_i}[-4] LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1] M^{s_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})) \cong \\ & \text{Hom}(M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-6] \oplus (M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-4])^{\oplus 2} \oplus M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-2]) \\ & = 0. \end{aligned}$$

The second to last isomorphism comes from using proposition 8 twice. This shows that the extension is non-trivial. Now we show that it is unique.

Consider the short exact sequence

$$0 \rightarrow M^{s_i}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow M^{s_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}) \rightarrow M^{s_i}(\underbrace{a_i, a_{i+2}, a_{i+1}})/M^{s_i}(\underbrace{a_{i+1}, a_{i+2}, a_i}) \rightarrow 0.$$

Applying $LZ_{s_i}^{t_i}$ to get a long exact sequence we easily see that

$$L_s Z_{s_i}^{t_i}(M^{s_i}(\underbrace{a_i, a_{i+2}, a_{i+1}})/M^{s_i}(\underbrace{a_{i+1}, a_{i+2}, a_i})) \cong M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$$

if $s = 1, 3$ and 0 otherwise.

Now,

$$\begin{aligned} & \text{Ext}^1(\epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), M^{s_i}(\underbrace{a_i, a_{i+2}, a_{i+1}})/M^{s_i}(\underbrace{a_{i+1}, a_{i+2}, a_i})) \cong \\ & \text{Hom}(\epsilon_{t_i}^{s_i} M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}), M^{s_i}(\underbrace{a_i, a_{i+2}, a_{i+1}})/M^{s_i}(\underbrace{a_{i+1}, a_{i+2}, a_i})[1]) \cong \\ & \text{Hom}(M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[0], M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[0] \oplus M^{t_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})[-2]) \cong \mathbb{C}. \end{aligned}$$

□

We would like to construct a natural transformation from $LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1]$ to $\text{Id} \oplus \epsilon_{t_i}^{s_i}[-2] LZ_{s_i}^{t_i}$.

The identity map gives the adjunction map α under the isomorphism $\text{Hom}(LZ^{s_{i+1}} \epsilon_{s_i}, LZ^{s_{i+1}} \epsilon_{s_i}) \cong \text{Hom}(LZ^{s_i} \epsilon_{s_{i+1}}[-1] LZ^{s_{i+1}} \epsilon_{s_i}[-1], \text{Id})$.

The identity map in $\text{Hom}(\epsilon_{s_i}, \epsilon_{s_i})$ gives an adjunction map in $\text{Hom}(LZ^{s_i} \epsilon_{s_i}, \text{Id})$. Apply $LZ_{s_i}^{t_i}$ to get a map in

$$\begin{aligned} \text{Hom}(LZ_{s_i}^{t_i} LZ^{s_i} \epsilon_{s_i}, LZ_{s_i}^{t_i}) & \cong \text{Hom}(LZ_{s_i}^{t_i} \epsilon_{s_i}, LZ_{s_i}^{t_i}) \\ & \cong \text{Hom}(LZ_{s_{i+1}}^{t_i} LZ^{s_{i+1}} \epsilon_{s_i}, LZ_{s_i}^{t_i}) \\ & \cong \text{Hom}(LZ^{s_{i+1}} \epsilon_{s_i}, \epsilon_{t_i}^{s_{i+1}} LZ_{s_i}^{t_i}). \end{aligned}$$

Now apply the functor $\epsilon_{s_{i+1}}$ to get an element of

$$\begin{aligned} \text{Hom}(\epsilon_{s_{i+1}} LZ^{s_{i+1}} \epsilon_{s_i}, \epsilon_{s_{i+1}} \epsilon_{t_i}^{s_{i+1}} LZ_{s_i}^{t_i}) & \cong \text{Hom}(\epsilon_{s_{i+1}} LZ^{s_{i+1}} \epsilon_{s_i}, \epsilon_{t_i} LZ_{s_i}^{t_i}) \\ & \cong \text{Hom}(\epsilon_{s_{i+1}} LZ^{s_{i+1}} \epsilon_{s_i}, \epsilon_{s_i} \epsilon_{t_i}^{s_i} LZ_{s_i}^{t_i}) \\ & \cong \text{Hom}(LZ^{s_i} \epsilon_{s_{i+1}} LZ^{s_{i+1}} \epsilon_{s_i}, \epsilon_{t_i}^{s_i} LZ_{s_i}^{t_i}). \end{aligned}$$

Now apply the shift functor $[-2]$ to get a morphism β in

$$\mathrm{Hom}(LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1], \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i}[-2] LZ_{\mathfrak{s}_i}^{\mathfrak{t}_i}).$$

Therefore $\alpha \oplus \beta: LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] \rightarrow \mathrm{Id} \oplus \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i}[-2] LZ_{\mathfrak{s}_i}^{\mathfrak{t}_i}$.

Now we come to the main result of the section.

Proposition 12. *The map*

$$\alpha \oplus \beta: LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] \rightarrow \mathrm{Id} \oplus \epsilon_{\mathfrak{t}_i}^{\mathfrak{s}_i}[-2] LZ_{\mathfrak{s}_i}^{\mathfrak{t}_i}$$

is an isomorphism.

Proof. Lemmas 23-27 show that the cohomology functors of these morphisms are isomorphisms when applied to generalized Verma modules. Thus the morphisms are isomorphisms when applied to projective modules by induction on the length of the generalized Verma flag. Finally, they are isomorphisms on arbitrary modules by considering projective resolutions. $\square$

The above theorem is the Koszul dual to lemma 4 of [BFK]. The correction term there vanishes when restricted to the appropriate parabolic subcategory just as in the case here where the correction terms vanishes when restricted to the appropriate singular block. This result is not a surprise in light of [Rh] or [MOS] where it was shown that projective functors and Zuckerman functors are Koszul dual. We end this section with an immediate corollary which is the functorial version of diagram 5.

Corollary 10. $\epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i} \oplus \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \cong$
 $\epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i} \epsilon_{\mathfrak{s}_{i+1}}[-1] LZ^{\mathfrak{s}_{i+1}} \oplus \epsilon_{\mathfrak{s}_i}[-1] LZ^{\mathfrak{s}_i}.$

5. GRADED CATEGORY $\mathcal{O}$

In the previous sections, various aspects of the representation theory of $\mathfrak{sl}_k$ were categorified. Topological invariants arise from the representation theory of $\mathcal{U}_q(\mathfrak{sl}_k)$. The categorification of quantum groups is accomplished through graded representation theory. We will treat category $\mathcal{O}$ as a category of graded modules. Then a shift in this grading descends to multiplication by q in the Grothendieck group. The idea of graded category $\mathcal{O}$ originates from [Soe2]. It was continued in [Str1] where among other things, it was shown how to construct graded lifts of translation functors.

Definition 14. Let $P_{\mathbf{d}} = \bigoplus_{x \in S_n/S_{\mathbf{d}}} P(x, \omega_{\mathbf{d}})$ where $\mathbf{d} = (d_{k-1}, \dots, d_0)$, x is a minimal coset representative, and $\omega_{\mathbf{d}} = (\underbrace{k-1, \dots, k-1}_{d_{k-1}}, \dots, \underbrace{0, \dots, 0}_{d_0})$.

Then $P_{\mathbf{d}}$ is a minimal projective generator of $\mathcal{O}_{\mathbf{d}}(\mathfrak{gl}_n)$. There is an equivalence of categories $\mathcal{O}_{\mathbf{d}}(\mathfrak{gl}_n) \cong \mathrm{mod} - A_{\mathbf{d}}$ where $A_{\mathbf{d}} = \mathrm{End}_{\mathcal{O}}(P_{\mathbf{d}})$ and $\mathrm{mod} - A_{\mathbf{d}}$ is the category of finitely generated right $A_{\mathbf{d}}$ -modules. We will interpret $A_{\mathbf{d}}$ as a graded algebra.

Lemma 27. *Let R and S be any rings. There is an equivalence of categories:*

$$\{\text{right exact functors compatible with direct sums} : (\mathrm{mod} - R \rightarrow \mathrm{mod} - S)\} \rightarrow R\text{-mod} - S.$$

Under this equivalence a functor F gets mapped to $F(R)$. In the other direction, a bimodule X gets mapped to $\bullet \otimes_R X$.

Proof. This is in [Bass] as well as lemma 3.4 of [Str1]. $\square$

Let α and β be parabolic subalgebras of $\mathfrak{gl}_n$ such that $\beta \subset \alpha$. Let $P(x) \in \mathcal{O}_{\mathbf{d}}(\mathfrak{gl}_n)$ be an indecomposable projective object. Then $Z^{\alpha} P(x)$ is either 0 or an indecomposable projective object in $\mathcal{O}_{\mathbf{d}}^{\alpha}(\mathfrak{gl}_n)$.

Denote by $P_{\mathbf{d}}^{\alpha}$ a minimal projective generator of $\mathcal{O}_{\mathbf{d}}^{\alpha}(\mathfrak{gl}_n)$. Let $A_{\mathbf{d}}^{\alpha} = \mathrm{End}(P_{\mathbf{d}}^{\alpha})$ be its endomorphism algebra. Then there is an equivalence of categories $\mathcal{O}_{\mathbf{d}}^{\alpha}(\mathfrak{gl}_n) \cong \mathrm{mod} - A_{\mathbf{d}}^{\alpha}$. Let $S = S(\mathfrak{h})$ be the symmetric algebra associated to the Cartan subalgebra. Let $C = S/S_{+}^W$ be the associated coinvariant algebra. Let C^{λ} be the subalgebra invariant under W_{λ} . Let w_0^{λ} be the longest element in the set of shortest coset representatives of W/W_{λ} . There is a well known isomorphism due to Soergel [Soe] between the endomorphism algebra of the indecomposable projective-injective module and this subalgebra of invariants: $\mathrm{End}(P(w_0^{\lambda}, \lambda)) \cong C^{\lambda}$.

Definition 15. Let $\mathbb{V}_\lambda: \mathcal{O}_\lambda \rightarrow \text{mod} - C^\lambda$ be the Soergel functor defined by $M \mapsto \text{Hom}_{\mathfrak{g}}(P(w_0^\lambda \cdot \lambda), M)$.

Soergel showed that the functor $\mathbb{V}_\lambda$ is fully faithful on projective objects [Soe]. Let I and J be compositions of n . If $I = i_1 + \dots + i_r = n$, associate to I the Young subgroup of S_n , $S_{i_1} \times \dots \times S_{i_r}$. Let μ_I and μ_J be integral dominant weights stabilized by the subgroups associated to I and J . Suppose $J \subset I$, (there is a containment of the associated subgroups.) Then we can define the translation functor θ_I^J from $\mathcal{O}_{\mu_I}(\mathfrak{gl}_n)$ to $\mathcal{O}_{\mu_J}(\mathfrak{gl}_n)$. It is the projective functor given by tensoring with the finite dimensional, irreducible module with highest weight $\mu_J - \mu_I$ and then projecting onto the block $\mathcal{O}_{\mu_J}(\mathfrak{gl}_n)$. Translation functors and Soergel functors are related by the following lemma.

Lemma 28. Let $\text{Res}_J^I: \text{mod} - C^J \rightarrow \text{mod} - C^I$ denote the restriction functor. Then

- (1) $\mathbb{V}_J \theta_I^J \cong C^J \otimes_{C^I} \mathbb{V}_I$.
- (2) $\mathbb{V}_I \theta_J^I \cong \text{Res}_J^I \mathbb{V}_J$.

Proof. This is proposition 3.3 of [FKS]. □

This lemma together with the fact that $\mathbb{V}_\lambda$ is a faithful functor on projective objects allows us to consider the endomorphism ring of a minimal projective generator of $\mathcal{O}$ as a graded ring [Str1]. A projective object is a direct summand of a sequence of projective functors applied to a dominant Verma module. Thus by the previous lemma, a projective object P , $\mathbb{V}_\lambda P$ becomes a graded C^λ -module. Then $\text{End}(\mathbb{V}_\lambda P)$ becomes a graded ring so there is a grading on $A_{\mathbf{d}}$.

In [Str] it was actually shown that it is a graded quotient of $A_{\mathbf{d}}$. Now we can consider $\mathcal{O}_{\mathbf{d}}(\mathfrak{gl}_n)$ and $\mathcal{O}_{\mathbf{d}}^\alpha(\mathfrak{gl}_n)$ as graded categories by considering $\text{gmod} - A_{\mathbf{d}}$ and $\text{gmod} - A_{\mathbf{d}}^\alpha$ respectively.

By lemma 27, the dual Zuckerman functor becomes

$$Z_\beta^\alpha: \text{mod} - A_{\mathbf{d}}^\beta \rightarrow \text{mod} - A_{\mathbf{d}}^\alpha$$

given by

$$\bullet \otimes_{A_{\mathbf{d}}^\beta} A_{\mathbf{d}}^\alpha.$$

Definition 16. The graded functor $\widetilde{Z}_\beta^\alpha: \text{gmod} - A_{\mathbf{d}}^\beta \rightarrow \text{gmod} - A_{\mathbf{d}}^\alpha$ is given by $\bullet \otimes_{A_{\mathbf{d}}^\beta} A_{\mathbf{d}}^\alpha \langle -(\dim(\alpha) - \dim(\beta))/2 \rangle$.

We may now also consider

$$L\widetilde{Z}_\beta^\alpha: D^b(\text{gmod} - A_{\mathbf{d}}^\beta) \rightarrow D^b(\text{gmod} - A_{\mathbf{d}}^\alpha).$$

We fix a graded lift of a generalized Verma module $\widetilde{M}^\mathbb{P}(x \cdot \lambda)$ so that its head is concentrated in degree zero. The proof of proposition 5.2 of [FKS] gives a graded inclusion $\widetilde{M}(x \cdot \lambda) \langle l(x) \rangle \rightarrow \widetilde{M}(\lambda)$ of Verma modules. This implies that the standard maps in a generalized BGG resolution are homogeneous of degree 1. A Koszul module is an object that admits a projective resolution such that the object in degree i is generated by its degree i subspace. We will need the following result later.

Proposition 13. Generalized Verma modules are Koszul.

Proof. This is theorem 3.11.4 of [BGS]. □

Proposition 14. Let $\beta \subset \alpha$ be two parabolic subalgebras. Assume $\dim(\alpha) - \dim(\beta) = 2d$. Then

- (1) The right adjoint of ϵ_α^β is $L\widetilde{Z}_\beta^\alpha[-2d] \langle -d \rangle$.
- (2) The right adjoint of $L\widetilde{Z}_\beta^\alpha$ is $\widetilde{\epsilon}_\alpha^\beta \langle d \rangle$.

Proof. (1) In the ungraded case the right adjoint of ϵ_α^β is $LZ_\beta^\alpha[-2d]$. Therefore if the right adjoint of $\widetilde{\epsilon}_\beta^\alpha$ exists, it must be a graded lift of $LZ_\beta^\alpha[-2d]$. Note that $\widetilde{\epsilon}_\alpha^\beta M$ becomes a right A^β -module via the quotient map $A^\beta \rightarrow A^\alpha$. Therefore

$$\text{Hom}_{\text{gmod} - A^\beta}(\widetilde{\epsilon}_\alpha^\beta M, N) \cong \text{Hom}_{\text{gmod} - A^\alpha}(M, \text{Hom}_{\text{gmod} - A^\beta}(A^\alpha, N)).$$

Now $\text{Hom}_{\text{gmod}-A^\beta}(A^\alpha, \bullet)$ is clearly a functor from $\text{gmod}-A^\beta$ to $\text{gmod}-A^\alpha$. The right adjoint is a direct sum of graded functors $\oplus_i F_i$ which in the ungraded category is simply $L_{2d}Z_\beta^\alpha$. Let $\nabla(x)$ denote a dual Verma module. We have an isomorphism:

$$\text{Hom}_{\text{gmod}-A^\beta}(\tilde{\epsilon}_\alpha^\beta \tilde{\nabla}^\alpha(x), \tilde{\nabla}^\beta(x)) \cong \text{Hom}_{\text{gmod}-A^\alpha}(\tilde{\nabla}^\alpha(x), \oplus_i F_i \tilde{\nabla}^\beta(x)).$$

If $F_i \tilde{\nabla}^\beta(x) \neq 0$, then $F_i \tilde{\nabla}^\beta(x) \cong L_{2d} \tilde{Z}_\beta^\alpha \tilde{\nabla}^\beta(x) \langle k_i \rangle$. By the definition of $\tilde{Z}_\beta^\alpha$ and the fact that generalized Verma modules are Koszul, we get that the above is isomorphic to $\tilde{\nabla}^\alpha(x) \langle d + k_i \rangle$. Since the space of morphisms is one dimensional, $k_i = -d$.

- (2) There is a degree zero homogenous map of A^β -modules $\psi: M \rightarrow M \otimes_{A^\beta} A^\alpha$ where $\psi(m) = m \otimes 1$. Now suppose $\phi: M \otimes_{A^\beta} A^\alpha \rightarrow N$ is a degree zero homogeneous A^β -module morphism. Then $\phi \circ \psi: M \rightarrow N$ is a degree zero homogeneous A^β -module morphism. Therefore we have a map $f: \text{Hom}_{\text{gmod}-A^\alpha}(M \otimes_{A^\beta} A^\alpha, N) \rightarrow \text{Hom}_{\text{gmod}-A^\beta}(M, N)$. It is clear that f is injective. We must verify that it is an isomorphism. Consider a two step free resolution of M , $M_2 \rightarrow M_1 \rightarrow M \rightarrow 0$. Then we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \text{Hom}_{\text{gmod}-A^\alpha}(M_2 \otimes_{A^\beta} A^\alpha, N) & \longleftarrow & \text{Hom}_{\text{gmod}-A^\alpha}(M_1 \otimes_{A^\beta} A^\alpha, N) & \longleftarrow & \text{Hom}_{\text{gmod}-A^\alpha}(M \otimes_{A^\beta} A^\alpha, N) & \longleftarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ \text{Hom}_{\text{gmod}-A^\beta}(M_2, N) & \longleftarrow & \text{Hom}_{\text{gmod}-A^\beta}(M_1, N) & \longleftarrow & \text{Hom}_{\text{gmod}-A^\beta}(M, N) & \longleftarrow & 0 \end{array}$$

Therefore it suffices to show that f is surjective for free modules M . It is clear for free modules that both spaces of morphisms have the same dimension so it must be a surjection. $\square$

5.1. Graded Projective Functors and Quantum Serre Relations. Define the k -tuple $\mathbf{d} + t_0\epsilon_0 + \dots + t_{k-1}\epsilon_{k-1}$ to be $(d_{k-1} + t_{k-1}, \dots, d_0 + t_0)$ where the t_i are integers. Assume $t > 0$. Define $(\mathbf{d}; \mathbf{d} + t\epsilon_i - t\epsilon_{i-1})$ to be the $(k+1)$ -tuple $(d_{k-1}, \dots, d_i, t, d_{i-1} - t, d_{i-2}, \dots, d_0)$. Define $(\mathbf{d}; \mathbf{d} - t\epsilon_i + t\epsilon_{i-1})$ to be the $(k+1)$ -tuple $(d_{k-1}, \dots, d_i - t, t, d_{i-1}, d_{i-2}, \dots, d_0)$.

Lemma 29. $\mathcal{E}_i^{(t)}: \mathcal{O}_{\mathbf{d}} \rightarrow \mathcal{O}_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}$ is isomorphic to $\theta_{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}^{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} \theta_{\mathbf{d}}^{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}$.

Proof. See proposition 3.2b of [FKS]. $\square$

Lemma 30. $\mathcal{F}_i^{(t)}: \mathcal{O}_{\mathbf{d}} \rightarrow \mathcal{O}_{\mathbf{d}-t\epsilon_i+t\epsilon_{i-1}}$ is isomorphic to $\theta_{\mathbf{d}; \mathbf{d}-t\epsilon_i+t\epsilon_{i-1}}^{\mathbf{d}-t\epsilon_i+t\epsilon_{i-1}} \theta_{\mathbf{d}}^{\mathbf{d}; \mathbf{d}-t\epsilon_i+t\epsilon_{i-1}}$.

Proof. Same as the previous lemma. $\square$

Now we are prepared to introduce the graded lifts $\tilde{\mathcal{E}}_i^{(t)}$, $\tilde{\mathcal{F}}_i^{(t)}$, $\tilde{\mathcal{H}}$, and $\tilde{\mathcal{H}}^{-1}$. By lemma 27, $\mathcal{E}_i^{(t)}: \text{mod}-A_{\mathbf{d}} \rightarrow \text{mod}-A_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}$ is given by

$$\bullet \otimes_{\text{mod}-A_{\mathbf{d}}} \text{Hom}_g(P_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}, \mathcal{E}_i^{(t)} P_{\mathbf{d}}).$$

By lemma 29 this is

$$\bullet \otimes_{\text{mod}-A_{\mathbf{d}}} \text{Hom}_g(P_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}, \theta_{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}^{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} \theta_{\mathbf{d}}^{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} P_{\mathbf{d}}).$$

Then by lemma 28 this is isomorphic to

$$\bullet \otimes_{\text{gmod}-A_{\mathbf{d}}} \text{Hom}_{C^{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} P_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}, \text{Res}_{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}^{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} C^{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}}).$$

Definition 17. (1) Let $r_{i,t} = \sum_{r=1}^t d_{i-1} - r$. Then we let $\tilde{\mathcal{E}}_i^{(t)} =$

$$\bullet \otimes_{\text{gmod}-A_{\mathbf{d}}} \text{Hom}_{C^{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} P_{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}, \text{Res}_{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}}^{\mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} C^{\mathbf{d}; \mathbf{d}+t\epsilon_i-t\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}} \langle -r_{i,t} + (t-1) \rangle).$$

(2) Let $s_{i,t} = \sum_{s=1}^t d_i - s$. Then we let $\tilde{\mathcal{F}}_i^{(t)} =$

$$\bullet \otimes_{\text{gmod}-A_{\mathbf{d}}} \text{Hom}_{C^{\mathbf{d}-t\epsilon_i+t\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}-t\epsilon_i+t\epsilon_{i-1}} P_{\mathbf{d}-t\epsilon_i+t\epsilon_{i-1}}, \text{Res}_{\mathbf{d}; \mathbf{d}-t\epsilon_i+t\epsilon_{i-1}}^{\mathbf{d}-t\epsilon_i+t\epsilon_{i-1}} C^{\mathbf{d}; \mathbf{d}-t\epsilon_i+t\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}} \langle -s_{i,t} + (t-1) \rangle).$$

(3) $\tilde{\mathcal{H}} = \text{Id} \langle d_i - d_{i-1} \rangle$.

(4) $\tilde{\mathcal{H}}^{-1} = \text{Id} \langle -(d_i - d_{i-1}) \rangle$.

Lemma 31. $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \langle 1 \rangle \oplus \tilde{\mathcal{E}}_i^{(2)} \langle -1 \rangle$.

Proof. First we consider the ungraded case and compute in the Grothendieck group.

$$\begin{aligned} & [\mathcal{E}_i][\mathcal{E}_i][M(k-1, \dots, k-1, \dots, i+1, \dots, i+1, i, \dots, i, i-1, \dots, i-1, \dots, 0, \dots, 0)] = \\ & 2[P(k-1, \dots, k-1, \dots, i, \dots, i, \underbrace{i-1, \dots, i-1}_{d_{i-1}}, i, i, \dots, 0, \dots, 0)]. \end{aligned}$$

The indecomposable projective functor that takes the dominant Verma module to this projective object is $\mathcal{E}_i^{(2)}$. Thus there is an isomorphism of functors $\mathcal{E}_i \mathcal{E}_i \cong \mathcal{E}_i^{(2)} \oplus \mathcal{E}_i^{(2)}$. So there must be an isomorphism of graded functors $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \langle r \rangle \oplus \tilde{\mathcal{E}}_i^{(2)} \langle s \rangle$, for some integers r and s .

$\tilde{\mathcal{E}}_i^{(2)}$ is given by tensoring with the graded bimodule

$$\text{Hom}_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} P_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}} \langle -2d_{i-1} + 4 \rangle).$$

As a $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ -module, $C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ is free of rank $|W^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}/W^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}|$. A basis can be chosen homogeneous in the degrees twice the length of minimal coset representatives of

$$\begin{aligned} & |W^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}/W^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}| = \\ & S_{d_{k-1}} \times \dots \times S_{d_{i+1}} \times S_{d_i+2} \times S_{d_{i-1}-2} \times \dots \times S_{d_0}/S_{d_{k-1}} \times \dots \times S_{d_i} \times S_2 \times S_{d_{i-1}-2} \times \dots \times S_{d_0}. \end{aligned}$$

So now we want to consider minimal coset representatives in $S_{d_i+2}/(S_{d_i} \times S_2)$. Proposition A-2 in [Str] tells us the the length of these coset representatives. It gives

$$C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \cong \oplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s - 2 \rangle.$$

Now $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i$ is given by tensoring with the bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}} P_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}} \langle -d_{i-1} + 1 \rangle) \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} P_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \mathbb{V}_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}} P_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}} \langle -d_{i-1} + 2 \rangle). \end{aligned}$$

Thus we need to compute $C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}}$ as a $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ -module. As above, $C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}}$ is a free $C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}$ -module of rank $|W^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}/W^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}}|$ and a basis can be chosen homogeneous in degrees twice the length of minimal coset representatives. There is a similar description for $C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ as a $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ -module. Thus

$$C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \cong \oplus_{s=0}^{d_i+1} \oplus_{r=0}^{d_i} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s \rangle.$$

Therefore the grading of $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i$ is encoded by

$$C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \cong \oplus_{s=0}^{d_i+1} \oplus_{r=0}^{d_i} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s - 2d_{i-3} + 3 \rangle.$$

The grading of $\tilde{\mathcal{E}}_i^{(2)}$ is encoded by

$$C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \cong \oplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s - 2 - 2d_{i-1} + 4 \rangle.$$

By examining the highest and lowest degrees, $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \langle 1 \rangle \oplus \tilde{\mathcal{E}}_i^{(2)} \langle -1 \rangle$. □

Lemma 32. $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i^{(2)} \langle 1 \rangle \oplus \tilde{\mathcal{F}}_i^{(2)} \langle -1 \rangle$.

Proof. First we consider the ungraded case and compute in the Grothendieck group.

$$\begin{aligned} & [\mathcal{F}_i][\mathcal{F}_i][M(k-1, \dots, k-1, \dots, i+1, \dots, i+1, i, \dots, i, i-1, \dots, i-1, \dots, 0, \dots, 0)] = \\ & 2[P(k-1, \dots, k-1, \dots, i+1, \dots, i+1, \underbrace{i-1, i-1, i, \dots, i}_{d_i}, \dots, 0, \dots, 0)]. \end{aligned}$$

The indecomposable projective functor that takes the dominant Verma module to this projective object is $\mathcal{F}_i^{(2)}$. Thus there is an isomorphism of functors $\mathcal{F}_i \mathcal{F}_i \cong \mathcal{F}_i^{(2)} \oplus \mathcal{F}_i^{(2)}$. So there must be an isomorphism of graded functors $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i^{(2)} \langle r \rangle \oplus \tilde{\mathcal{F}}_i^{(2)} \langle s \rangle$, for some integers r and s .

$\tilde{\mathcal{F}}_i^{(2)}$ is given by tensoring with the graded bimodule

$$\text{Hom}_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} P_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}, \text{Res}_{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}} \langle -2d_i + 4 \rangle).$$

As a $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}-}$ module, $C^{\mathbf{d};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}$ is free of rank $|W^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}/W^{\mathbf{d};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}|$. A basis can be chosen homogenous in the degrees twice the length of minimal coset representatives of

$$|W^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}/W^{\mathbf{d};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}| = S_{d_{k-1}} \times \cdots \times S_{d_{i+1}} \times S_{d_i-2} \times S_{d_{i-1}+2} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_i-2} \times S_2 \times S_{d_{i-1}} \times \cdots \times S_{d_0}.$$

So now we want to consider minimal coset representatives in $S_{d_{i-1}+2}/(S_2 \times S_{d_{i-1}})$. Proposition A-2 in [Str] tells us the the length of these coset representatives. It gives

$$C^{\mathbf{d};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \cong \oplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle 2r+2s-2 \rangle.$$

Now $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_i$ is given by tensoring with the bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}} P_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}, C^{\mathbf{d};\mathbf{d}-\epsilon_i+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}_{\mathbf{d}} P_{\mathbf{d}} \langle -d_i+1 \rangle) \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}}(\mathbb{V}_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} P_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}, C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} \mathbb{V}_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}} \langle -d_i+2 \rangle). \end{aligned}$$

Thus we need to compute $C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} C^{\mathbf{d};\mathbf{d}-\epsilon_i+\epsilon_{i-1}}$ as a $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}-}$ module. As above, $C^{\mathbf{d};\mathbf{d}-\epsilon_i+\epsilon_{i-1}}$ is a free $C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}-}$ module of rank $|W^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}/W^{\mathbf{d};\mathbf{d}-\epsilon_i+\epsilon_{i-1}}|$ and a basis can be chosen homogeneous in degrees twice the length of minimal coset representatives. There is a similar description for $C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}$ as a $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}-}$ module.

Thus

$$C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} C^{\mathbf{d};\mathbf{d}-\epsilon_i+\epsilon_{i-1}} \cong \oplus_{s=0}^{d_{i-1}} \oplus_{r=0}^{d_{i-1}+1} C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle 2r+2s \rangle.$$

Therefore the grading of $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_i$ is encoded by

$$C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} C^{\mathbf{d};\mathbf{d}-\epsilon_i+\epsilon_{i-1}} \cong \oplus_{s=0}^{d_{i-1}} \oplus_{r=0}^{d_{i-1}+1} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle 2r+2s-2d_i+3 \rangle.$$

The grading of $\tilde{\mathcal{F}}_i^{(2)}$ is encoded by

$$C^{\mathbf{d};\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \cong \oplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle 2r+2s-2d_i+2 \rangle.$$

Thus $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i^{(2)} \langle 1 \rangle \oplus \tilde{\mathcal{F}}_i^{(2)} \langle -1 \rangle$. □

Lemma 33. *For some indecomposable projective functor G_1 , there are isomorphisms of graded functors:*

- (1) $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1} \oplus \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1} \oplus \tilde{G}_1.$
- (2) $\tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)} \cong \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1} \oplus \tilde{G}_1.$
- (3) $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1} \oplus \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)}.$

Proof. First we consider the ungraded case and compute in the Grothendieck group.

$$\begin{aligned} & [\mathcal{E}_i][\mathcal{E}_{i+1}][\mathcal{E}_i][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & 2[P(k-1, \dots, k-1, \dots, i+1, \dots, i+1, \underbrace{i, \dots, i, i+1}_{d_i}, \underbrace{i-1, \dots, i-1, i, i}_{d_{i-1}}, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, i, \dots, i, \underbrace{i-1, \dots, i-1, i, i+1}_{d_{i-1}}, \dots, 0, \dots, 0)]. \end{aligned}$$

We also have

$$\begin{aligned} & [\mathcal{E}_i^{(2)}][\mathcal{E}_{i+1}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i, \dots, i, i+1}_{d_i}, \underbrace{i-1, \dots, i-1, i, i}_{d_{i-1}}, \dots, 0, \dots, 0)] \end{aligned}$$

and

$$\begin{aligned} & [\mathcal{E}_{i+1}][\mathcal{E}_i^{(2)}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, i, \dots, i, i-1, \dots, i-1, i, i+1, \dots, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, i+1, \dots, i+1, i, \dots, i, i+1, i-1, \dots, i-1, i, i, \dots, 0, \dots, 0)]. \end{aligned}$$

Therefore we get the desired isomorphism of functors if we ignore the grading. There is an isomorphism

$$\tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)} \cong \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1} \langle a \rangle \oplus \tilde{G}_1$$

for some graded lift of the indecomposable projective functor G_1 and for some shift a .

The functor $\tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d};\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -2d_{i-1} + 4 \rangle) \otimes_{A_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1};\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle -d_i - 1 \rangle). \end{aligned}$$

Therefore we must consider

$$X = C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1};\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}} C^{\mathbf{d};\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle -2d_{i-1} - d_i + 3 \rangle$$

as a $C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}$ -module.

As a $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ -module, $C^{\mathbf{d};\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ is free of rank $|W^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}/W^{\mathbf{d};\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}|$ and a basis can be chosen in degrees twice the length of minimal coset representatives. Therefore we would like to consider minimal coset representatives in $S_{d_i+2}/S_{d_i} \times S_2$. Proposition A-2 of [Str] tells us the length of these coset representatives. We get $C^{\mathbf{d};\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \cong \bigoplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s - 2 \rangle$.

Similarly, as a $C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}$ -module, $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1};\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_k} \times \cdots \times S_{d_{i+1}+1} \times S_{d_i+1} \times S_{d_{i-1}-2} \times \cdots \times S_{d_0}/S_{d_k} \times \cdots \times S_{d_{i+1}} \times S_1 \times S_{d_i+1} \times S_{d_{i-1}-2} \times \cdots \times S_{d_0}.$$

Therefore,

$$X \cong \bigoplus_{t=0}^{d_i+1} \bigoplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s + 2t - 2d_{i-1} - d_i + 1 \rangle.$$

Next, $\tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}-\epsilon_i}, C^{\mathbf{d};\mathbf{d}+\epsilon_{i+1}-\epsilon_i} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -d_i + 1 \rangle) \otimes_{A_{\mathbf{d}+\epsilon_{i+1}-\epsilon_i}} \\ & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i;\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i}} \mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}-\epsilon_i} \langle -2d_{i-1} + 4 \rangle). \end{aligned}$$

Thus we must consider

$$Y = C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i;\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i}} C^{\mathbf{d};\mathbf{d}+\epsilon_{i+1}-\epsilon_i} \langle -2d_{i-1} - d_i + 5 \rangle$$

as a $C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}$ -module. As a $C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i}$ -module, $C^{\mathbf{d};\mathbf{d}+\epsilon_{i+1}-\epsilon_i}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives from

$$S_{d_{k-1}} \times \cdots \times S_{d_{i+1}+1} \times S_{d_i-1} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i+1}} \times S_1 \times S_{d_i-1} \times \cdots \times S_{d_0}.$$

So now we want to consider minimal coset representatives in $S_{d_{i+1}+1}/S_{d_{i+1}} \times S_1$. Proposition A-2 of [Str] gives

$$C^{\mathbf{d};\mathbf{d}+\epsilon_{i+1}-\epsilon_i} \cong \bigoplus_{t=0}^{d_i+1} C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i} \langle 2t \rangle.$$

Similarly, as a $C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}$ -module, $C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_i;\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \cdots \times S_{d_{i+1}+1} \times S_{d_i+1} \times S_{d_{i-1}-2} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i+1}+1} \times S_{d_i-1} \times S_2 \times S_{d_{i-1}-2} \times \cdots \times S_{d_0}.$$

Therefore,

$$Y \cong \bigoplus_{t=0}^{d_i+1} \bigoplus_{d_i \geq r > s \geq 0} C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \langle 2r + 2s + 2t - 2d_{i-1} - d_i + 3 \rangle.$$

Finally, $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}, C^{\mathbf{d};\mathbf{d}+\epsilon_i-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -d_{i-1} + 1 \rangle) \otimes_{A_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}}, C^{\mathbf{d};\mathbf{d}+\epsilon_i-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}} \langle -d_i \rangle) \otimes_{A_{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1};\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}} \langle -d_{i-1} + 2 \rangle). \end{aligned}$$

Thus we must consider

$$\begin{aligned} W = & C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1};\mathbf{d}+\epsilon_{i+1}-\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}}} C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1};\mathbf{d}+\epsilon_{i+1}-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \\ & C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1};\mathbf{d}+\epsilon_i-\epsilon_{i-1}} \langle -2d_{i-1} - d_i + 3 \rangle. \end{aligned}$$

As in all the other computations,

$$W \cong \bigoplus_{t=0}^{d_{i+1}} \bigoplus_{r=0}^{d_i} \bigoplus_{s=0}^{d_i} C^{\mathbf{d}+\epsilon_{i+1}+\epsilon_i-2\epsilon_{i-1}} \langle 2r+2s+2t-2d_{i-1}-d_i+3 \rangle.$$

The grading for $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i$ is given by the summation

$$\begin{aligned} & \sum_{t=0}^{d_{i+1}} \sum_{r=0}^{d_i} \sum_{s=0}^{d_i} q^{2r+2s+2t-2d_{i-1}-d_i+3} = \\ & 2 \sum_{t=0}^{d_{i+1}} \sum_{d_i \geq r > s \geq 0} q^{2r+2s+2t-2d_{i-1}-d_i+3} + \sum_{t=0}^{d_{i+1}} \sum_{r=0}^{d_i} q^{4r+2t-2d_{i-1}-d_i+3}. \end{aligned}$$

The grading for $\tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)}$ is given by

$$\sum_{t=0}^{d_{i+1}} \sum_{d_i+1 \geq r > s \geq 0} q^{2(r-1)+2s+2t-2d_{i-1}-d_i+3}.$$

Letting $u = r - 1$, we get

$$\begin{aligned} & \sum_{t=0}^{d_{i+1}} \sum_{d_i \geq u \geq s \geq 0} q^{2u+2s+2t-2d_{i-1}-d_i+3} = \\ & \sum_{t=0}^{d_{i+1}} \sum_{d_i \geq u > s \geq 0} q^{2u+2s+2t-2d_{i-1}-d_i+3} + \sum_{t=0}^{d_{i+1}} \sum_{u=0}^{d_i} q^{4u+2t-2d_{i-1}-d_i+3}. \end{aligned}$$

The grading for $\tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1}$ is given by

$$\sum_{t=0}^{d_{i+1}} \sum_{d_i \geq u > s \geq 0} q^{2u+2s+2t-2d_{i-1}-d_i+3}.$$

Therefore we get the desired isomorphisms of graded functors. $\square$

Lemma 34. *For some indecomposable projective functor G_2 , there are isomorphisms of graded functors:*

- (1) $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i^{(2)} \oplus \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i^{(2)} \oplus \tilde{G}_2.$
- (2) $\tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i-1} \cong \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i^{(2)} \oplus \tilde{G}_2.$
- (3) $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i-1} \oplus \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i^{(2)}.$

Proof. First we consider the ungraded case and compute in the Grothendieck group.

$$\begin{aligned} & [\mathcal{E}_i][\mathcal{E}_{i-1}][\mathcal{E}_i][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & 2[P(k-1, \dots, k-1, \dots, \underbrace{i, \dots, i}_{d_{i-1}}, \underbrace{i-1, \dots, i-1, i, i}_{d_{i-2}}, i-2, \dots, i-2, i-1, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i, \dots, i}_{d_{i-1}}, \underbrace{i-1, \dots, i-1, i, i}_{d_{i-2}}, i-2, \dots, i-2, i, \dots, 0, \dots, 0)]. \end{aligned}$$

We also have

$$\begin{aligned} & [\mathcal{E}_{i-1}][\mathcal{E}_i^{(2)}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-1, \dots, i-1, i}_{d_{i-1}}, \underbrace{i-2, \dots, i-2, i}_{d_{i-2}}, \dots, 0, \dots, 0)] \end{aligned}$$

and

$$\begin{aligned} & [\mathcal{E}_i^{(2)}][\mathcal{E}_{i-1}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \underbrace{i-1, \dots, i-1}_{d_{i-1}}, \underbrace{i, i, i-2, \dots, i-2}_{d_{i-2}}, \dots, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \underbrace{i-1, \dots, i-1}_{d_{i-1}}, \underbrace{i, i-2, \dots, i-2}_{d_{i-2}}, \dots, 0, \dots, 0)]. \end{aligned}$$

Therefore we get the desired isomorphism of functors if we ignore the grading. The functor $\tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i-1}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}-\epsilon_{i-2}-\epsilon_{i-1}+2\epsilon_i})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i-1}-2\epsilon_{i-2}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}, C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}}\langle -d_{i-2}+1 \rangle) \otimes_{A_{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}} \\ & \text{Hom}_{C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}}(\mathbb{V}P_{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}, C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}} \mathbb{V}P_{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}\langle -2d_{i-1}+2 \rangle). \end{aligned}$$

Therefore we must consider

$$X = C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}\langle -2d_{i-1}-d_{i-2}+3 \rangle$$

as a $C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}$ -module.

As a $C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}$ -module, $C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}$ is free of rank $|W^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}/W^{\mathbf{d}; \mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}|$ and a basis can be chosen in degrees twice the length of minimal coset representatives. Therefore we would like to consider minimal coset representatives in $S_{d_{i-1}+1}/S_{d_{i-1}} \times S_1$. Proposition A-2 of [Str] tells us the length of these coset representatives. We get $C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}} \cong \bigoplus_{t=0}^{d_{i-1}-1} C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}}\langle 2t \rangle$.

Similarly, as a $C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}$ -module, $C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_{i-2}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \dots \times S_{d_i+2} \times S_{d_{i-1}-1} \times S_{d_{i-2}-1} \times \dots \times S_{d_0}/S_{d_{k-1}} \times \dots \times S_{d_i} \times S_2 \times S_{d_{i-1}-1} \times S_{d_{i-2}-1} \times \dots \times S_{d_0}.$$

Therefore,

$$X \cong \bigoplus_{t=0}^{d_{i-1}-1} \bigoplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}\langle 2r+2s+2t-2d_{i-1}-d_{i-2}+1 \rangle.$$

Next, $\tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i^{(2)}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}}\langle -2d_{i-1}+4 \rangle) \otimes_{A_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}}(\mathbb{V}P_{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}, C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}\langle -d_{i-2}+1 \rangle). \end{aligned}$$

Thus we must consider

$$Y = C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}\langle -2d_{i-1}-d_{i-2}+5 \rangle.$$

As a $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ -module, $C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives from

$$S_{d_{k-1}} \times \dots \times S_{d_i+2} \times S_{d_{i-1}-2} \times \dots \times S_{d_0}/S_{d_{k-1}} \times \dots \times S_{d_i} \times S_2 \times S_{d_{i-1}-2} \times \dots \times S_{d_0}.$$

So now we want to consider minimal coset representatives in $S_{d_i+2}/S_{d_i} \times S_2$. Proposition A-2 of [Str] gives

$$C^{\mathbf{d}; \mathbf{d}+2\epsilon_i-2\epsilon_{i-1}} \cong \bigoplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}}\langle 2r+2s-2 \rangle.$$

Similarly, as a $C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}$ -module, $C^{\mathbf{d}+2\epsilon_i-2\epsilon_{i-1}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \dots \times S_{d_i+2} \times S_{d_{i-1}-1} \times S_{d_{i-2}-1} \times \dots \times S_{d_0}/S_{d_{k-1}} \times \dots \times S_{d_i+2} \times S_{d_{i-1}-2} \times S_1 \times S_{d_{i-2}-1} \times \dots \times S_{d_0}.$$

Therefore,

$$Y \cong \bigoplus_{t=0}^{d_{i-1}-2} \bigoplus_{d_i+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}\langle 2r+2s+2t-2d_{i-1}-d_{i-2}+3 \rangle.$$

Finally, $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}}\langle -d_{i-1}+1 \rangle) \otimes_{A_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-2}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i-2}}, C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+\epsilon_i-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}\langle -d_{i-2}+1 \rangle) \otimes_{A_{\mathbf{d}+\epsilon_i-\epsilon_{i-2}}} \\ & \text{Hom}_{C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}}}(\mathbb{V}P_{\mathbf{d}+2\epsilon_i+\epsilon_{i-1}-\epsilon_{i-2}}, C^{\mathbf{d}+\epsilon_i-\epsilon_{i-2}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-2\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-2}}} \mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i-2}}\langle -d_{i-1}+1 \rangle). \end{aligned}$$

Thus we must consider

$$W = C^{\mathbf{d}+\epsilon_i-\epsilon_{i-2}; \mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-2}}} C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}; \mathbf{d}+\epsilon_i-\epsilon_{i-2}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \langle -2d_{i-1}-d_{i-2}+3 \rangle.$$

As in all the other computations,

$$W \cong \oplus_{t=0}^{d_{i-1}-1} \oplus_{r=0}^{d_i} \oplus_{s=0}^{d_i+1} C^{\mathbf{d}+2\epsilon_i-\epsilon_{i-1}-\epsilon_{i-2}} \langle 2r+2s+2t-2d_{i-1}-d_{i-1}+3 \rangle.$$

The grading for $\tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i-1} \oplus \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i^{(2)}$ is given by

$$\begin{aligned} & \sum_{t=0}^{d_{i-1}} \sum_{d_i+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_{i-1}-d_{i-2}+1} + \sum_{t=0}^{d_{i-1}-2} \sum_{d_i+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_{i-1}-d_{i-2}+3} = \\ & \sum_{t=0}^{d_{i-1}-1} \sum_{d_i+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_{i-1}-d_{i-2}+1} + \sum_{t=0}^{d_{i-1}-1} \sum_{d_i+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_{i-1}-d_{i-2}+3} = \\ & (1+q^{-2}) \sum_{t=0}^{d_{i-1}-1} \sum_{d_i+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_{i-1}-d_{i-2}+3} = \\ & \sum_{t=0}^{d_{i-1}-1} \sum_{r=0}^{d_i+1} \sum_{s=0}^{d_i} q^{2r+2s+2t-2d_{i-1}-d_{i-2}+3}. \end{aligned}$$

This is the grading for $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i$ so we have the desired isomorphism of graded functors. $\square$

Lemma 35. *For some indecomposable projective functor G_3 , there are isomorphisms of graded functors:*

- (1) $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i^{(2)} \oplus \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i^{(2)} \oplus \tilde{G}_3.$
- (2) $\tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i+1} \cong \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i^{(2)} \oplus \tilde{G}_3.$
- (3) $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i+1} \oplus \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i^{(2)}.$

Proof. First we consider the ungraded case and compute in the Grothendieck group.

$$\begin{aligned} & [\mathcal{F}_i][\mathcal{F}_{i+1}][\mathcal{F}_i][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & 2[P(k-1, \dots, k-1, \dots, \underbrace{i, i+1, i+1, \dots, i+1}_{d_{i+1}}, \underbrace{i-1, i-1, i, \dots, i-1}_{d_i}, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-1, i+1, i+1, \dots, i+1}_{d_i}, \dots, 0, \dots, 0)]. \end{aligned}$$

We also have

$$\begin{aligned} & [\mathcal{F}_{i+1}][\mathcal{F}_i^{(2)}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i, i+1, i+1, \dots, i+1}_{d_{i+1}}, \underbrace{i-1, i-1, i, \dots, i}_{d_i}, \dots, 0, \dots, 0)] \end{aligned}$$

and

$$\begin{aligned} & [\mathcal{F}_i^{(2)}][\mathcal{F}_{i+1}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-1, i+1, i+1, \dots, i+1}_{d_{i+1}}, \underbrace{i-1, i, \dots, i}_{d_i}, \dots, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i, i+1, i+1, \dots, i+1}_{d_i}, \underbrace{i-1, i-1, i, \dots, i}_{d_i}, \dots, 0, \dots, 0)]. \end{aligned}$$

Therefore we get the desired isomorphism of functors if we ignore the grading.

The functor $\tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i+1}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}+\epsilon_i-2\epsilon_{i+1}}}(\mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i+1}}, C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -d_{i+1}+1 \rangle) \otimes_{A_{\mathbf{d}+\epsilon_i-\epsilon_{i+1}}} \\ & \text{Hom}_{C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}}(\mathbb{V}P_{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}, C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}}} \mathbb{V}P_{\mathbf{d}+\epsilon_i-\epsilon_{i+1}} \langle -2d_i+2 \rangle). \end{aligned}$$

Therefore we must consider

$$X = C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i+1}} \langle -2d_i - d_{i+1} + 3 \rangle$$

as a $C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} -$ module.

As a $C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}} -$ module, $C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i+1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives. Therefore we would like to consider minimal coset representatives in $S_{d_{i+1}}/S_1 \times S_{d_i}$. Proposition A-2 of [Str] tells us the length of these coset representatives. We get $C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i+1}} \cong \bigoplus_{t=0}^{d_i} C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}} \langle 2t \rangle$.

Similarly, as a $C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} -$ module, $C^{\mathbf{d}+\epsilon_i-\epsilon_{i+1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \cdots \times S_{d_{i+1}-1} \times S_{d_i-1} \times S_{d_{i-1}+2} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i+1}-1} \times S_{d_i-1} \times S_2 \times S_{d_{i-1}} \times \cdots \times S_{d_0}.$$

Therefore,

$$X \cong \bigoplus_{t=0}^{d_i} \bigoplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \langle 2r + 2s + 2t - 2d_i - d_{i+1} + 3 \rangle.$$

Next, $\tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i^{(2)}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}) -$ bimodule

$$\text{Hom}_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}} (\mathbb{V}P_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -2d_i + 4 \rangle) \otimes_{A_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}}$$

$$\text{Hom}_{C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}} (\mathbb{V}P_{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}, C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle -d_{i+1} + 1 \rangle).$$

Thus we must consider

$$Y = C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle -2d_i - d_{i+1} + 5 \rangle.$$

As a $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} -$ module, $C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \cdots \times S_{d_{i-2}} \times S_{d_{i-1}+2} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i-2}} \times S_2 \times S_{d_{i-1}} \times \cdots \times S_{d_0}.$$

So now we want to consider minimal coset representatives in $S_{d_{i-1}+2}/S_2 \times S_{d_{i-1}}$. Proposition A-2 of [Str] gives

$$C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \cong \bigoplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle 2r + 2s - 2 \rangle.$$

Similarly, as a $C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} -$ module, $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \cdots \times S_{d_{i+1}-1} \times S_{d_i-1} \times S_{d_{i-1}+2} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i+1}-1} \times S_1 \times S_{d_i-2} \times S_{d_{i-1}+2} \times \cdots \times S_{d_0}.$$

Therefore,

$$Y \cong \bigoplus_{t=0}^{d_i-2} \bigoplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \langle 2r + 2s + 2t - 2d_i - d_{i+1} + 3 \rangle.$$

Finally, $\tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}) -$ bimodule

$$\text{Hom}_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} (\mathbb{V}P_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}-\epsilon_i+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -d_i + 1 \rangle) \otimes_{A_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}}$$

$$\text{Hom}_{C^{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}}} (\mathbb{V}P_{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}}, C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}; \mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}} \langle -d_{i+1} + 1 \rangle) \otimes_{A_{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}}}$$

$$\text{Hom}_{C^{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}} (\mathbb{V}P_{\mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}}, C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}} \langle -d_i + 1 \rangle).$$

Thus we must consider

$$W = C^{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}; \mathbf{d}+2\epsilon_{i-1}-\epsilon_i-\epsilon_{i+1}} \otimes_{C^{\mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}}} C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}; \mathbf{d}-\epsilon_{i+1}+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}-\epsilon_i+\epsilon_{i-1}} \langle -2d_i - d_{i+1} + 3 \rangle.$$

As in all the other computations,

$$W \cong \bigoplus_{t=0}^{d_i-1} \bigoplus_{r=0}^{d_i-1} \bigoplus_{s=0}^{d_i-1} C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \langle 2r + 2s + 2t - 2d_i - d_{i+1} + 3 \rangle.$$

The grading for $\tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i+1} \oplus \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i^{(2)}$ is given by the summation

$$\begin{aligned} & \sum_{t=0}^{d_i} \sum_{d_{i-1}+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i+1}+1} + \sum_{t=0}^{d_i-2} \sum_{d_{i-1}+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i+1}+3} = \\ & \sum_{t=0}^{d_i-1} \sum_{d_{i-1}+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i+1}+1} + \sum_{t=0}^{d_i-1} \sum_{d_{i-1}+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i+1}+3} = \end{aligned}$$

$$(q^{-2} + 1) \sum_{t=0}^{d_i-1} \sum_{d_{i-1}+1 \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i-1}+3}.$$

This is the grading for $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i$ so we have the desired isomorphism of graded functors. $\square$

Lemma 36. *For some indecomposable projective functor G_4 , there are isomorphisms of graded functors:*

- (1) $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i-1} \oplus \tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i-1} \oplus \tilde{G}_4.$
- (2) $\tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i^{(2)} \cong \tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i-1} \oplus \tilde{G}_4.$
- (3) $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i-1} \oplus \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i^{(2)}.$

Proof. First we consider the ungraded case and compute in the Grothendieck group.

$$\begin{aligned} & [\mathcal{F}_i][\mathcal{F}_{i-1}][\mathcal{F}_i][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & 2[P(k-1, \dots, k-1, \dots, \underbrace{i-1, i-1, i, \dots, i}_{d_i}, \underbrace{i-2, i-1, \dots, i-1}_{d_{i-1}}, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-2, i-1, i, \dots, i}_{d_{i-1}}, \dots, 0, \dots, 0)]. \end{aligned}$$

We also have

$$\begin{aligned} & [\mathcal{F}_i^{(2)}][\mathcal{F}_{i-1}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-2, i-1, i, \dots, i}_{d_i}, \underbrace{i-2, i-1, \dots, i-1}_{d_{i-1}}, \dots, 0, \dots, 0)] \end{aligned}$$

and

$$\begin{aligned} & [\mathcal{F}_{i-1}][\mathcal{F}_i^{(2)}][M(k-1, \dots, k-1, \dots, 0, \dots, 0)] = \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-1, i-1, i, \dots, i}_{d_i}, \underbrace{i-2, i-1, \dots, i-1}_{d_{i-1}}, \dots, 0, \dots, 0)] + \\ & [P(k-1, \dots, k-1, \dots, \underbrace{i-2, i-1, i, \dots, i}_{d_i}, \underbrace{i-1, \dots, i-1}_{d_{i-1}}, \dots, 0, \dots, 0)]. \end{aligned}$$

Therefore we get the desired isomorphism of functors if we ignore the grading.

The functor $\tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i^{(2)}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i})$ -bimodule

$$\begin{aligned} & \text{Hom}_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}} \langle -2d_i + 4 \rangle) \otimes_{A_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}} \\ & \text{Hom}_{C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}, C^{\mathbf{d}+2\epsilon_{i-1}-2\epsilon_i; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \otimes_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle -d_{i-1} - 1 \rangle). \end{aligned}$$

Therefore we must consider

$$X = C^{\mathbf{d}+2\epsilon_{i-1}-2\epsilon_i; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \otimes_{C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle -2d_i - d_{i-1} + 3 \rangle$$

as a $C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}$ -module.

As a $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}$ -module, $C^{\mathbf{d}, \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}$ is free of rank $|W^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}/W^{\mathbf{d}, \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}}|$ and a basis can be chosen in degrees twice the length of minimal coset representatives. Therefore we would like to consider minimal coset representatives in $S_{d_{i-1}+2}/S_2 \times S_{d_{i-1}}$. Proposition A-2 of [Str] tells us the length of these coset representatives. We get $C^{\mathbf{d}, \mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \cong \bigoplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}} \langle 2r + 2s - 2 \rangle$.

Similarly, as a $C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}$ -module, $C^{\mathbf{d}-2\epsilon_i+2\epsilon_{i-1}, \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_k} \times \dots \times S_{d_{i-2}} \times S_{d_{i-1}+1} \times S_{d_{i-2}+1} \times \dots \times S_{d_0}/S_{d_k} \times \dots \times S_{d_{i-2}} \times S_{d_{i-1}+1} S_1 \times S_{d_{i-2}} \times \dots \times S_{d_0}.$$

Therefore,

$$X \cong \bigoplus_{t=0}^{d_{i-2}} \bigoplus_{d_{i-1}+1 \geq r > s \geq 0} C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \langle 2r + 2s + 2t - 2d_i - d_{i-1} + 1 \rangle.$$

Next, $\tilde{\mathcal{F}}_i^{(2)} \tilde{\mathcal{F}}_{i-1}$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i})$ -bimodule

$$\text{Hom}_{C^{\mathbf{d}-\epsilon_{i-1}+\epsilon_{i-2}}}(\mathbb{V}P_{\mathbf{d}-\epsilon_{i-1}+\epsilon_{i-2}}, C^{\mathbf{d}; \mathbf{d}-\epsilon_{i-1}+\epsilon_{i-2}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}}\langle -d_{i-1}+1 \rangle) \otimes_{A_{\mathbf{d}-\epsilon_{i-1}+\epsilon_{i-2}}}$$

$$\text{Hom}_{C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}, C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \otimes_{C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}-\epsilon_{i-2}-\epsilon_{i-1}}\langle -2d_i+4 \rangle).$$

Thus we must consider

$$Y = C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}+2\epsilon_i} \otimes_{C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}}\langle -2d_i-d_{i-1}+5 \rangle$$

as a $C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}$ -module. As a $C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}}$ -module, $C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \cdots \times S_{d_{i-1}-1} \times S_{d_{i-2}+1} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i-1}-1} \times S_1 \times S_{d_{i-2}} \times \cdots \times S_{d_0}.$$

So now we want to consider minimal coset representatives in $S_{d_{i-2}+1}/S_1 \times S_{d_{i-2}}$. Proposition A-2 of [Str] gives

$$C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}} \cong \bigoplus_{t=0}^{d_{i-2}} C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}}\langle 2t \rangle.$$

Similarly, as a $C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}$ -module, $C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_{i-1}; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}$ is free and a basis can be chosen in degrees twice the length of minimal coset representatives of

$$S_{d_{k-1}} \times \cdots \times S_{d_{i-2}} \times S_{d_{i-1}+1} \times S_{d_{i-2}+1} \times \cdots \times S_{d_0}/S_{d_{k-1}} \times \cdots \times S_{d_{i-2}} \times S_2 \times S_{d_{i-1}-1} \times S_{d_{i-2}+1} \times \cdots \times S_{d_0}.$$

Therefore,

$$Y \cong \bigoplus_{t=0}^{d_{i-2}} \bigoplus_{d_{i-1} \geq r > s \geq 0} C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}\langle 2r+2s+2t-2d_i-d_{i-1}+3 \rangle.$$

Finally, $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i$ is given by tensoring with the graded $(A_{\mathbf{d}}, A_{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i})$ -bimodule

$$\text{Hom}_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}}(\mathbb{V}P_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}, C^{\mathbf{d}; \mathbf{d}-\epsilon_i+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{V}P_{\mathbf{d}}\langle -d_i+1 \rangle) \otimes_{A_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}}$$

$$\text{Hom}_{C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_i}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i-2}-\epsilon_i}, C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}; \mathbf{d}-\epsilon_i+\epsilon_{i-2}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} \mathbb{V}P_{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}\langle -d_{i-1} \rangle) \otimes_{A_{\mathbf{d}+\epsilon_{i-2}-\epsilon_i}}$$

$$\text{Hom}_{C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}}(\mathbb{V}P_{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}, C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_i; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \otimes_{C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_i}} \mathbb{V}P_{\mathbf{d}+\epsilon_{i-2}-\epsilon_i}\langle -d_i+2 \rangle).$$

Thus we must consider

$$W = C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_i; \mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i} \otimes_{C^{\mathbf{d}+\epsilon_{i-2}-\epsilon_i}} C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_i; \mathbf{d}+\epsilon_{i-2}-\epsilon_i} \otimes_{C^{\mathbf{d}+\epsilon_{i-1}-\epsilon_i}} C^{\mathbf{d}; \mathbf{d}+\epsilon_{i-1}-\epsilon_i}\langle -2d_i-d_{i-1}+3 \rangle.$$

As in all the other computations,

$$W \cong \bigoplus_{t=0}^{d_{i-2}} \bigoplus_{r=0}^{d_{i-1}} \bigoplus_{s=0}^{d_{i-1}} C^{\mathbf{d}+\epsilon_{i-2}+\epsilon_{i-1}-2\epsilon_i}\langle 2r+2s+2t-2d_i-d_{i-1}+3 \rangle.$$

The grading for $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i$ is given by

$$\begin{aligned} & \sum_{t=0}^{d_{i-2}} \sum_{r=0}^{d_{i-1}} \sum_{s=0}^{d_{i-1}} q^{2r+2s+2t-2d_i-d_{i-1}+3} = \\ & 2 \sum_{t=0}^{d_{i-2}} \sum_{d_{i-1} \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i-1}+3} + \sum_{t=0}^{d_{i-2}} \sum_{r=0}^{d_{i-1}} q^{4r+2t-2d_i-d_{i-1}+3}. \end{aligned}$$

The grading for $\tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i^{(2)}$ is given by

$$\sum_{t=0}^{d_{i-2}} \sum_{d_{i-1}+1 \geq r > s \geq 0} q^{2(r-1)+2s+2t-2d_i-d_{i-1}+3}.$$

Letting $u = r - 1$, this is equal to

$$\begin{aligned} & \sum_{t=0}^{d_{i-2}} \sum_{d_{i-1} \geq u \geq s \geq 0} q^{2u+2s+2t-2d_i-d_{i-1}+3} = \\ & \sum_{t=0}^{d_{i-2}} \sum_{d_{i-1} \geq u > s \geq 0} q^{2u+2s+2t-2d_i-d_{i-1}+3} + \sum_{t=0}^{d_{i-2}} \sum_{u=0}^{d_{i-1}} q^{4u+2t-2d_i-d_{i-1}+3}. \end{aligned}$$

Since the grading for $\tilde{\mathcal{F}}_i^{(2)}$ is given by $\sum_{t=0}^{d_{i-2}} \sum_{d_{i-1} \geq r > s \geq 0} q^{2r+2s+2t-2d_i-d_{i-1}+3}$, we get the desired isomorphism of graded functors. $\square$

Now we may state a categorification of the quantum Serre relations.

Theorem 5. *There are isomorphisms of graded projective functors:*

- (1) $\tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_i^{-1} \cong \text{Id} \cong \tilde{\mathcal{H}}_i^{-1} \tilde{\mathcal{H}}_i.$
- (2) $\tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j \cong \tilde{\mathcal{H}}_j \tilde{\mathcal{H}}_i.$
- (3) If $|i - j| > 1$, $\tilde{\mathcal{H}}_i \tilde{\mathcal{E}}_j \cong \tilde{\mathcal{E}}_j \tilde{\mathcal{H}}_i.$
- (4) $\tilde{\mathcal{H}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{H}}_i \langle 2 \rangle.$
- (5) If $|i - j| = 1$, $\tilde{\mathcal{H}}_i \tilde{\mathcal{E}}_j \cong \tilde{\mathcal{E}}_j \tilde{\mathcal{H}}_i \langle -1 \rangle.$
- (6) If $|i - j| > 1$, $\tilde{\mathcal{H}}_i \tilde{\mathcal{F}}_j \cong \tilde{\mathcal{F}}_j \tilde{\mathcal{H}}_i.$
- (7) $\tilde{\mathcal{H}}_i \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i \tilde{\mathcal{H}}_i \langle -2 \rangle.$
- (8) If $|i - j| = 1$, $\tilde{\mathcal{H}}_i \tilde{\mathcal{F}}_j \cong \tilde{\mathcal{F}}_j \tilde{\mathcal{H}}_i \langle 1 \rangle.$
- (9) If $i \neq j$, $\tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_j \cong \tilde{\mathcal{F}}_j \tilde{\mathcal{E}}_i.$
- (10) If $|i - j| > 1$, $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_j \cong \tilde{\mathcal{E}}_j \tilde{\mathcal{E}}_i.$
- (11) If $|i - j| > 1$, $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_j \cong \tilde{\mathcal{F}}_j \tilde{\mathcal{F}}_i.$
- (12) $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \oplus \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \langle 1 \rangle \oplus \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \langle -1 \rangle.$
- (13) $\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \oplus \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i \langle 1 \rangle \oplus \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i-1} \tilde{\mathcal{E}}_i \langle -1 \rangle.$
- (14) $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i+1} \oplus \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i \langle 1 \rangle \oplus \tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i+1} \tilde{\mathcal{F}}_i \langle -1 \rangle.$
- (15) $\tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \oplus \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i \langle 1 \rangle \oplus \tilde{\mathcal{F}}_i \tilde{\mathcal{F}}_{i-1} \tilde{\mathcal{F}}_i \langle -1 \rangle.$
- (16) If $d_{i-1} > d_i$ then

$$\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i \oplus_{r=0}^{d_{i-1}-d_i-1} \text{Id} \langle d_{i-1} - d_i - 1 - 2r \rangle.$$

If $d_i > d_{i-1}$ then

$$\tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \oplus_{r=0}^{d_i-d_{i-1}-1} \text{Id} \langle d_i - d_{i-1} - 1 - 2r \rangle.$$

If $d_i = d_{i-1}$, then $\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i.$

Proof. (1) This is obvious.

(2) This is obvious.

(3) If $|i - j| > 1$, then the value of $|d_i - d_{i-1}|$ remains unchanged after applying $\tilde{\mathcal{E}}_j$.

(4) After applying $\tilde{\mathcal{E}}_i$, d_i increases by 1 and d_{i-1} decreases by 1 so there is a shift of 2.

(5) If $j = i + 1$, then d_i decreases by 1 and d_{i-1} remains unchanged. If $j = i - 1$, then d_{i-1} increases by 1 and d_i remains unchanged so there is a shift of -1.

(6) This is the same as the proof of 3.

(7) After applying $\tilde{\mathcal{F}}_i$, d_i decreases by 1 and d_{i-1} increases by 1 so there is a shift of -2.

(8) If $j = i + 1$, then d_i increases by 1 and d_{i-1} remains unchanged. If $j = i - 1$, then d_{i-1} decreases by 1 and d_i remains unchanged.

(9) We would like to prove that $\mathcal{E}_i \mathcal{F}_j$ and $\mathcal{F}_j \mathcal{E}_i$ are indecomposable projective functors if $i \neq j$. We show this by computing $[\mathcal{E}_i \mathcal{F}_j]$ and $[\mathcal{F}_j \mathcal{E}_i]$ on $[M(k-1, \dots, k-1, \dots, 0, \dots, 0)]$. If $i - j > 1$, both are equal to

$$[P(k-1, \dots, k-1, \dots, \underbrace{i, \dots, i}_{d_{i-1}}, \underbrace{i-1, \dots, i-1}_{d_j}, \dots, j-1, j, \dots, j, \dots, 0, \dots, 0)].$$

The case of $i - j < 1$ is similar. If $i - j = 1$, then both are equal to

$$[P(k-1, \dots, k-1, \dots, \underbrace{i, \dots, i}_{d_{i-1}}, i-2, i-1, i-1, \dots, i-1, i, i-2, \dots, i-2, \dots, 0, \dots, 0)].$$

The case of $i - j = -1$ is similar. Thus $\mathcal{E}_i \mathcal{F}_j$ and $\mathcal{F}_j \mathcal{E}_i$ are indecomposable projective functors that are equal on the Grothendieck group so they are isomorphic. Since their lowest and highest gradings obviously coincide, their graded lifts are also isomorphic to each other with no shift.

(10) Again, we first compute the ungraded functors on the dominant generalized Verma module in the Grothendieck group to get

$$[P(k-1, \dots, k-1, \dots, \underbrace{j-1, \dots, j-1}_{d_{j-1}}, \underbrace{j, \dots, j}_{d_{i-1}}, \dots, i-1, i-1, \dots, i-1, i, \dots, 0, \dots, 0)].$$

Thus $\mathcal{E}_i \mathcal{E}_j$ and $\mathcal{E}_j \mathcal{E}_i$ are indecomposable isomorphic projective functors. Since their lowest and highest gradings coincide, their graded lifts are isomorphic with no shift.

- (11) The proof of this is the same as the previous one. Applying these functors to the dominant generalized Verma module in the Grothendieck group gives

$$[P(k-1, \dots, k-1, \dots, \underbrace{j-1, j, \dots, j}_{d_j}, \dots, \underbrace{i-1, i, \dots, i}_{d_i}, \dots, 0, \dots, 0)].$$

The rest of the proof is the same.

- (12) Lemma 31 implies both

$$\begin{aligned} \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i &\cong \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)} \langle 1 \rangle \oplus \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)} \langle -1 \rangle \\ \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} &\cong \tilde{\mathcal{E}}_i^{(2)} \langle 1 \rangle \tilde{\mathcal{E}}_{i+1} \oplus \tilde{\mathcal{E}}_i^{(2)} \langle -1 \rangle \tilde{\mathcal{E}}_{i+1}. \end{aligned}$$

Then lemma 33 gives

$$\tilde{\mathcal{E}}_i \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i^{(2)} \tilde{\mathcal{E}}_{i+1} \oplus \tilde{\mathcal{E}}_{i+1} \tilde{\mathcal{E}}_i^{(2)}.$$

These facts now easily prove the claim.

- (13) This is similar to the proof of 12 using lemmas 31 and 34.
(14) This is similar to the proof of 12 using lemmas 32 and 35.
(15) This is similar to the proof of 12 using lemmas 32 and 36.
(16) We compute $[\mathcal{E}_i][\mathcal{F}_i](M(k-1, \dots, k-1, \dots, 0, \dots, 0))$. In general,

$$[\mathcal{E}_i][\mathcal{F}_i](M(a_1, \dots, a_n)) = \sum_{r=1; a_r=i}^n \sum_{s=1; a_s=i-1}^n [M(a_1, \dots, a'_r, \dots, a''_s, \dots, a_n)] + \sum_{r=1; a_r=i}^n [M(a_1, \dots, a_n)]$$

where $a'_r = i-1$ and $a''_s = i$. Applied to the dominant Verma module it is equal to

$$[P(k-1, \dots, k-1, \dots, \underbrace{i-1, i, \dots, i}_{d_i}, \underbrace{i-1, i-1, \dots, i-1, i}_{d_{i-1}}, 0, \dots, 0)] + \sum_{r=1; a_r=i}^n [M(k-1, \dots, 0)].$$

On the other hand,

$$[\mathcal{F}_i][\mathcal{E}_i](M(a_1, \dots, a_n)) = \sum_{r=1; a_r=i}^n \sum_{s=1; a_s=i-1}^n [M(a_1, \dots, a'_r, \dots, a''_s, \dots, a_n)] + \sum_{s=1; a_s=i-1}^n [M(a_1, \dots, a_n)].$$

So for the dominant Verma module it is equal to

$$[P(k-1, \dots, k-1, \dots, \underbrace{i-1, i, \dots, i}_{d_i}, \underbrace{i-1, i-1, \dots, i-1, i}_{d_{i-1}}, 0, \dots, 0)] + \sum_{s=1; a_s=i-1}^n [M(k-1, \dots, 0)].$$

Thus

$$\mathcal{E}_i \mathcal{F}_i \cong G \oplus \bigoplus_{r=0}^{d_i-1} \text{Id}$$

and

$$\mathcal{F}_i \mathcal{E}_i \cong G \oplus \bigoplus_{s=0}^{d_{i-1}-1} \text{Id}$$

where G is an indecomposable projective functor.

As in lemma 3.4 of [FKS], as a $C^{\mathbf{d}}$ -module, $C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \cong \bigoplus_{r=0}^{d_{i-1}-1} C^{\mathbf{d}} \langle 2r \rangle$ and $C^{\mathbf{d}; \mathbf{d}-\epsilon_i+\epsilon_{i-1}} \cong \bigoplus_{r=0}^{d_i-1} C^{\mathbf{d}} \langle 2r \rangle$. Thus

$$(2) \quad C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}+\epsilon_i-\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}+\epsilon_i-\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{C} \cong \bigoplus_{l=0}^{d_i} \bigoplus_{k=0}^{d_{i-1}-1} \mathbb{C} \langle 2k+2l \rangle$$

$$(3) \quad C^{\mathbf{d}; \mathbf{d}-\epsilon_i+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}-\epsilon_i+\epsilon_{i-1}}} C^{\mathbf{d}; \mathbf{d}-\epsilon_i+\epsilon_{i-1}} \otimes_{C^{\mathbf{d}}} \mathbb{C} \cong \bigoplus_{l=0}^{d_{i-1}} \bigoplus_{k=0}^{d_i-1} \mathbb{C} \langle 2k+2l \rangle.$$

Since their lowest degrees coincide and G is indecomposable, there is an isomorphism of graded functors

$$\tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i \cong \tilde{G} \oplus \bigoplus_{r=0}^{d_i-1} \text{Id} \langle m_r \rangle$$

and

$$\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \cong \tilde{G} \oplus \bigoplus_{s=0}^{d_{i-1}-1} \text{Id} \langle n_r \rangle$$

Now $\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i$ is given by equation 2 but shifted by $\langle -(d_i-1) \rangle \langle -(d_{i-1}) \rangle$ and $\tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i$ is given by equation 3 but shifted by $\langle -(d_{i-1}) \rangle \langle -(d_i-1) \rangle$.

Hence $[\tilde{\mathcal{F}}_i][\tilde{\mathcal{E}}_i] - [\tilde{\mathcal{E}}_i][\tilde{\mathcal{F}}_i] =$

$$\sum_{l=0}^{d_i} \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - 2l + d_i + d_{i-1} - 1 \rangle] - \sum_{k=0}^{d_{i-1}} \sum_{l=0}^{d_i-1} [\mathbb{C}\langle -2k - 2l + d_i + d_{i-1} - 1 \rangle].$$

Suppose $d_i < d_{i-1}$. Then the above is equal to

$$\begin{aligned} & \sum_{l=0}^{d_{i-1}-1} \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - 2l + d_i + d_{i-1} - 1 \rangle] + \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - 2d_i + d_i + d_{i-1} - 1 \rangle] - \\ & \sum_{k=0}^{d_{i-1}-1} \sum_{l=0}^{d_i-1} [\mathbb{C}\langle -2k - 2l + d_i + d_{i-1} - 1 \rangle] - \sum_{l=0}^{d_i-1} [\mathbb{C}\langle -2l - 2d_{i-1} + d_i + d_{i-1} - 1 \rangle] = \\ & \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - 2d_i + d_i + d_{i-1} - 1 \rangle] - \sum_{l=0}^{d_i-1} [\mathbb{C}\langle -2l - 2d_{i-1} + d_i + d_{i-1} - 1 \rangle] = \\ & \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - d_i + d_{i-1} - 1 \rangle] - \sum_{l=0}^{d_i-1} [\mathbb{C}\langle -2l - d_{i-1} + d_i - 1 \rangle]. \end{aligned}$$

Let $m = l - d_i + d_{i-1}$. Then the above is equal to

$$\begin{aligned} & \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - d_i + d_{i-1} - 1 \rangle] - \sum_{m=d_{i-1}-d_i}^{d_{i-1}-1} [\mathbb{C}\langle -2m - 2d_i + 2d_{i-1} + d_i - d_{i-1} - 1 \rangle] = \\ & \sum_{k=0}^{d_{i-1}-1} [\mathbb{C}\langle -2k - d_i + d_{i-1} - 1 \rangle] - \sum_{m=d_{i-1}-d_i}^{d_{i-1}-1} [\mathbb{C}\langle -2m + d_{i-1} - d_i - 1 \rangle] = \\ & \sum_{k=0}^{d_{i-1}-d_i-1} [\mathbb{C}\langle -2k + d_{i-1} - d_i - 1 \rangle] = [d_{i-1} - d_i]. \end{aligned}$$

Thus

$$\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i \oplus_{r=0}^{d_{i-1}-d_i-1} \text{Id}\langle d_{i-1} - d_i - 1 - 2r \rangle.$$

Suppose $d_i = d_{i-1}$. Then clearly $\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \cong \tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i$.

Finally suppose $d_i > d_{i-1}$. As in the case $d_{i-1} > d_i$,

$$[\tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i] - [\tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i] = [d_i - d_{i-1}].$$

Thus

$$\tilde{\mathcal{E}}_i \tilde{\mathcal{F}}_i \cong \tilde{\mathcal{F}}_i \tilde{\mathcal{E}}_i \oplus_{r=0}^{d_i-d_{i-1}-1} \text{Id}\langle d_i - d_{i-1} - 1 - 2r \rangle.$$

□

5.2. Graded Equivalence of Categories. We would like to prove a graded version of the equivalence given in section 3. In the ungraded case there is an isomorphism of functors

$$LZ_{\mathfrak{q}_j}^{\mathfrak{p}_j} \epsilon_{\mathfrak{p}_{j+1}}^{\mathfrak{q}_j} [-(k-1)] LZ_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} \epsilon_{\mathfrak{p}_j}^{\mathfrak{q}_j} [-(k-1)] \cong \text{Id}.$$

Thus in the graded case there is an isomorphism for some shift a :

$$L\tilde{Z}_{\mathfrak{q}_j}^{\mathfrak{p}_j} \tilde{\epsilon}_{\mathfrak{p}_{j+1}}^{\mathfrak{q}_j} [-(k-1)] L\tilde{Z}_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} \tilde{\epsilon}_{\mathfrak{p}_j}^{\mathfrak{q}_j} [-(k-1)] \cong \text{Id}\langle a \rangle.$$

To determine the shift, we compute on a generalized Verma module.

Lemma 37. *In the notation of section 3.2, $L_s \tilde{Z}_{\mathfrak{q}_j}^{\mathfrak{p}_{j+1}} \tilde{\epsilon}_{\mathfrak{p}_j}^{\mathfrak{q}_j} \tilde{M}^{\mathfrak{p}_j}(\alpha) \cong \tilde{M}^{\mathfrak{p}_{j+1}}(\beta)$ if $s = k - 1$ and 0 otherwise.*

Proof. We know that it is zero for $s \neq k-1$ so we only need to concentrate on the case that $s = k-1$. Consider the graded sequences from lemma 5:

$$\begin{aligned}
0 &\rightarrow \tilde{K}_0 \rightarrow \tilde{M}^{q_j}(\alpha) \rightarrow \tilde{M}^{p_j}(\alpha) \rightarrow 0. \\
0 &\rightarrow \tilde{K}_1 \rightarrow \tilde{M}^{q_j}(\sigma_1.\alpha)\langle 1 \rangle \rightarrow \tilde{K}_0 \rightarrow 0. \\
0 &\rightarrow \tilde{K}_2 \rightarrow \tilde{M}^{q_j}(\sigma_1\sigma_2.\alpha)\langle 2 \rangle \rightarrow \tilde{K}_1 \rightarrow 0. \\
&\dots \\
0 &\rightarrow \tilde{K}_{k-2} \rightarrow \tilde{M}^{q_j}(\sigma_1 \cdots \sigma_{k-2}.\alpha)\langle k-2 \rangle \rightarrow \tilde{K}_{k-3} \rightarrow 0. \\
0 &\rightarrow \tilde{K}_{k-1} \rightarrow \tilde{M}^{q_j}(\sigma_1 \cdots \sigma_{k-2}\sigma_{k-1}.\alpha)\langle k-1 \rangle \rightarrow \tilde{K}_{k-2} \rightarrow 0.
\end{aligned}$$

As in the ungraded we get

$$\begin{aligned}
L_s \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{q_j}(\sigma_1 \cdots \sigma_l.\alpha)\langle l \rangle &\cong L_s \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{K}_{l-1} \cong L_{s+1} \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{K}_{l-2} \cong \cdots \cong L_{s+l-1} \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{K}_0 \\
&\cong L_{s+l} \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{p_j}(\alpha).
\end{aligned}$$

We know $L_s \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{q_j}(\sigma_1 \cdots \sigma_l.\alpha)\langle l \rangle \cong 0$ if $s \neq k-l-1$. If $s = k-l-1$, it is isomorphic to

$$\tilde{M}^{p_{j+1}}(\beta)\langle l \rangle \langle k-l-1 \rangle \langle -(k-1) \rangle \cong \tilde{M}^{p_{j+1}}(\beta).$$

The second shift comes from the fact the $\tilde{M}^{p_{j+1}}(\beta)$ is a Koszul module. The third shift comes from the grading convention for the lift of the dual Zuckerman functor. Thus there is no shift overall. $\square$

We let $\tilde{\Delta}$ denote a composition of graded equivalences.

5.3. Graded Diagram Relation 1. It suffices to repeat the calculation of lemma 8 with the grading.

Lemma 38. *If $i = 2r$,*

$$L_i \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{p_j}(\alpha) \cong \tilde{M}^{p_j}(\alpha)\langle 2r+1-k \rangle.$$

Proof. From the graded exact sequences of the previous subsection we easily see that

$$L_i \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{p_j}(\alpha) \cong \cdots \cong L_{i-r} \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{q_j}(\sigma_1 \cdots \sigma_r.\alpha)\langle r \rangle.$$

Now, $\tilde{M}^{q_j}(\sigma_1 \cdots \sigma_r.\alpha)$ is a Koszul module and has a projective resolution whose r th component is generated in degree r . Thus

$$L_{i-r} \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_{j+1}}^{q_j} \tilde{M}^{q_j}(\sigma_1 \cdots \sigma_r.\alpha)\langle r \rangle \cong \tilde{M}^{p_j}(\alpha)\langle r \rangle \langle r \rangle \langle -(k-1) \rangle \cong \tilde{M}^{p_j}(\alpha)\langle 2r+1-k \rangle.$$

The second shift comes from the fact that the module is Koszul, and the third shift comes from the definition of the functor. $\square$

Corollary 11. *There is an isomorphism of graded functors:*

$$L \tilde{Z}_{q_j}^{p_j} \tilde{\epsilon}_{p_j}^{q_j} [-(k-1)] \cong \oplus_{r=0}^{k-1} \text{Id}[2r-(k-1)]\langle 2r-(k-1) \rangle.$$

5.4. Graded Diagram Relation 2. In this subsection we include only the i and $(i+1)$ st entries of the weights for the Verma modules.

Lemma 39. *There is an isomorphism, of graded objects*

$$L \tilde{Z}^{\tilde{s}_i} \tilde{\epsilon}_{\tilde{s}_i} \tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}}) \cong \tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}})\langle -1 \rangle \oplus \tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}})[2]\langle 1 \rangle.$$

Proof. Consider the object $\tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}})$. There is an exact sequence

$$0 \rightarrow \tilde{M}(\underbrace{a_{i+1}, a_i})\langle 1 \rangle \rightarrow \tilde{M}(\underbrace{a_i, a_{i+1}}) \rightarrow \tilde{\epsilon}_{\tilde{s}_i} \tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}}) \rightarrow 0.$$

Now apply the functor $L \tilde{Z}^{\tilde{s}_i}$ to get a long exact sequence. It follows that

$$L_0 \tilde{Z}^{\tilde{s}_i} \tilde{\epsilon}_{\tilde{s}_i} \tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}}) \cong \tilde{M}^{\tilde{s}_i}(\underbrace{a_i, a_{i+1}})\langle -1 \rangle.$$

The shift comes from the convention taken for the functor. Thus,

$$L_2 \widetilde{Z}^{\mathfrak{s}_i} \widetilde{\epsilon}_{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}) \cong \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}) \langle -1 \rangle \langle 1 \rangle \langle 1 \rangle.$$

The second shift comes from the exact sequence and the third shift comes from that the module is Koszul.. Therefore

$$L \widetilde{Z}^{\mathfrak{s}_i} \widetilde{\epsilon}_{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}) \cong \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}) \langle -1 \rangle \oplus \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}}) [2] \langle 1 \rangle.$$

□

Corollary 12. *There is an isomorphism of graded functors: $L \widetilde{Z}^{\mathfrak{s}_i} \widetilde{\epsilon}_{\mathfrak{s}_i} [-1] \cong \text{Id}[-1] \langle -1 \rangle \oplus \text{Id}[1] \langle 1 \rangle$.*

5.5. Graded Diagram Relation 3. As in section 4.3, we study the graded generalized Verma module of highest weight α , $\widetilde{M}^{\mathfrak{p}_{i+1}}(a_1, \dots, a_i, k-1, \dots, 0, a_{i+k}, \dots, a_n)$.

Lemma 40. *Let $a_i = l$. Suppose $s = 0$ or $s = 2$. Then*

$$\begin{aligned} L_s \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{\epsilon}_{\mathfrak{p}_{i+1}}^{\mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) &= 0. \\ L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{\epsilon}_{\mathfrak{p}_{i+1}}^{\mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0, a_{i+k}) &\cong L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(e.\alpha) / L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_0 \\ L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_0 &\cong L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1.\alpha) \langle 1 \rangle / L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_1 \\ \dots & \\ L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l-3} &\cong L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l-2}.\alpha) \langle k-l-2 \rangle / L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l-2} \\ L_s \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l-2} &\cong L_{s-1} \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l-1} \\ L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l-1} &\cong L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l}.\alpha) \langle k-l \rangle / L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l} \\ L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l} &\cong L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-l+1}.\alpha) \langle k-l+1 \rangle / L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-l+1} \\ \dots & \\ L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-4} &\cong L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-3}.\alpha) \langle k-3 \rangle / L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-3} \\ L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{K}_{k-3} &\cong L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-2}.\alpha) \langle k-2 \rangle / L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-1}.\alpha) \langle k-1 \rangle. \end{aligned}$$

Proof. This easily follows from the proof of lemma 10. □

Lemma 41. *Suppose $s \neq k-2, k$. Then*

$$L_s \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{p}_{i+1}} \widetilde{\epsilon}_{\mathfrak{s}_i + \mathfrak{q}_{i+1}}^{\mathfrak{q}_{i+1}} (L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-2}.\alpha) \langle k-2 \rangle / L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-1}.\alpha) \langle k-1 \rangle) \cong 0.$$

For $s = k-2$, it is isomorphic to $\widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \langle k-4 \rangle$. For $s = k$, it is isomorphic to $\widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \langle k-2 \rangle$.

Proof. Note that if $s = k-2$,

$$\begin{aligned} L_s \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{p}_{i+1}} \widetilde{\epsilon}_{\mathfrak{s}_i + \mathfrak{q}_{i+1}}^{\mathfrak{q}_{i+1}} L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-2}.\alpha) \langle k-2 \rangle &\cong \\ \widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \langle (k-2) \rangle \langle -1 \rangle \langle (k-2) \rangle \langle -(k-1) \rangle. \end{aligned}$$

The second shift comes from the innermost functor. The third shift comes from Koszulness of the module. The fourth shift comes from the outermost functor. For all other s it is zero.

Also if $s = k-1$,

$$\begin{aligned} L_s \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{p}_{i+1}} \widetilde{\epsilon}_{\mathfrak{s}_i + \mathfrak{q}_{i+1}}^{\mathfrak{q}_{i+1}} L_0 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{q}_{i+1}}(\sigma_1 \dots \sigma_{k-1}.\alpha) \langle k-1 \rangle &\cong \\ \widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \langle (k-1) \rangle \langle -1 \rangle \langle (k-1) \rangle \langle -(k-1) \rangle. \end{aligned}$$

For all other s it is zero. Then the lemma follows from the of lemma 11. □

Lemma 42. *If $s = 1, 3, \dots, 2(k-1) - 1$, then*

$$L_s \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{p}_{i+1}} \widetilde{\epsilon}_{\mathfrak{s}_i + \mathfrak{q}_{i+1}}^{\mathfrak{q}_{i+1}} L_1 \widetilde{Z}_{\mathfrak{q}_{i+1}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1}} \widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \cong \widetilde{M}^{\mathfrak{p}_{i+1}}(a_i, k-1, \dots, 0) \langle -k+1+s \rangle.$$

Otherwise it is zero.

Proof. This follows from the previous lemma and following the exact sequences of lemmas 12, 13, 14 and corollaries 5 and 6. $\square$

The graded version of proposition 10 now holds.

Proposition 15. *There is an isomorphism of graded functors*

$$L\tilde{Z}_{q_{i+1}}^{p_{i+1}+1}\tilde{\epsilon}_{s_i+q_{i+1}}^{q_{i+1}}[-1]L\tilde{Z}_{q_{i+1}}^{s_i+q_{i+1}+1}\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)] \cong \oplus_{r=1}^{k-1} \text{Id}[2r-k]\langle 2r-k \rangle.$$

5.6. Graded Diagram Relation 4. We begin with a generalized Verma module

$$\widetilde{M}^\alpha(\underbrace{k-1, \dots, 0}_{\text{width } k}, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k}).$$

Suppose $a_{i+k} = l$. Let us denote the module above by $\widetilde{M}^\alpha(l)$.

There are short exact sequences coming from the generalized BGG resolution:

$$\begin{aligned} 0 \rightarrow \tilde{K}_0 \rightarrow \widetilde{M}^\beta(\underbrace{k-1, \dots, \hat{0}}_{\text{width } k}, 0, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k})\langle 0 \rangle \rightarrow \widetilde{M}^\alpha(l) \rightarrow 0 \\ 0 \rightarrow \tilde{K}_1 \rightarrow \widetilde{M}^\beta(\underbrace{k-1, \dots, \hat{1}}_{\text{width } k}, 1, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k})\langle 1 \rangle \rightarrow \tilde{K}_0 \rightarrow 0 \\ \dots \\ 0 \rightarrow \tilde{K}_{l-1} \rightarrow \widetilde{M}^\beta(\underbrace{k-1, \dots, \widehat{l-1}}_{\text{width } k}, 0, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k})\langle l-1 \rangle \rightarrow \tilde{K}_{l-2} \rightarrow 0 \\ 0 \rightarrow \tilde{K}_l \rightarrow \widetilde{M}^\beta(\underbrace{k-1, \dots, \hat{l}}_{\text{width } k}, l, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k})\langle l \rangle \rightarrow \tilde{K}_{l-1} \rightarrow 0 \\ 0 \rightarrow \tilde{K}_{l+1} \rightarrow \widetilde{M}^\beta(\underbrace{k-1, \dots, \widehat{l+1}}_{\text{width } k}, 0, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k})\langle l+1 \rangle \rightarrow \tilde{K}_l \rightarrow 0 \\ \dots \\ 0 \rightarrow \tilde{K}_{k-1} \rightarrow \widetilde{M}^\beta(\underbrace{\widehat{k-1}, \dots, 0}_{\text{width } k}, k-1, \underbrace{a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k})\langle k-1 \rangle \rightarrow \tilde{K}_{k-2} \rightarrow 0. \end{aligned}$$

As in the ungraded case, the following lemma follows immediately.

Lemma 43. *There are exact sequences:*

$$\begin{aligned} 0 \rightarrow L_1\tilde{Z}_\beta^\gamma\tilde{K}_0 \rightarrow \widetilde{M}^\gamma(\underbrace{k-1, \dots, \hat{0}}_{\text{width } k}, \underbrace{a_{i+k}, 0}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle 0 \rangle \rightarrow L_1\tilde{Z}_\beta^\alpha\tilde{\epsilon}_\alpha^\beta\widetilde{M}^\alpha(l) \rightarrow 0 \\ 0 \rightarrow L_1\tilde{Z}_\beta^\gamma\tilde{K}_1 \rightarrow \widetilde{M}^\gamma(\underbrace{k-1, \dots, \hat{1}}_{\text{width } k}, \underbrace{a_{i+k}, 1}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle 1 \rangle \rightarrow L_1\tilde{Z}_\beta^\gamma\tilde{K}_0 \rightarrow 0 \\ \dots \\ 0 \rightarrow L_1\tilde{Z}_\beta^\gamma\tilde{K}_{l-1} \rightarrow \widetilde{M}^\gamma(\underbrace{k-1, \dots, \widehat{l-1}}_{\text{width } k}, \underbrace{a_{i+k}, l-1}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle l-1 \rangle \rightarrow L_1\tilde{Z}_\beta^\gamma\tilde{K}_{l-2} \rightarrow 0 \\ L_i\tilde{Z}_\beta^\gamma\tilde{K}_l = L_{i+1}\tilde{Z}_\beta^\gamma\tilde{K}_{l-1}, \forall i \\ 0 \rightarrow L_0\tilde{Z}_\beta^\gamma\tilde{K}_{l+1} \rightarrow \widetilde{M}^\gamma(\underbrace{k-1, \dots, \widehat{l+1}}_{\text{width } k}, \underbrace{l+1, a_{i+k}}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle l \rangle \rightarrow L_0\tilde{Z}_\beta^\gamma\tilde{K}_l \rightarrow 0 \\ \dots \\ 0 \rightarrow L_0\tilde{Z}_\beta^\gamma\tilde{K}_{k-1} \rightarrow \widetilde{M}^\gamma(\underbrace{\widehat{k-1}, \dots, 0}_{\text{width } k}, \underbrace{k-1, a_{i+k}}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle k-2 \rangle \rightarrow L_0\tilde{Z}_\beta^\gamma\tilde{K}_{k-2} \rightarrow 0 \end{aligned}$$

Proof. The shifts in the grading come from the definition of the functors and Koszulness of the modules. $\square$

Corollary 13. *There are exact sequences:*

$$\begin{aligned} 0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_1\tilde{Z}_\beta^\gamma\tilde{K}_0 \rightarrow \tilde{\epsilon}_\gamma^\beta\widetilde{M}^\gamma(\underbrace{k-1, \dots, \hat{0}}_{\text{width } k}, \underbrace{a_{i+k}, 0}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle 0 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1\tilde{Z}_\beta^\alpha\tilde{\epsilon}_\alpha^\beta\widetilde{M}^\alpha(l) \rightarrow 0 \\ 0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_1\tilde{Z}_\beta^\gamma\tilde{K}_1 \rightarrow \tilde{\epsilon}_\gamma^\beta\widetilde{M}^\gamma(\underbrace{k-1, \dots, \hat{1}}_{\text{width } k}, \underbrace{a_{i+k}, 1}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle 1 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1\tilde{Z}_\beta^\gamma\tilde{K}_0 \rightarrow 0 \\ \dots \\ 0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_1\tilde{Z}_\beta^\gamma\tilde{K}_{l-1} \rightarrow \tilde{\epsilon}_\gamma^\beta\widetilde{M}^\gamma(\underbrace{k-1, \dots, \widehat{l-1}}_{\text{width } k}, \underbrace{a_{i+k}, l-1}_{\text{width } 1}, \underbrace{a_{i+k+1}, \dots, a_{i+2k-1}}_{\text{width } k-1})\langle l-1 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1\tilde{Z}_\beta^\gamma\tilde{K}_{l-2} \rightarrow 0 \\ \tilde{\epsilon}_\gamma^\beta L_i\tilde{Z}_\beta^\gamma\tilde{K}_l \cong \tilde{\epsilon}_\gamma^\beta L_{i+1}\tilde{Z}_\beta^\gamma\tilde{K}_{l-1}, \forall i \end{aligned}$$

$$\begin{aligned}
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{K}_{l+1} &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{k-1, \dots, \widehat{l+1}, 0}_{\dots}, \underbrace{l+1, a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\dots}) \langle l \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{K}_l \rightarrow 0 \\
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{K}_{k-1} &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{k-1, \dots, 0}_{\dots}, \underbrace{k-1, a_{i+k}, a_{i+k+1}, \dots, a_{i+2k-1}}_{\dots}) \langle k-2 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{K}_{k-2} \rightarrow 0
\end{aligned}$$

Lemma 44. Let $\{a_{i+k+1}, \dots, a_{i+2k-1}\} = \{k-1, \dots, \hat{m}, \dots, 0\}$. Let $m \neq l$. If $s = k-1$, then

$$L_s \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\alpha^\beta \tilde{M}^\alpha(l) = \tilde{M}^\delta(\underbrace{k-1, \dots, \hat{m}, \dots, 0}_{\dots}, \underbrace{a_{i+k}, k-1, \dots, 0}_{\dots}) \langle 0 \rangle.$$

Otherwise it is zero.

Proof. This follows easily from the previous corollary and the Koszul property of the modules. $\square$

Now consider $\tilde{M}^\delta(\underbrace{a_i, \dots, a_{i+k-2}, l}_{\dots}, \underbrace{k-1, \dots, 0}_{\dots}) = \tilde{M}^\delta(l)$. There is a graded resolution for this module as there was for $\tilde{M}^\alpha(l)$. Similarly we get the following lemma.

Lemma 45. There are exact sequences:

$$\begin{aligned}
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_0 &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{a_i, \dots, a_{i+k-2}, k-1}_{\dots}, \underbrace{a_{i+k-1}, \widehat{k-1}, \dots, 0}_{\dots}) \langle 0 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{M}^\delta(l) \rightarrow 0 \\
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_1 &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{a_i, \dots, a_{i+k-2}, k-2}_{\dots}, \underbrace{a_{i+k-1}, k-1, \widehat{k-2}, \dots, 0}_{\dots}) \langle 1 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_0 \rightarrow 0 \\
&\dots \\
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-2} &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{a_i, \dots, a_{i+k-2}, l+1}_{\dots}, \underbrace{a_{i+k-1}, k-1, \widehat{l+1}, \dots, 0}_{\dots}) \langle k-l-2 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-3} \rightarrow 0 \\
&\tilde{\epsilon}_\gamma^\beta L_i \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-1} \cong \tilde{\epsilon}_\gamma^\beta L_{i+1} \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-2}, \forall i \\
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l} &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{a_i, \dots, a_{i+k-2}, a_{i+k-1}, l-1}_{\dots}, \underbrace{k-1, \widehat{l-1}, \dots, 0}_{\dots}) \langle k-l-1 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-1} \rightarrow 0 \\
&\dots \\
0 \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{J}_{k-1} &\rightarrow \tilde{\epsilon}_\gamma^\beta \tilde{M}^\gamma(\underbrace{a_i, \dots, a_{i+k-2}, a_{i+k-1}, 0}_{\dots}, \underbrace{k-1, \dots, \hat{0}}_{\dots}) \langle k-2 \rangle \rightarrow \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{J}_{k-2} \rightarrow 0
\end{aligned}$$

Lemma 46. Let $\{a_i, \dots, a_{i+k-2}\} = \{k-1, \dots, \hat{m}, \dots, 0\}$. Let $m \neq l$. If $s = k-1$, then

$$L_s \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\delta^\beta \tilde{M}^\delta(l) = \tilde{M}^\alpha(\underbrace{k-1, \dots, 0}_{\dots}, \underbrace{a_{i+k-1}, k-1, \dots, \hat{m}, \dots, 0}_{\dots}) \langle 0 \rangle.$$

Otherwise it is zero.

Proof. This follows from the previous lemma. $\square$

We must now study the case when $l = m$.

Lemma 47. Let $m = l$. Then $L_s \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\alpha^\beta \tilde{M}^\alpha(l) \cong$

$$\begin{aligned}
&\tilde{M}^\delta(0) \langle -l+1 \rangle / \dots / \tilde{M}^\delta(k-1) \langle k-l \rangle & \text{if } s = k-l \\
&L_{k-l+1} \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} & \text{if } s = k-1
\end{aligned}$$

and there is an exact sequence

$$0 \rightarrow \tilde{M}^\delta(l) \langle 1 \rangle / \dots / \tilde{M}^\delta(k-1) \langle k-l \rangle \rightarrow L_{k-l+1} \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} \rightarrow \tilde{M}^\delta(l+1) \langle 0 \rangle / \dots / \tilde{M}^\delta(k-1) \langle k-l-2 \rangle \rightarrow 0.$$

Proof. By corollary 13, we get $L_s \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{K}_{k-2} \cong$

$$\begin{aligned}
&\tilde{M}^\delta(k-1) \langle k-2-l \rangle & \text{if } s = k-l-1 \\
&0 & \text{if } s \neq k-l-1.
\end{aligned}$$

Continuing in this way and using corollary 13 we get $L_s \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{K}_l \cong$

$$\begin{aligned}
&\tilde{M}^\delta(l+1) \langle 0 \rangle / \dots / \tilde{M}^\delta(k-1) \langle k-2-l \rangle & \text{if } s = k-l-1 \\
&0 & \text{if } s \neq k-l-1
\end{aligned}$$

and then $L_s \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-1} \cong$

$$\begin{aligned} & \tilde{M}^\delta(l+1)\langle 0 \rangle / \dots / \tilde{M}^\delta(k-1)\langle k-l-2 \rangle & \text{if } s = k-l-1 \\ & 0 & \text{if } s \neq k-l-1. \end{aligned}$$

Corollary 13 then gives an exact sequence

$$0 \rightarrow \tilde{M}^\delta(l-1)\langle 0 \rangle \rightarrow L_{k-l} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-2} \rightarrow \tilde{M}^\delta(l+1)\langle 0 \rangle / \dots / \tilde{M}^\delta(k-1)\langle k-l-2 \rangle \rightarrow 0.$$

Next, corollary 13 gives the following diagram:

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ & & \tilde{M}^\delta(l-1)\langle 0 \rangle & & & & \\ & & \downarrow & & & & \\ L_{k-l+1} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} & \xrightarrow{\quad} & L_{k-l} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-2} & \xrightarrow{\quad} & \tilde{M}^\delta(l-2)\langle -1 \rangle & \twoheadrightarrow & L_{k-l} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} \\ & & \downarrow & & & & \\ & & \tilde{M}^\delta(l+1)\langle 0 \rangle / \dots / \tilde{M}^\delta(k-1)\langle k-l-2 \rangle & & & & \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

Thus $L_{k-l} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} \cong \tilde{M}^\delta(l-2)\langle -1 \rangle / \dots / \tilde{M}^\delta(k-1)\langle k-l \rangle$ and there is an exact sequence

$$0 \rightarrow \tilde{M}^\delta(l)\langle 1 \rangle / \dots / \tilde{M}^\delta(k-1)\langle k-l \rangle \rightarrow L_{k-l+1} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} \rightarrow \tilde{M}^\delta(l+1)\langle 0 \rangle / \dots / \tilde{M}^\delta(k-1)\langle k-l-2 \rangle \rightarrow 0.$$

Continuing in this manner and using the corollary gives us the lemma. $\square$

Lemma 48. *Let $m = l$. Then $L_s \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\delta^\beta \tilde{M}^\delta(l) \cong$*

$$\begin{aligned} & \tilde{M}^\alpha(k-1)\langle l-k+2 \rangle / \dots / \tilde{M}^\alpha(0)\langle l+1 \rangle & \text{if } s = l+1 \\ & L_{l+2} \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-4} & \text{if } s = k-1 \end{aligned}$$

and there is an exact sequence

$$0 \rightarrow \tilde{M}^\alpha(l)\langle 1 \rangle / \dots / \tilde{M}^\alpha(0)\langle l+1 \rangle \rightarrow L_{l+2} \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-4} \rightarrow \tilde{M}^\alpha(l-1)\langle 0 \rangle / \dots / \tilde{M}^\alpha(0)\langle l-1 \rangle \rightarrow 0.$$

Proof. We use the exact sequences of lemma 45.

If $s = l$, $L_s \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{J}_{k-2} \cong \tilde{M}^\alpha(0)\langle l-1 \rangle$ and is zero otherwise.

If $s = l$, $L_s \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{J}_{k-3} \cong \tilde{M}^\alpha(1)\langle l-2 \rangle / \tilde{M}^\alpha(0)\langle l-1 \rangle$ and is zero otherwise.

Continuing in this way,

If $s = l$, $L_s \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_0 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-1} \cong \tilde{M}^\alpha(l-1)\langle 0 \rangle / \dots / \tilde{M}^\alpha(0)\langle l-1 \rangle$ and is zero otherwise.

Next we get an exact sequence

$$0 \rightarrow \tilde{M}^\alpha(l+1)\langle 0 \rangle \rightarrow L_{l+1} \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{J}_{k-l-3} \rightarrow \tilde{M}^\alpha(l-1)\langle 0 \rangle / \dots / \tilde{M}^\alpha(0)\langle l-1 \rangle \rightarrow 0.$$

Lemma 45 then produces the following diagram:

$$\begin{array}{ccccccc}
& & 0 & & & & \\
& & \downarrow & & & & \\
& & \widetilde{M}^\alpha(l+1)\langle 0 \rangle & & & & \\
& & \downarrow & & & & \\
L_{l+2}\widetilde{Z}_\beta^\alpha\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{J}_{k-l-4} & \xrightarrow{\quad} & L_{l+1}\widetilde{Z}_\beta^\alpha\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{J}_{k-l-3} & \xrightarrow{\quad} & \widetilde{M}^\alpha(l+2)\langle -1 \rangle & \xrightarrow{\quad} & L_{l+1}\widetilde{Z}_\beta^\alpha\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{J}_{k-l-4} \xrightarrow{\quad} 0 \\
& & \downarrow & & & & \\
& & \widetilde{M}^\alpha(l-1)\langle 0 \rangle / \dots / \widetilde{M}^\alpha(0)\langle l-1 \rangle & & & & \\
& & \downarrow & & & & \\
& & 0 & & & &
\end{array}$$

Thus $L_{l+1}\widetilde{Z}_\beta^\alpha\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{J}_{k-l-4} \cong \widetilde{M}^\alpha(l+2)\langle -1 \rangle / \dots / \widetilde{M}^\alpha(0)\langle l+1 \rangle$ and there is an exact sequence

$$0 \rightarrow \widetilde{M}^\alpha(l)\langle 1 \rangle / \dots / \widetilde{M}^\alpha(0)\langle l+1 \rangle \rightarrow L_{l+2}\widetilde{Z}_\beta^\alpha\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{J}_{k-l-4} \rightarrow \widetilde{M}^\alpha(l-1)\langle 0 \rangle / \dots / \widetilde{M}^\alpha(0)\langle l-1 \rangle \rightarrow 0.$$

Using the rest of the exact sequences of lemma 45, we easily obtain this lemma. (The details are the same as lemma 47.) $\square$

$$\text{Let } F = L\widetilde{Z}_\beta^\alpha\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{\epsilon}_\delta^\beta.$$

Lemma 49. For $s = 0, 1, \dots, k-l-2$, $H^{-r}F\widetilde{M}^\delta(l)\langle l \rangle / \dots / \widetilde{M}^\delta(k-1)\langle k-1 \rangle \cong$

$$\begin{aligned}
& (\widetilde{M}^\alpha(k-1)\langle 0 \rangle / \dots / \widetilde{M}^\delta(0)\langle k-1 \rangle) \langle k - (2s+1) - l - 1 \rangle & \text{if } r = 2(k-1) - l - (2s+1) \\
& \widetilde{M}^\alpha(l-1)\langle l \rangle / \dots / \widetilde{M}^\alpha(0)\langle 2l-1 \rangle & \text{if } r = k-1.
\end{aligned}$$

Proof. We will verify this by induction on l . By proposition 15,

$$F\widetilde{M}^\delta(0)\langle 0 \rangle / \dots / \widetilde{M}^\delta(k-1)\langle k-1 \rangle \cong \oplus_{r=0}^{k-2} \widetilde{M}^\alpha(k-1)\langle 0 \rangle / \dots / \widetilde{M}^\alpha(0)\langle k-1 \rangle [2r+1] \langle 2r-k+2 \rangle.$$

This is the base case.

Consider the short exact sequence

$$0 \rightarrow \widetilde{M}^\delta(l)\langle l \rangle / \dots / \widetilde{M}^\delta(k-1)\langle k-1 \rangle \rightarrow \widetilde{M}^\delta(l-1)\langle l-1 \rangle \rightarrow \widetilde{M}^\delta(l-1)\langle l-1 \rangle / \dots / \widetilde{M}^\delta(k-1)\langle k-1 \rangle \rightarrow 0.$$

By the induction hypothesis, for $s = 0, 1, \dots, k-l-1$, $H^{-r}F\widetilde{M}^\delta(l-1)\langle l-1 \rangle / \dots / \widetilde{M}^\delta(k-1)\langle k-1 \rangle \cong$

$$\begin{aligned}
& (\widetilde{M}^\alpha(k-1)\langle 0 \rangle / \dots / \widetilde{M}^\delta(0)\langle k-1 \rangle) \langle k - (2s+1) - l \rangle & \text{if } r = 2(k-1) - l + 1 - (2s+1) \\
& \widetilde{M}^\alpha(l-2)\langle 1 \rangle / \dots / \widetilde{M}^\alpha(0)\langle l-1 \rangle & \text{if } r = k-1.
\end{aligned}$$

By lemma 48, $H^{-s}F\widetilde{M}^\delta(l-1) \cong$

$$\begin{aligned}
& \widetilde{M}^\alpha(k-1)\langle l-k+1 \rangle / \dots / \widetilde{M}^\delta(0)\langle l \rangle & \text{if } s = l \\
& X_{l-1} & \text{if } s = k-1,
\end{aligned}$$

such that X_{l-1} fits into the following diagram:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \widetilde{M}^\alpha(l-1)\langle l \rangle / \dots / \widetilde{M}^\alpha(0)\langle 2l-1 \rangle & \xrightarrow{f} & X_{l-1} & \xrightarrow{g} & \widetilde{M}^\alpha(l-2)\langle l-1 \rangle / \dots / \widetilde{M}^\alpha(0)\langle 2l-3 \rangle \longrightarrow 0 \\
& & & & \downarrow h & & \\
& & & & \widetilde{M}^\alpha(l-2)\langle l-1 \rangle / \dots / \widetilde{M}^\alpha(0)\langle 2l-3 \rangle. & &
\end{array}$$

The fact that the maps g and h are the same follows from the proof of the ungraded case. This finishes the lemma as in the ungraded case as well. $\square$

Lemma 50. $H^{-s}FL\widetilde{Z}_\beta^\delta\widetilde{\epsilon}_\gamma^\beta L_1\widetilde{Z}_\beta^\gamma\widetilde{\epsilon}_\alpha^\beta \widetilde{M}^\alpha(l) \cong$

$$\begin{aligned}
& (\widetilde{M}^\alpha(k-1)\langle 0 \rangle / \dots / \widetilde{M}^\delta(0)\langle k-1 \rangle) \langle s - 2k + l + 1 \rangle & \text{if } s = k-l+1, k-l+3, \dots, k-l+2(k-1)-3 \\
& \widetilde{M}^\alpha(l) & \text{if } s = 2(k-1).
\end{aligned}$$

Proof. By lemma 47, there is a distinguished triangle

$$L_{k-l+1} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} [k-1] \rightarrow L \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\alpha^\beta \tilde{M}^\alpha(l) \rightarrow \tilde{M}^\delta(0) \langle -l+1 \rangle / \dots / \tilde{M}^\delta(k-1) \langle k-l \rangle [k-l].$$

By proposition 15, we know that $H^{-s} F \tilde{M}^\delta(0) \langle 0 \rangle / \dots / \tilde{M}^\delta(k-1) \langle k-1 \rangle [k-l] \cong (\tilde{M}^\alpha(k-1) \langle 0 \rangle / \dots / \tilde{M}^\alpha(0) \langle k-1 \rangle) \langle s-2k+l+1 \rangle$ for $s = k-l+1, k-l+3, \dots, k-l+2(k-1)-1$.

We easily compute that for $s = 2(k-1)$,

$$H^{-s} F L_{k-l+1} \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{K}_{l-3} [k-1] \cong \tilde{M}^\alpha(l) \langle 0 \rangle.$$

Therefore the long exact sequence for the distinguished triangle and F gives $H^{-s} F L \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta L_1 \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\alpha^\beta \tilde{M}^\alpha(l) \cong$

$$\begin{aligned} & (\tilde{M}^\alpha(k-1) \langle 0 \rangle / \dots / \tilde{M}^\delta(0) \langle k-1 \rangle) \langle s-2k+2 \rangle & \text{if } s = k-l+1, k-l+3, \dots, k-l+2(k-1)-3 \\ & \tilde{M}^\alpha(l) & \text{if } s = 2(k-1). \end{aligned}$$

□

This computation along with proposition 11 gives the following corollary.

Corollary 14. *There is an isomorphism of graded functors:*

$$L \tilde{Z}_\beta^\alpha \tilde{\epsilon}_\gamma^\beta [-1] L \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\delta^\beta [-(k-1)] L \tilde{Z}_\beta^\delta \tilde{\epsilon}_\gamma^\beta [-1] L \tilde{Z}_\beta^\gamma \tilde{\epsilon}_\alpha^\beta [-(k-1)] \cong \text{Id} \oplus \bigoplus_{r=1}^{k-2} \tilde{\epsilon}_{\alpha+\delta}^\alpha [-(k-1)] L \tilde{Z}_\alpha^{\alpha+\delta} [2r-(k-1)] \langle 2r-(k-1) \rangle.$$

5.7. Graded Diagram Relation 5. We need only compute in the graded case on a convenient generalized Verma module. We choose $\tilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}})$ with $a_i > a_{i+2} > a_{i+1}$. Now we just repeat most of the arguments of lemma 25.

Proposition 16. *There is an isomorphism of graded functors*

$$L \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} [-1] L \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i} [-1] \cong \text{Id} \oplus \tilde{\epsilon}_{\mathfrak{t}_i}^{\mathfrak{s}_i} [-2] L \tilde{Z}^{\mathfrak{t}_i}.$$

Proof. Consider the short exact sequence

$$0 \rightarrow \tilde{M}(a_{i+1}, a_i, a_{i+2}) \langle 1 \rangle \rightarrow \tilde{M}(a_i, a_{i+1}, a_{i+2}) \rightarrow \tilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow 0.$$

This gives rise to the short exact sequence

$$(4) \quad 0 \rightarrow \tilde{M}^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}) \rightarrow L_1 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i} \tilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow \tilde{M}^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}) \rightarrow 0.$$

There are two shifts in the first module of this sequence. One comes from the Koszul property of the module. The other is due to the definition of the functor, but they cancel each other out.

Now consider the exact sequence

$$0 \rightarrow \tilde{M}(a_{i+1}, a_{i+2}, a_i) \langle 1 \rangle \rightarrow \tilde{M}(a_{i+1}, a_i, a_{i+2}) \rightarrow \tilde{M}^{\mathfrak{s}_{i+1}}(a_{i+1}, \underbrace{a_i, a_{i+2}}) \rightarrow 0.$$

Applying the functor $L \tilde{Z}^{\mathfrak{s}_i}$ gives

$$\begin{aligned} L_1 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \tilde{M}^{\mathfrak{s}_{i+1}}(a_{i+1}, a_i, a_{i+2}) &\cong \tilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) / \tilde{M}^{\mathfrak{s}_i}(\underbrace{a_{i+1}, a_{i+2}, a_i} \rangle 1) \\ L_0 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \tilde{M}^{\mathfrak{s}_{i+1}}(a_{i+1}, a_i, a_{i+2}) &\cong L_2 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \tilde{M}^{\mathfrak{s}_{i+1}}(a_{i+1}, a_i, a_{i+2}) \cong 0. \end{aligned}$$

Next we look at

$$0 \rightarrow \tilde{M}(a_i, a_{i+1}, a_{i+2}) \langle 1 \rangle \rightarrow \tilde{M}(a_i, a_{i+2}, a_{i+1}) \rightarrow \tilde{M}^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}) \rightarrow 0.$$

This leads to the exact sequence

$$\begin{aligned} 0 \rightarrow L_1 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \tilde{M}^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}) &\rightarrow \tilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \rightarrow \tilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}) \langle -1 \rangle \rightarrow \\ L_0 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \tilde{M}^{\mathfrak{s}_{i+1}}(a_i, \underbrace{a_{i+2}, a_{i+1}}) &\rightarrow 0. \end{aligned}$$

As in the ungraded case, the middle map is the standard map so

$$\begin{aligned} L_0 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \widetilde{M}^{\mathfrak{s}_{i+1}}(a_i, a_{i+2}, a_{i+1}) &\cong \tilde{\epsilon}_{\mathfrak{t}_i}^{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{\langle -1 \rangle}) \\ L_1 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} \widetilde{M}^{\mathfrak{s}_{i+1}}(a_i, a_{i+2}, a_{i+1}) &\cong \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{\langle 1 \rangle}). \end{aligned}$$

There is a long exact sequence coming from equation 4. Substituting in the above into this exact sequence, we get

$$\begin{aligned} 0 \rightarrow \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_{i+2}, a_{i+1}, a_i}_{\langle 1 \rangle}) &\rightarrow L_1 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} L_1 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -1 \rangle}) \rightarrow \\ \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -1 \rangle}) / \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_{i+1}, a_{i+2}, a_i}_{\langle 1 \rangle}) &\rightarrow \tilde{\epsilon}_{\mathfrak{t}_i}^{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{\langle -1 \rangle}) \rightarrow \\ L_0 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} L_1 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -1 \rangle}) &\rightarrow 0. \end{aligned}$$

Thus as in the ungraded case,

$$L_0 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} L_1 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -1 \rangle}) \cong \tilde{\epsilon}_{\mathfrak{t}_i}^{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{\langle -1 \rangle})$$

and

$$L_1 \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}} L_1 \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -1 \rangle}) \cong \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -1 \rangle}).$$

Now,

$$\tilde{\epsilon}_{\mathfrak{t}_i}^{\mathfrak{s}_i}[-2] L \tilde{Z}_{\mathfrak{s}_i}^{\mathfrak{t}_i} \widetilde{M}^{\mathfrak{s}_i}(\underbrace{a_i, a_{i+1}, a_{i+2}}_{\langle -2 \rangle}) \cong \widetilde{M}^{\mathfrak{t}_i}(\underbrace{a_i, a_{i+2}, a_{i+1}}_{\langle -2 \rangle}) \langle 1 \rangle.$$

The first shift comes from the definition of the functor. The second shift comes from the fact that module is Koszul. Thus

$$L \tilde{Z}^{\mathfrak{s}_i} \tilde{\epsilon}_{\mathfrak{s}_{i+1}}[-1] L \tilde{Z}^{\mathfrak{s}_{i+1}} \tilde{\epsilon}_{\mathfrak{s}_i}[-1] \cong \text{Id} \oplus \tilde{\epsilon}_{\mathfrak{t}_i}^{\mathfrak{s}_i}[-2] L \tilde{Z}_{\mathfrak{s}_i}^{\mathfrak{t}_i}.$$

□

6. CROSSINGS

The category $\mathcal{T}$ of oriented tangles has finite sequences of $+$, $-$ signs for objects and isotopy classes of oriented tangles as morphisms. This is a strict tensor category. Oriented caps and cups are clearly morphisms in this category. All eight different types of oriented crossings are morphisms as well. It is shown in [Ka], theorem XII.2.2 that the morphisms are generated by oriented cups and caps and only two oriented crossings. That theorem also gives a complete list of the relations that are satisfied, (along with planar isotopies for tangles without crossings.) The goal of this section is to assign functors to the two crossings which serve as generators and show that the defining relations are satisfied on a functorial level. That is the main result of this paper.

First we state the functorial relations guaranteeing invariance under isotopy for tangles without crossings. This follows from theorem 6.2 of [Str2] except for the base case in the induction hypothesis of proposition 6.4. This follows from lemma 11.80 of [KV].

Theorem 6. *Let $j \geq i$. Then there are isomorphisms of functors*

- (1) $\cap_{i+1, \pm, r+2} \circ \cup_{i, \mp, r} \cong \text{Id}$
- (2) $\cap_{i, \pm, r+2} \circ \cup_{i+1, \mp, r} \cong \text{Id}$
- (3) $\cap_{j, \pm, r-2} \circ \cap_{i, \pm, r} \cong \cap_{i, \pm, r-2} \circ \cap_{j+2, \pm, r}$
- (4) $\cup_{j, \pm, r-2} \circ \cap_{i, \pm, r} \cong \cap_{i, \pm, r+2} \circ \cup_{j+2, \pm, r}$
- (5) $\cup_{i, \pm, r-2} \circ \cap_{j, \pm, r} \cong \cap_{j+2, \pm, r+2} \circ \cup_{i, \pm, r}$

(6)

$$\cup_{i,\pm,r+2} \circ \cup_{j,\pm,r} \cong \cup_{j+2,\pm,r+2} \circ \cup_{i,\pm,r}$$

(7)

$$\cap_{i,\pm,r+2} \circ \cup_{i,\pm,r} \cong \text{Id} \oplus \text{Id}.$$

Now to the crossing



we assign the functor

$$\Omega_i = \text{Cone}(\tilde{\epsilon}_{s_i}[-1]L\tilde{Z}_{s_i} \rightarrow \text{Id}[1]\langle 1 \rangle)[-k]\langle -k \rangle.$$

To the opposite crossing



we assign the functor

$$\Pi_i = \text{Cone}(\text{Id}\langle -1 \rangle \rightarrow \tilde{\epsilon}_{s_i}L\tilde{Z}_{s_i})[k-2]\langle k \rangle.$$

For ease of notation, we will assume that all arcs not shown in the diagrams are straight arrows oriented up. Now we must check the following eight relations.

Theorem 7. *The functors assigned to the generators of the set of morphisms for the category $\mathcal{T}$ satisfy the following relations:*

(1)

$$\cap_{i+1,+,r+2} \circ \cup_{i,-,r} \cong \text{Id} \cong \cap_{i,-,r+2} \circ \cup_{i+1,+,r}.$$




FIGURE 14. Relation 1

(2)

$$\cap_{i+1,-,r+2} \circ \cup_{i,+,r} \cong \text{Id} \cong \cap_{i,+,r+2} \circ \cup_{i+1,-,r}.$$




FIGURE 15. Relation 2

(3)

$$\begin{aligned} \cap_{i-2,-,r+2} \circ \cap_{i-1,-,r+4} \circ \Omega_{i+2k-4} \circ \cup_{i+1,+,r+2} \circ \cup_{i,+,r} &\cong \\ \cap_{i,+,r+2} \circ \cap_{i+1,+,r+4} \circ \Omega_{i+2k-4} \circ \cup_{i-1,-,r+2} \circ \cup_{i-2,-,r} & \end{aligned}$$

$$\begin{aligned} \cap_{i-2,-,r+2} \circ \cap_{i-1,-,r+4} \circ \Omega_{i+2k-4} \circ \cup_{i+1,+,r+2} \circ \cup_{i,+,r} &\cong \\ \cap_{i,+,r+2} \circ \cap_{i+1,+,r+4} \circ \Omega_{i+2k-4} \circ \cup_{i-1,-,r+2} \circ \cup_{i-2,-,r} & \end{aligned}$$




FIGURE 16. Relation 3

(4)

$$\Omega_i \circ \Pi_i \cong \text{Id} \cong \Pi_i \circ \Omega_i$$




FIGURE 17. Relation 4

(5)

$$\Omega_i \circ \Omega_{i+1} \circ \Omega_i \cong \Omega_{i+1} \circ \Omega_i \circ \Omega_{i+1}$$

(6)

$$\cap_{i+1,+,r+2} \circ \Omega_i \circ \cup_{i+1,+,r} \cong \text{Id} \cong \cap_{i+1,+,r+2} \circ \Pi_{i+k-2} \circ \cup_{i+1,+,r}$$

(7)

$$\begin{aligned} \cap_{i,-,r+2} \circ \Omega_{i+k-1} \circ \cup_{i+2,+,r} \circ \cap_{i+2,+,r+2} \circ \Pi_{i+k-1} \circ \cup_{i,-,r} &\cong \text{Id} \cong \\ \cap_{i,-,r+2} \circ \Pi_{i+k-1} \circ \cup_{i+2,+,r} \circ \cap_{i+2,+,r+2} \circ \Omega_{i+k-1} \circ \cup_{i,-,r} & \end{aligned}$$




FIGURE 18. Relation 5




FIGURE 19. Relation 6




FIGURE 20. Relation 7

(8)

$$\begin{aligned} \cap_{i,+ ,r+2} \circ \Omega_{i+k-3} \circ \cup_{i-2,- ,r} \circ \cap_{i-2,- ,r+2} \circ \Pi_{i+k-3} \circ \cup_{i,+ ,r} &\cong \text{Id} \cong \\ \cap_{i,+ ,r+2} \circ \Omega_{i+k-3} \circ \cup_{i-2,- ,r} \circ \cap_{i-2,- ,r+2} \circ \Pi_{i+k-3} \circ \cup_{i,+ ,r} & \end{aligned}$$

6.1. Relation 1. The goal of this section is to prove that the functors assigned to the tangles in diagram 14 are isomorphic. We interpret the integers a_i and $-a_i$, mod k .

Lemma 51. $\cap_{i+1,-a_i,r+2} \circ \cup_{i,a_i,r} \cong \text{Id}$.

Proof. The left hand side is isomorphic to

$$\nu \circ \tilde{\Delta}_{(a_1, \dots, a_i, k, \dots, a_r)}^{(k, a_1, \dots, a_r)} \circ L\tilde{Z}_{(a_1, \dots, a_{i-1}, a_i, -a_i, a_i, \dots, a_r)}^{(a_1, \dots, a_i, k, \dots, a_r)} \circ \tilde{\epsilon}_{(a_1, \dots, a_{i-1}, k, a_i, \dots, a_r)}^{(a_1, \dots, a_{i-1}, a_i, -a_i, a_i, \dots, a_r)}[-(k-1)] \circ \tilde{\Delta}_{(k, a_1, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, a_i, \dots, a_r)} \circ \zeta.$$

This is easily seen to be isomorphic to

$$\cong \nu \circ \tilde{\Delta}_{(a_1, \dots, a_i, k, \dots, a_r)}^{(k, a_1, \dots, a_r)} \circ \tilde{\Delta}_{(k, a_1, \dots, a_r)}^{(a_1, \dots, a_i, k, a_{i+1}, \dots, a_r)} \circ \zeta \cong \text{Id}.$$




FIGURE 21. Relation 8

□

Lemma 52. $\cap_{i,a_i,r+2} \circ \cup_{i+1,-a_i,r} \cong \text{Id}$.

Proof. The left hand side is isomorphic to

$$\nu \circ \tilde{\Delta}_{(a_1, \dots, a_{i-1}, k, \dots, a_r)}^{(k, a_1, \dots, a_r)} \circ L\tilde{Z}_{(a_1, \dots, a_i, -a_i, a_i, a_{i+1}, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, \dots, a_r)} \circ \tilde{e}_{(a_1, \dots, a_i, k, \dots, a_r)}^{(a_1, \dots, a_i, -a_i, a_i, a_{i+1}, \dots, a_r)} [-(k-1)] \circ \tilde{\Delta}_{(k, a_1, \dots, a_r)}^{(a_1, \dots, a_i, k, \dots, a_r)} \circ \zeta.$$

This is easily seen to be isomorphic to

$$\cong \nu \circ \tilde{\Delta}_{(a_1, \dots, a_{i-1}, k, \dots, a_r)}^{(k, a_1, \dots, a_r)} \circ \tilde{\Delta}_{(k, a_1, \dots, a_r)}^{(a_1, \dots, a_{i-1}, k, \dots, a_r)} \circ \zeta \cong \text{Id}.$$

□

Now we easily get relation 1.

Corollary 15. $\cap_{i+1,+,r+2} \circ \cup_{i,-,r} \cong \text{Id} \cong \cap_{i,-,r+2} \circ \cup_{i+1,+,r}$.

Proof. Take $a_i = -1$ and apply the two lemmas. □

6.2. Relation 2. There is the tangle relation for diagram 39. As in the previous subsection there is an analogous isomorphism of functors.

Corollary 16. $\cap_{i+1,-,r+2} \circ \cup_{i,+,r} \cong \text{Id} \cong \cap_{i,+,r+2} \circ \cup_{i+1,-,r}$.

Proof. Take $a_i = 1$ from the two lemmas in the previous subsection. □

6.3. Relation 3. In this section we define functors for crossings with both arrows pointing down. It will be shown that these functors are consistent in a certain way with the functors defined for the other types of crossings. First we must define functors categorifying intertwiners $\Lambda^{k-1}V_{k-1} \otimes \Lambda^{k-1}V_{k-1} \rightarrow \Lambda^{k-2} \otimes \Lambda^k V_{k-1}$ and $\Lambda^{k-2} \otimes \Lambda^k V_{k-1} \rightarrow \Lambda^{k-1}V_{k-1} \otimes \Lambda^{k-1}V_{k-1}$. Let F_1 and F_2 be the functors for the diagrams on the left and right respectively of figure 22.




FIGURE 22.

Explicitly,

$$F_1 = L\tilde{Z}_{\mathbf{q}_i + \mathbf{p}_{i+k}}^{\mathbf{p}_i + \mathbf{p}_{i+k}} L\tilde{Z}_{\mathbf{r}_{i+2} + \mathbf{p}_{i+k}}^{\mathbf{q}_i + \mathbf{p}_{i+k}}$$

$$F_2 = L\tilde{Z}_{\mathbf{p}_i+\mathbf{q}_i+k}^{\mathbf{p}_i+\mathbf{p}_i+k} L\tilde{Z}_{\mathbf{q}_i+\mathbf{q}_i+k}^{\mathbf{p}_i+\mathbf{q}_i+k} \tilde{\epsilon}_{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} [-(k-1)] L\tilde{Z}_{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}} L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{r}_{i+2}+\mathbf{p}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-1)].$$

If $\phi: R \rightarrow S$ is a natural transformation, then there is a natural transformation $\phi^*: S^* \rightarrow R^*$ of adjoint functors. There is a map constructed by taking the adjoint of an adjunction morphism:

$$\alpha_1: F_1 \rightarrow L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{p}_i+k} \tilde{\epsilon}_{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(2k-3)] L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{r}_{i+2}+\mathbf{p}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-1)][k-2]\langle k-2 \rangle.$$

We may rewrite the above line as

$$\alpha_1: F_1 \rightarrow L\tilde{Z}_{\mathbf{q}_i+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{p}_i+k} L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{q}_i+\mathbf{q}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-2)] \tilde{\epsilon}_{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} [-(k-1)] L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{r}_{i+2}+\mathbf{p}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-1)][k-2]\langle k-2 \rangle = F_3.$$

Since

$$L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{q}_i+\mathbf{q}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} \cong \oplus_{j=0}^{k-2} \text{Id}[2j]\langle -(k-2) + 2j \rangle,$$

there is a projection $\alpha_2: F_3 \rightarrow F_2$.

Lemma 53. *The composite $\alpha_2 \circ \alpha_1: F_1 \rightarrow F_2$ is an isomorphism.*

Proof. The map coming from adjunction is clearly non-zero. Since α_2 is just projection, $\alpha_2 \circ \alpha_1$ is non-zero. An easily calculation shows that F_1 an F_2 applied to a generalized Verma module

$$M^{\mathbf{r}_{i+2}+\mathbf{p}_{i+k}}(a_1, \dots, a_{i-1}, a_i, a_{i+1}, \underbrace{a_{i+2}, \dots, a_{i+k-1}}_{k-1}, \dots, 0, a_{i+2k}, \dots, a_n),$$

gives a generalized Verma module

$$M^{\mathbf{p}_i+\mathbf{p}_{i+k}}(a_1, \dots, a_{i-1}, \underbrace{k-1, \dots, 0}_{k-1}, \underbrace{k-1, \dots, 0}_{k-1}, a_{i+2}, \dots, a_n)$$

or zero. Therefore $\alpha_2 \circ \alpha_1$ is a non-zero endomorphism of a generalized Verma module (when it is non-trivial) so it is an isomorphism. This composition of maps is then an isomorphism when the functors are applied to a projective object by considering a filtration with subquotients of generalized Verma modules. Then it is an isomorphism of functors because a bounded complex is quasi-isomorphic to a complex of projective objects. $\square$

Let G_1 and G_2 be functors for the diagrams on the left and right respectively of figure 23.




FIGURE 23.

Explicitly,

$$\begin{aligned} G_1 &= L\tilde{Z}_{\mathbf{p}_i+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{p}_{i+k}} L\tilde{Z}_{\mathbf{q}_i+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} [-(k-1)] L\tilde{Z}_{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{s}_i+\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}} [-1] \\ G_2 &= L\tilde{Z}_{\mathbf{p}_i+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{p}_{i+k}} L\tilde{Z}_{\mathbf{s}_i+\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{s}_i+\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-2)]. \end{aligned}$$

There is a morphism constructed similarly to α_1 above:

$$\begin{aligned} \beta_1: G_2 &\rightarrow L\tilde{Z}_{\mathbf{q}_i+\mathbf{q}_{i+k}}^{\mathbf{p}_i+\mathbf{p}_{i+k}} L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{q}_i+\mathbf{q}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-2)] \tilde{\epsilon}_{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{q}_i+\mathbf{q}_{i+k}} [-(k-1)] \circ \\ &\quad L\tilde{Z}_{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{p}_{i+1}+\mathbf{q}_{i+k}} L\tilde{Z}_{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}}^{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}} \tilde{\epsilon}_{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{r}_{i+2}+\mathbf{q}_{i+k}} [-(k-2)] \tilde{\epsilon}_{\mathbf{s}_i+\mathbf{q}_{i+1}+\mathbf{q}_{i+k}}^{\mathbf{q}_{i+1}+\mathbf{q}_{i+k}} [-1][2k-4]\langle -2k+4 \rangle. \end{aligned}$$

Then there is an obvious projection β_2 from the latter functor to G_1 .

Lemma 54. *The composite $\beta_2 \circ \beta_1: G_2 \rightarrow G_1$ is an isomorphism.*

Proof. The map β_1 is non-zero for an object when the functors applied to the object do not vanish because it comes from adjunction. Since β_2 is projection, it too is non-zero when the functors are non-zero. An easy calculation shows that G_1 and G_2 applied to the generalized Verma module

$$M^{\mathfrak{s}_i \mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}(a_1, \dots, a_i, \underbrace{a_{i+1}, a_{i+2}, \dots, a_{i+k}}_{\text{shifted}}, \underbrace{a_{i+k-1}, \dots, a_{i+2k-1}}_{\text{shifted}}, a_{i+2k}, \dots, a_n)$$

is isomorphic to a shifted generalized Verma module

$$M^{\mathfrak{p}_i + \mathfrak{p}_{i+k}}(a_1, \dots, a_{i-1}, \underbrace{k-1, \dots, 0}_{\text{shifted}}, \underbrace{k-1, \dots, 0}_{\text{shifted}}, a_{i+2k}, \dots, a_n)$$

or zero. Therefore for generalized Verma modules $\beta_2 \circ \beta_1$ is an isomorphism. It is then an isomorphism for projective objects by considering a Verma flag. It is then an isomorphism for all bounded complex because there exists a complex of projective quasi-isomorphic to it. $\square$




FIGURE 24.

Corollary 17. *There is an isomorphism of functors for the two diagrams in figure 24.*

$$G_1 \circ L\tilde{Z}_{\mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}} \cong F_2 \circ L\tilde{Z}_{\mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}^{\mathfrak{r}_{i+2} + \mathfrak{p}_{i+k}} \tilde{\epsilon}_{\mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}^{\mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}[-(k-2)].$$

Proof. By lemma 54, the left hand side is isomorphic to

$$L\tilde{Z}_{\mathfrak{p}_i + \mathfrak{q}_{i+k}}^{\mathfrak{p}_i + \mathfrak{p}_{i+k}} L\tilde{Z}_{\mathfrak{s}_i + \mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}^{\mathfrak{p}_i + \mathfrak{q}_{i+k}} \tilde{\epsilon}_{\mathfrak{s}_i + \mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}^{\mathfrak{s}_i + \mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}[-(k-2)] L\tilde{Z}_{\mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}^{\mathfrak{s}_i + \mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}.$$

Commuting the last two functors in the expression above, we get that it is isomorphic to

$$L\tilde{Z}_{\mathfrak{p}_i + \mathfrak{q}_{i+k}}^{\mathfrak{p}_i + \mathfrak{p}_{i+k}} L\tilde{Z}_{\mathfrak{s}_i + \mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}^{\mathfrak{p}_i + \mathfrak{q}_{i+k}} L\tilde{Z}_{\mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}^{\mathfrak{s}_i + \mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}} \tilde{\epsilon}_{\mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}^{\mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}[-(k-2)].$$

Lemma 53 implies that this is isomorphic to the right hand side of the desired isomorphism. $\square$

The next corollary gives an isomorphism between the two diagrams in figure 25.




FIGURE 25.

Corollary 18. *Let G_1^* and F_2^* denote the adjoint functors. Then*

$$\tilde{\epsilon}_{\mathfrak{s}_i + \mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}^{\mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}}[-1] \circ G_1^* \cong L\tilde{Z}_{\mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}^{\mathfrak{q}_{i+1} + \mathfrak{q}_{i+k}} \circ \tilde{\epsilon}_{\mathfrak{r}_{i+2} + \mathfrak{p}_{i+k}}^{\mathfrak{r}_{i+2} + \mathfrak{q}_{i+k}}[-(k-1)] \circ F_2^*.$$

Proof. This follows from the previous corollary by taking adjoints. $\square$

Let H_1 be the functor for the diagram on the left and H_2 be the functor for the diagram on the right of figure 26.




FIGURE 26.

Let H_3 be the functor for the diagram on the left and H_4 be the functor for the diagram on the right of figure 27.




FIGURE 27.

Let $G = L\tilde{Z}_{\mathbf{r}_i+\mathbf{q}_i+k-2}^{\mathbf{q}_{i-1}+\mathbf{q}_i+k-2} \epsilon_{\mathbf{r}_i+\mathbf{p}_i+k-1}^{\mathbf{r}_i+\mathbf{q}_i+k-2} [-(k-1)] L\tilde{Z}_{\mathbf{r}_i+\mathbf{q}_i+k-2}^{\mathbf{r}_i+\mathbf{p}_i+k-1} \epsilon_{\mathbf{q}_{i-1}+\mathbf{q}_i+k-2}^{\mathbf{r}_i+\mathbf{q}_i+k-2} [-(k-2)]$. It is the functor for the diagram in figure 28.




FIGURE 28.

Corollary 19. $H_1 \cong H_2 \cong G$.

Proof. The two corollaries, imply that $H_1 \cong H_3$ and $H_2 \cong H_4$. Then, using the first relation, both of these functors are seen to be isomorphic to G . $\square$

Now we are prepared to prove the main relations of this section. Let A and B denote the functors assigned to the diagrams on the left and right respectively of figure 29.

Let C and D denote the functors assigned to the diagrams on the left and right respectively of figure 30.

Proposition 17. (1) *The functors A and B are isomorphic.*
 (2) *The functors C and D are isomorphic.*




FIGURE 29.




FIGURE 30.

Proof. The first pair of functors are cones of morphisms $G \rightarrow \text{Id}[1]\langle 1 \rangle$. The second pair are cones of morphisms $\text{Id}[-1]\langle -1 \rangle \rightarrow G$. Clearly,

$$\text{Hom}(G, \text{Id}[1]\langle 1 \rangle) \cong \text{Hom}(\text{Id}[-1]\langle -1 \rangle, G) \cong \text{End}(L\tilde{Z}_{(k-2,1,k-1)}^{(k-2,k)} \tilde{\epsilon}_{(k-1,k-1)}^{(k-2,1,k-1)}).$$

By theorem 34 of [MOS], this space is isomorphic to a space of endomorphisms of a projective functor:

$$\text{End}(\tilde{\theta}_{(k-2,1,k-1)}^{(k-2,k)} \tilde{\theta}_{(k-1,k-1)}^{(k-2,1,k-1)}).$$

By proposition 8.7 of [Str2], this is isomorphic to

$$\text{End}_{C^{(k-2,k)} - \text{gmod} - C^{(k-1,k-1)}}(C^{(k-2,1,k-1)} \otimes_{C^{(k-1,k-1)}} C^{(k-1,k-1)}).$$

As a graded $(C^{(k-2,k)}, C^{(k-1,k-1)})$ -bimodule, $C^{(k-2,1,k-1)} \otimes_{C^{(k-1,k-1)}} C^{(k-1,k-1)}$ is generated by $1 \otimes 1$. Thus the degree zero component of this space of endomorphisms is one dimensional. Since there is a unique homogenous morphism up to scalar of degree 0 between the functors, the cones must be isomorphic. $\square$

6.4. Relation 4. We will show that the functors assigned to the crossings, Π_i and Ω_i are auto-equivalences of the derived category.

Let $\alpha: \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} \rightarrow \text{Id}[2]\langle 1 \rangle$ and $\beta: \text{Id} \rightarrow \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} \langle 1 \rangle$.

Proposition 18. $\Pi_i \circ \Omega_i \cong \text{Id}$.

Proof. We would like to show that $\text{Cone}(\beta)\text{Cone}(\alpha) \cong \text{Id}[r]\langle s \rangle$ for some r and s . Consider the following commutative diagram.

$$\begin{array}{ccccc} \text{Cone}(\beta)\tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} & \xrightarrow{g} & \text{Cone}(\beta)\text{Id}[2]\langle 1 \rangle & \longrightarrow & \text{Cone}(\beta)\text{Cone}(\alpha) \\ \uparrow f & & \uparrow & & \uparrow \\ \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} \langle 1 \rangle \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} & \longrightarrow & \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} \langle 1 \rangle \text{Id}[2]\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} \langle 1 \rangle \text{Cone}(\alpha) \\ \uparrow h & & \uparrow & & \uparrow \\ \text{Id} \circ \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} & \longrightarrow & \text{Id} \circ \text{Id}[2]\langle 1 \rangle & \longrightarrow & \text{Id} \circ \text{Cone}(\alpha) \end{array}$$

By properties of adjoint functors, h is split inclusion so f is surjection onto $\tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} [2]\langle 2 \rangle$. By commutativity, g is the map fitting into the distinguished triangle

$$\text{Id}[2]\langle 1 \rangle \xrightarrow{\beta[2]\langle 1 \rangle} \tilde{\epsilon}_{s_i} L\tilde{Z}_{s_i} [2]\langle 2 \rangle \xrightarrow{g} \text{Cone}(\beta)[2]\langle 1 \rangle.$$

Therefore $\text{Cone}(\beta)\text{Cone}(\alpha) \cong \text{Id}[3]\langle 1 \rangle$. Then with the shifts built into the definition of Π_i and Ω_i , we have $\Pi_i \circ \Omega_i \cong \text{Id}$. $\square$

Proposition 19. $\Omega_i \circ \Pi \cong \text{Id}$.

Proof. As in the previous proposition we would like to show that $\text{Cone}(\alpha)\text{Cone}(\beta) \cong \text{Id}[r]\langle s \rangle$ for some r and s . Consider the following commutative diagram.

$$\begin{array}{ccccc}
\text{Cone}(\alpha) & \longrightarrow & \text{Cone}(\alpha)\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha)\text{Cone}(\beta) \\
\uparrow & & \uparrow & & \uparrow \\
\text{Id}[2]\langle 1 \rangle & \longrightarrow & \text{Id}[2]\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Id}[2]\langle 1 \rangle\text{Cone}(\beta) \\
\uparrow & & \uparrow & & \uparrow h \\
\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i} & \xrightarrow{f} & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\text{Cone}(\beta)
\end{array}$$

Note that $f = \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\text{adj}$ and $g: \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle \rightarrow \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}$ is given by $\text{Id}_{|\tilde{\epsilon}_{s_i}}\text{adj}'\text{Id}_{|L\tilde{Z}^{s_i}\langle 1 \rangle}$, where $\text{adj}: \text{Id} \rightarrow \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}$ is the obvious adjunction map and $\text{adj}': L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_i} \rightarrow \text{Id}$ is an adjunction map as well. By properties of adjunction maps, $g \circ f = \text{Id}$. Thus f is split injective so $\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\text{Cone}(\beta) \cong \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}[2]\langle 2 \rangle$. As in the previous proposition, the map h fits into the distinguished triangle

$$\text{Id}[2]\langle 1 \rangle \xrightarrow{\beta[2]\langle 1 \rangle} \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}[2]\langle 2 \rangle \xrightarrow{h} \text{Cone}(\beta)[2]\langle 1 \rangle.$$

Therefore $\text{Cone}(\alpha)\text{Cone}(\beta) \cong \text{Id}$ and we have shown that Π_i and Ω_i are inverse auto-equivalences. $\square$

6.5. Relation 5.

Lemma 55. Suppose we have the following morphism of distinguished triangles where f and h are split inclusion maps.

$$\begin{array}{ccccc}
X & \xrightarrow{\alpha} & Y & \xrightarrow{\beta} & Z \\
\downarrow f & & \downarrow h & & \downarrow j \\
X' & \xrightarrow{\alpha'} & Y' & \xrightarrow{\beta'} & Z'
\end{array}$$

Then j is also a split inclusion.

Proof. See exercise 7.2 of [Sch]. $\square$

Proposition 20. There is a functorial braid relation $\Pi_i \circ \Pi_{i+1} \circ \Pi_i \cong \Pi_{i+1} \circ \Pi_i \circ \Pi_{i+1}$.

Proof. First we will consider the commutative diagram expressing the functor $\Pi_i \circ \Pi_{i+1}$.

$$\begin{array}{ccccc}
\text{Cone}(\alpha_{i+1}) & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\text{Cone}(\alpha_{i+1}) & \longrightarrow & \text{Cone}(\alpha_i)\text{Cone}(\alpha_{i+1}) \\
\uparrow & & \uparrow & & \uparrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle \\
\uparrow \alpha_{i+1} & & \uparrow & & \uparrow \\
\text{Id} & \xrightarrow{\alpha_i} & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha_i)
\end{array}$$

Apply $\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}$ to $\Pi_i \circ \Pi_{i+1}$ and to the above diagram to get the following diagram.

$$\begin{array}{ccccc}
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_{i+1}) & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\text{Cone}(\alpha_{i+1}) & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_i)\text{Cone}(\alpha_{i+1}) \\
\uparrow & & \uparrow & & \uparrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle \\
\uparrow & & \uparrow & & \uparrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\alpha_{i+1} & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\alpha_i & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_i) \\
\uparrow & & \uparrow & & \uparrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Id} & \xrightarrow{\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\alpha_i} & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_i)
\end{array}$$

There is an obvious morphism from the first diagram to the second one. All of the squares commute. Now apply $\text{Cone}(\alpha_{i+1})$ to the first diagram to get the following commutative diagram.

$$\begin{array}{ccccc}
\text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_{i+1}) & \longrightarrow & \text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle\text{Cone}(\alpha_{i+1}) & \longrightarrow & \text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_i)\text{Cone}(\alpha_{i+1}) \\
\uparrow & & \uparrow & & \uparrow \\
\text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle & \longrightarrow & \text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle & \longrightarrow & \text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle \\
\uparrow & & \uparrow & & \uparrow \\
\text{Cone}(\alpha_{i+1})\alpha_{i+1} & \xrightarrow{\text{Cone}(\alpha_{i+1})\text{Id}} & \text{Cone}(\alpha_{i+1})\alpha_i & \xrightarrow{\text{Cone}(\alpha_{i+1})\alpha_i} & \text{Cone}(\alpha_{i+1})\alpha_i \\
\text{Cone}(\alpha_{i+1})\text{Id} & \longrightarrow & \text{Cone}(\alpha_{i+1})\tilde{\epsilon}_iL\tilde{Z}_i\langle 1\rangle & \longrightarrow & \text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_i)
\end{array}$$

There is a morphism from the second diagram to this one coming from the distinguished triangle

$$\text{Id} \rightarrow \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle \rightarrow \text{Cone}(\alpha_{i+1}).$$

All of the resulting squares commute.

$$\text{Let } F = \text{Cone}((\text{Cone}(\text{Id} \rightarrow \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle)) \rightarrow \text{Cone}(\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle \rightarrow \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle)).$$

Then $\Pi_{i+1} \circ \Pi_i \circ \Pi_{i+1} = \text{Cone}(F \rightarrow \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle F)$. This is isomorphic to

$$\text{Cone}(F \rightarrow \text{Cone}(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle \rightarrow \text{Cone}(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle \rightarrow \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle \oplus \tilde{\epsilon}_{t_i}L\tilde{Z}^{t_i}[1]\langle 3\rangle))).$$

We claim that

$$\text{Cone}(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle \rightarrow \text{Cone}(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle \rightarrow \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle \oplus \tilde{\epsilon}_{t_i}L\tilde{Z}^{t_i}[1]\langle 3\rangle))$$

is isomorphic to $\text{Cone}(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle \rightarrow \tilde{\epsilon}_{t_i}L\tilde{Z}^{t_i}[1]\langle 3\rangle)$.

We have the following commutative diagram

$$\begin{array}{ccccc}
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle & \xrightarrow{\rho} & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle & \xrightarrow{\pi} & \text{Cone}(\rho) \\
\downarrow & & \downarrow f & & \downarrow \nu \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle & \xrightarrow{\mu} & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle & \xrightarrow{g} & \text{Cone}(\mu)
\end{array}$$

By properties of adjunction maps, π maps $\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle$ onto itself and $\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle$ to zero. Thus,

$$\nu\pi(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle) \cong \nu(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle) \cong g \circ f(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle).$$

Let us examine the map f . There is an adjunction map $\text{Id}[2]\langle 1\rangle \rightarrow L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_i}$. There is another map $L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_i} \rightarrow L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}$ which also comes from adjunction. From general principles (see [Mac]), the composition of these two maps is adjunction $\text{Id}[2]\langle 1\rangle \rightarrow L\tilde{Z}^{s_i}\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}$. We know from proposition 12 that this map is a split inclusion. This is how the map f is constructed. This then gives the following commutative diagram of distinguished triangles.

$$\begin{array}{ccccc}
0 & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle & \longrightarrow & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}[2]\langle 3\rangle \\
\downarrow & & \downarrow & & \downarrow \delta \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle & \xrightarrow{\mu} & \tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle & \xrightarrow{g} & \text{Cone}(\mu) \\
\downarrow & & \downarrow & & \downarrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1\rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle & \longrightarrow & \tilde{\epsilon}_{t_i}L\tilde{Z}^{t_i}[1]\langle 3\rangle & \longrightarrow & \text{Cone}(\delta)
\end{array}$$

This implies the claim.

The functor $\Pi_i \circ \Pi_{i+1} \circ \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1\rangle$ is expressed by the following commutative diagram. There is a morphism from the first diagram to this one such that all the resulting squares commute.

$$\begin{array}{ccccc}
\text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha_i)\text{Cone}(\alpha_{i+1})\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle \\
\uparrow & & \uparrow & & \uparrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha_i)\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle \\
\uparrow & & \uparrow & & \uparrow \\
\alpha_{i+1}\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \xrightarrow{\alpha_i\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle} & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha_i)\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle \\
\uparrow & & \uparrow & & \uparrow \\
\text{Id } \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \xrightarrow{\alpha_i\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle} & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle & \longrightarrow & \text{Cone}(\alpha_i)\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle
\end{array}$$

The functor $\Pi_i \circ \Pi_{i+1} \circ \Pi_i$ is expressed through the following diagram.

$$\begin{array}{ccccc}
\text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_i) & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_i) & \longrightarrow & \text{Cone}(\alpha_i)\text{Cone}(\alpha_{i+1})\text{Cone}(\alpha_i) \\
\uparrow & & \uparrow & & \uparrow \\
\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_i) & \longrightarrow & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_i) & \longrightarrow & \text{Cone}(\alpha_i)\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\text{Cone}(\alpha_i) \\
\uparrow & & \uparrow & & \uparrow \\
\alpha_{i+1}\text{Cone}(\alpha_i) & \xrightarrow{\alpha_i\text{Cone}(\alpha_i)} & \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle\text{Cone}(\alpha_i) & \longrightarrow & \text{Cone}(\alpha_i)\text{Cone}(\alpha_i)
\end{array}$$

By definition, $\Pi_i \circ \Pi_{i+1} \circ \Pi_i = \text{Cone}(F \rightarrow F\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle)$. We repeat the computation done for $\Pi_{i+1} \circ \Pi_i \circ \Pi_{i+1}$, and find this is isomorphic to $\text{Cone}(F \rightarrow \text{Cone}(\tilde{\epsilon}_{s_{i+1}}L\tilde{Z}^{s_{i+1}}\langle 1 \rangle\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle \rightarrow \tilde{\epsilon}_{t_i}L\tilde{Z}^{t_i}[1]\langle 3 \rangle))$.

Now we just have to verify that the maps in both cones are the same. Consider the commutative diagram

$$\begin{array}{ccc}
F & \longrightarrow & F\tilde{\epsilon}_iL\tilde{Z}_i\langle 1 \rangle \\
\downarrow & & \downarrow \\
\tilde{\epsilon}_{i+1}L\tilde{Z}_{i+1}\langle 1 \rangle F & \longrightarrow & \tilde{\epsilon}_{i+1}L\tilde{Z}_{i+1}\langle 1 \rangle F\tilde{\epsilon}_iL\tilde{Z}_i\langle 1 \rangle
\end{array}$$

Since $\tilde{\epsilon}_{i+1}L\tilde{Z}_{i+1}\langle 1 \rangle F \rightarrow \tilde{\epsilon}_{i+1}L\tilde{Z}_{i+1}\langle 1 \rangle F\tilde{\epsilon}_iL\tilde{Z}_i\langle 1 \rangle$ and $F\tilde{\epsilon}_iL\tilde{Z}_i\langle 1 \rangle \rightarrow \tilde{\epsilon}_{i+1}L\tilde{Z}_{i+1}\langle 1 \rangle F\tilde{\epsilon}_iL\tilde{Z}_i\langle 1 \rangle$ are split inclusions by lemma 55, both maps in the cone are the same. □

Corollary 20. *There is an isomorphism*

$$\Omega_i\Omega_{i+1}\Omega_i \cong \Omega_{i+1}\Omega_i\Omega_{i+1}.$$

Proof. This follows immediately from the proposition by taking the adjoint of both sides. □

6.6. Relation 6.

Proposition 21. *There is an isomorphism $\cap_{i+1,+ , r+2} \circ \Omega_i \circ \cup_{i+1,+ , r} \cong \text{Id}$.*

Proof. The left hand sides is

$$\begin{aligned}
& L\tilde{Z}_{q_{i+1}}^{p_{i+1}}\text{Cone}(\tilde{\epsilon}_{s_i}[-1]L\tilde{Z}^{s_i} \rightarrow \text{Id}[1]\langle 1 \rangle)\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)][-k]\langle -k \rangle \cong \\
& \text{Cone}(L\tilde{Z}_{q_{i+1}}^{p_{i+1}}\tilde{\epsilon}_{s_i}[-1]L\tilde{Z}^{s_i}\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)] \rightarrow L\tilde{Z}_{q_{i+1}}^{p_{i+1}}\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)+1]\langle 1 \rangle)[-k]\langle -k \rangle).
\end{aligned}$$

Now apply proposition 4.19 to the left hand side and corollary 4.5 to the right to get

$$\text{Cone}(\oplus_{j=0}^{k-2} \text{Id}[k-2-2j]\langle k-2-2j \rangle \rightarrow \oplus_{j=0}^{k-1} \text{Id}[k-2j]\langle k-2j \rangle)[-k]\langle -k \rangle \cong \text{Id}.$$

□

Proposition 22. *There is an isomorphism $\cap_{i+1,+ , r+2} \circ \Pi_i \circ \cup_{i+1,+ , r} \cong \text{Id}$.*

Proof. The left hand side is

$$\begin{aligned}
& L\tilde{Z}_{q_{i+1}}^{p_{i+1}}\text{Cone}(\text{Id}\langle -1 \rangle \rightarrow \tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\langle 1 \rangle)\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)][k-2]\langle k \rangle \cong \\
& \text{Cone}(L\tilde{Z}_{q_{i+1}}^{p_{i+1}}\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)]\langle -1 \rangle \rightarrow L\tilde{Z}_{q_{i+1}}^{p_{i+1}}\tilde{\epsilon}_{s_i}L\tilde{Z}^{s_i}\tilde{\epsilon}_{p_{i+1}}^{q_{i+1}}[-(k-1)][k-2]\langle k \rangle).
\end{aligned}$$

Now apply corollary 10 to the left side and proposition 12 to the right to get

$$\text{Cone}(\oplus_{j=0}^{k-1} \text{Id}[k-1-2j]\langle k-2-2j \rangle \rightarrow \oplus_{j=0}^{k-2} \text{Id}[k-1-2j]\langle k-2-2j \rangle)[k-2]\langle k \rangle \cong \text{Id}.$$

□

6.7. Relation 7. For purposes of this subsection, we have the following abbreviations for the relevant subalgebras.

- Definition 18.**
- (1) $\mu = \mathfrak{p}_i + \mathfrak{q}_{i+k}$.
 - (2) $\nu = \mathfrak{q}_{i-1} + \mathfrak{q}_{i+k}$.
 - (3) $\gamma = \mathfrak{q}_{i-1} + \mathfrak{s}_{i+k-1} + \mathfrak{q}_{i+k}$.
 - (4) $\delta = \mathfrak{q}_{i-1} + \mathfrak{p}_{i+k}$.

We let $\alpha: \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \rightarrow \text{Id}[2]\langle 1 \rangle$ and $\beta: \text{Id} \rightarrow \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \langle 1 \rangle$.

Proposition 23. *There are functorial isomorphisms:*

- (1)
$$\cap_{i,-,r+2} \circ \Omega_{i+k-1} \circ \cup_{i+2,+,r} \circ \cap_{i+2,+,r+2} \circ \Pi_{i+k-1} \circ \cup_{i,-,r} \cong \text{Id}$$
- (2)
$$\cap_{i,-,r+2} \circ \Pi_{i+k-1} \circ \cup_{i+2,+,r} \circ \cap_{i+2,+,r+2} \circ \Omega_{i+k-1} \circ \cup_{i,-,r} \cong \text{Id}.$$

Proof. (1) The left hand side is the functor given below shifted by $[-3]\langle -1 \rangle$.

$$L\tilde{Z}_\nu^\mu \text{Cone}(\text{Id} \xrightarrow{\beta} \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \langle 1 \rangle) \tilde{\epsilon}_\delta^\nu[-(k-1)] L\tilde{Z}_\nu^\delta \text{Cone}(\tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \xrightarrow{\alpha} \text{Id}[2]\langle 1 \rangle) \tilde{\epsilon}_\mu^\nu[-(k-1)].$$

This expression gives rise to the following commutative diagram.

$$\begin{array}{ccccc}
L\tilde{Z}_\nu^\mu \text{Cone}(\beta) \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \tilde{\epsilon}_\mu^\nu[-2(k-1)] & \longrightarrow & L\tilde{Z}_\nu^\mu \text{Cone}(\beta) \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \tilde{\epsilon}_\mu^\nu[-2k+4]\langle 1 \rangle & \longrightarrow & L\tilde{Z}_\nu^\mu \text{Cone}(\beta) \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \text{Cone}(\alpha) \tilde{\epsilon}_\mu^\nu[-2(k-1)] \\
\uparrow & & \uparrow & & \uparrow \\
L\tilde{Z}_\nu^\mu \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \tilde{\epsilon}_\mu^\nu[-2(k-1)]\langle 1 \rangle & \longrightarrow & L\tilde{Z}_\nu^\mu \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \tilde{\epsilon}_\mu^\nu[-2k+4]\langle 2 \rangle & \longrightarrow & L\tilde{Z}_\nu^\mu \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \text{Cone}(\alpha) \tilde{\epsilon}_\mu^\nu[-2(k-1)]\langle 1 \rangle \\
\uparrow & & \uparrow & & \uparrow \\
L\tilde{Z}_\nu^\mu \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \tilde{\epsilon}_\gamma^\nu L\tilde{Z}_\nu^\gamma \tilde{\epsilon}_\mu^\nu[-2(k-1)] & \longrightarrow & L\tilde{Z}_\nu^\mu \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \tilde{\epsilon}_\mu^\nu[-2k+4]\langle 1 \rangle & \longrightarrow & L\tilde{Z}_\nu^\mu \tilde{\epsilon}_\delta^\nu L\tilde{Z}_\nu^\delta \text{Cone}(\alpha) \tilde{\epsilon}_\mu^\nu[-2(k-1)] \\
\text{Using the isomorphisms constructed earlier, the above diagram is the same as} & & & & \\
\tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[-2k+4]\langle -k+2 \rangle \oplus \text{Id}[2]\langle 1 \rangle & \longrightarrow & \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[-2k+5]\langle -k+2 \rangle & \longrightarrow & X \\
\uparrow & & \uparrow & & \uparrow \\
\oplus_{j=0}^{k-3} \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[-2j]\langle k-2-2j \rangle \oplus \text{Id}[2]\langle 1 \rangle & \longrightarrow & \oplus_{j=0}^{k-2} \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[2-2j]\langle k-2j \rangle & \longrightarrow & \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[2]\langle k \rangle \oplus \text{Id} \\
\uparrow & & \uparrow & & \uparrow \\
\oplus_{j=0}^{k-2} \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[-2j]\langle k-2-2j \rangle & \longrightarrow & \oplus_{j=0}^{k-1} \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[2-2j]\langle k-2j \rangle & \longrightarrow & \tilde{\epsilon}_{\mathfrak{p}_i}^{\mathfrak{q}_i} L\tilde{Z}_{\mathfrak{q}_i}^{\mathfrak{p}_i}[2]\langle k \rangle
\end{array}$$

Therefore $X \cong \text{Id}[3]\langle 1 \rangle$ and so the composition of the functors with the its built in shifts is the identity.

- (2) The proof is the same. One just has to reflect the commutative diagrams in the diagonal going from the bottom left to the top right.

□

6.8. Relation 8.

Proposition 24. *There are the following functorial isomorphisms:*

- (1)
$$\cap_{i,+,r+2} \circ \Omega_{i+k-3} \circ \cup_{i-2,-,r} \circ \cap_{i-2,-,r+2} \circ \Pi_{i+k-3} \circ \cup_{i,+,r} \cong \text{Id}$$
- (2)
$$\cap_{i,+,r+2} \circ \Omega_{i+k-3} \circ \cup_{i-2,-,r} \circ \cap_{i-2,-,r+2} \circ \Pi_{i+k-3} \circ \cup_{i,+,r} \cong \text{Id}.$$

Proof. One verifies these isomorphisms as in the proofs of relation seven. The only differences are the subscripts of the parabolic subalgebras. □

6.9. Skein Relations. The images of the functors assigned to the crossings in the Grothendieck group satisfy the skein relations for the HOMFLYPT polynomial. The following propositions follows directly from the definitions of the functors involved.

Theorem 8. *The following relations are satisfied in the Grothendieck group:*

- (1) $[\Pi_i] = (-1)^k q^k ([\tilde{\epsilon}_{s_i} L \tilde{Z}^{s_i}] - q^{-1} [\text{Id}]).$
- (2) $[\Omega_i] = (-1)^k q^{-k} (-q [\text{Id}] + [\tilde{\epsilon}_{s_i} L \tilde{Z}^{s_i}])$
- (3) $q^k [\Omega_i] - q^{-k} [\Pi_i] = (-1)^{k-1} (q^1 - q^{-1}) [\text{Id}]$

REFERENCES

- [AndStr] H. Andersen, C. Stroppel, Twisting functors on $\mathcal{O}$, *Represent. Theory* **7**, (2003), 681–699 (electronic).
- [Ark] S. Arkhipov, Algebraic construction of contragradient quasi-Verma modules in positive characteristic. Representation theory of algebraic groups and quantum groups, *Adv. Stud. Pure Math.*, **40**, Math. Soc. Japan, Tokyo, (2004), 27–68.
- [Bac] E. Backelin, Koszul duality for parabolic and singular category $\mathcal{O}$, *Representation Theory*, (1999) 139–152.
- [Bass] H. Bass, Algebraic K-theory, Benjamin 1968.
- [BFK] J. Bernstein, I. Frenkel, M. Khovanov, A Categorification of the Temperley-Lieb Algebra and Schur Quotients of $U(\mathfrak{sl}_2)$ via Projective and Zuckerman Functors, *Selecta Mathematica*, 1999.
- [BG] J. N. Bernstein, S. I. Gelfand, Tensor products of finite and infinite dimensional representations of semisimple Lie algebras, *Compositio Math.* **41**, (1980), 245–285.
- [BGG] J.N. Bernstein, I.M. Gelfand and S.I. Gelfand, Category of $\mathfrak{g}$ -modules, *Functional Anal. and Appl.* **10**, 2 (1976) 87–92.
- [BGS] A. Beilinson, V. Ginzburg and W. Soergel, Koszul Duality patterns in representation theory, *Journal of the AMS* **9** (1996), 473–527.
- [Br] J. Brundan, Symmetric functions, parabolic category $\mathcal{O}$ and the Springer fiber, math.RT/0608235.
- [ES] T. Enright, B. Shelton, Categories of highest weight modules: applications to classical Hermitian symmetric pairs, *Memoirs of the American Mathematical Society* **367**, (1987), 1–94.
- [EW] T. Enright, N. Wallach, Notes on Homological Algebra and Representations of Lie Algebras, *Duke Math. J.* **47**, 1 (1980), 1–15.
- [FH] W. Fulton, J. Harris, Representation Theory A First Course, Springer-Verlag, 1991.
- [FK] I. Frenkel, M. Khovanov, Canonical Bases In Tensor Products And Graphical Calculus For $U_q(\mathfrak{sl}_2)$, *Duke Math J.* **87**, 3 (1997), 409–480.
- [FKS] I. Frenkel, M. Khovanov, C. Stroppel, A Categorification of Finite-Dimensional Irreducible Representations of Quantum $\mathfrak{sl}_2$ and Their Tensor Products, math.QA/0511467.
- [GM] S.I. Gelfand, Yu. I. Manin, Methods of Homological Algebra, Springer-Verlag, 1996.
- [J] J.C. Jantzen, Einhüllende Algebren halbeinfacher Lie-Algebren. Ergebnisse der Mathematik und ihrer Grenzgebiete, Berlin **3**, 1983.
- [Jac] M. Jacobsson, An invariant of link cobordisms from Khovanov homology, *Algebr. Geom. Topol.*, **4** (2004) 1211–1251.
- [Jo1] A. Joseph, The Enright Functor on the Bernstein-Bernstein-Gelfand Category $\mathcal{O}$, *Invent. Math.*, **67**, (1982), 423–445.
- [Jo2] A. Joseph, Completion Functors in the $\mathcal{O}$ Category, In: Non Commutative Harmonic Analysis and Lie Groups, Springer Lecture Notes, **1020**, 1983, 80–106.
- [Ka] C. Kassel, Quantum Groups, Springer-Verlag, 1995.
- [KasSch] M. Kashiwara, P. Schapira, Sheaves on Manifolds, Springer-Verlag, 1994.
- [Khom] O. Khomenko, Categories with Projective Functors, *Proc. London Math. Soc.*, **90**, 3, 711–737.
- [Khov] M. Khovanov, A categorification of the Jones polynomial, *Duke Math. J.*, 101(3): 359–426, 2000.
- [Khov2] M. Khovanov, Crossingless matchings and the cohomology of (n,n) Springer varieties, *Communications in Contemporary Math.*, **6** (2004) no.2, 561–577.
- [Khov3] M. Khovanov, An invariant of tangle cobordisms, *Trans. Amer. Math. Soc.*, **358** (2006), 315–327.
- [Khov4] M. Khovanov, Triply-graded link homology and Hochschild homology of Soergel bimodules, math.GT/0510265.
- [Khov5] M. Khovanov, A functor-valued invariant of tangles, *Algebr. Geom. Topol.*, **2** (2002), 665–741.
- [KhoMaz] O. Khomenko, V. Mazorchuk, On Arkhipov’s and Enright’s functors, *Math. Z.* **249** 2, (2005), 357–386.
- [KV] A. Knapp, D. Vogan, Cohomological Induction and Unitary Representations, Princeton University Press, 1995.
- [KR] M. Khovanov, L. Rozansky, Matrix Factorization and Link Homology, math.QA 0401268.
- [Le] J. Lepowsky, A generalization of the Bernstein-Gelfand-Gelfand resolution, *J. Algebra*, **49**, 2 (1977), 496–511.
- [Mac] S. Mac Lane, Categories for the working mathematician, Springer-Verlag, 1998.
- [MazStr1] V. Mazorchuk, C. Stroppel, On Functors Associated to a Simple Root, math.RT/0410339.
- [MazStr2] V. Mazorchuk, C. Stroppel, Translation and shuffling of projectively presentable modules and a categorification of a parabolic Hecke module, *Trans. Amer. Math. Soc.*, **357** (2005), no. 7, 2939–2973.
- [MOS] V. Mazorchuk, S. Ovsienko, C. Stroppel, Quadratic duals, Koszul dual functors, and applications, math.RT/0603475.
- [MOY] H. Murakami, T. Ohtsuki, S. Yamada, HOMFLY Polynomial via an Invariant of Colored Plane Graphs, *Enseign. Math* (2) **44** (1998), no. 3–4, 325–360.
- [MP] D. Milicic, P. Pandzic, On degeneration of the spectral sequence for the composition of Zuckerman functors, *Glas. Mat. Ser. III* **32**(52), 2(1997), 179–199.
- [Ro] A. Rocha-Caridi, Splitting criteria for $\mathfrak{g}$ -modules induced from a parabolic and Bernstein-Gelfand-Gelfand resolution of a finite-dimensional, irreducible $\mathfrak{g}$ -module, *Trans. AMS* **262**, 2 (1980), 335–366.

- [Rh] S. Ryom-Hansen, Koszul duality of translation and Zuckerman functors, *J. Lie Theory*, **14** (2004), no. 1, 151-163.
- [Sch] P. Schapira, Categories and homological algebra, 2006.
- [Sh] B. Shelton, Extensions Between Generalized Verma Modules: The Hermitian Symmetric Case, *Math. Z.* **197**, (1988), 305-318.
- [Soe] W. Soergel, Kategorie $\mathcal{O}$, perverse Garben und Moduln über den Koinvarianten zur Weylgruppe, *J. Amer. Math. Soc.*, **3** (1990), no. 2, 421-445.
- [Soe2] W. Soergel, Gradings on representation categories, Proceedings of the International Congress of Mathematicians, Zurich, Switzerland, 1994.
- [Str1] C. Stroppel, Category $\mathcal{O}$: Gradings and Translation Functors., *J. Algebra* **268**, (2003), no. 1, 301-326.
- [Str2] C. Stroppel, Categorification of the Temperley-Lieb category, tangles, and cobordisms via projective functors, *Duke Math. J.* **126**, (2005), no. 3, 547-596.
- [Str3] C. Stroppel, Category $\mathcal{O}$: Quivers and endomorphism rings of projectives, *Representation Theory* **7**, 2003.
- [Str4] C. Stroppel, TQFT with corners and tilting functors in the Kac-Moody case, RT/0605103.
- [Str5] C. Stroppel, Perverse sheaves on Grassmannians, Springer fibres and Khovanov homology, math.RT/0608234.
- [Z] G. Zuckerman, Tensor products of finite and infinite dimensional representations of semisimple Lie groups, *Ann. Math.*, **106** (1977), no. 2, 295-308.